

**UNIVERSIDADE FEDERAL DE ALAGOAS
INSTITUTO DE CIÊNCIAS BIOLÓGICAS E DA SAÚDE
Programa de Pós-Graduação em Diversidade Biológica e Conservação nos
Trópicos**

CIRO RAMON FÉLIX DOS SANTOS SILVA

**Microbiologia da filosfera: Explorando a biodiversidade, ecologia e taxonomia no
sistema leveduras-bromélias em escala global e no semiárido brasileiro**

**MACEIÓ - ALAGOAS
Março/2023**

CIRO RAMON FÉLIX DOS SANTOS SILVA

Microbiologia da filosfera: Explorando a biodiversidade, ecologia e taxonomia no sistema leveduras-bromélias em escala global e no semiárido brasileiro

Tese apresentada ao Programa de Pós-Graduação em Diversidade Biológica e Conservação nos Trópicos, Instituto de Ciências Biológicas e da Saúde, Universidade Federal de Alagoas, como requisito para obtenção do título de Doutor em CIÉNCIAS BIOLÓGICAS, área de concentração em Conservação da Biodiversidade Tropical.

Orientadora: Prof^a. Dr^a. Melissa Fontes Landell

MACEIÓ – ALAGOAS
Março/2023

**Catalogação na Fonte
Universidade Federal de Alagoas
Biblioteca Central
Divisão de Tratamento Técnico**

Bibliotecário: Marcelino de Carvalho Freitas Neto – CRB-4 - 1767

S586m Silva, Ciro Ramon Félix dos Santos.

Microbiologia da filosfera : explorando a biodiversidade, ecologia e taxonomia no sistema leveduras-bromélias em escala global e no semiárido brasileiro / Ciro Ramon Félix dos Santos Silva. – 2023.

206 f. : il.

Orientadora: Melissa Fontes Landell.

Tese (Doutorado em Ciências Biológicas) – Universidade Federal de Alagoas. Instituto de Ciências Biológicas e da Saúde. Programa de Pós-Graduação em Diversidade Biológica e Conservação nos Trópicos. Maceió, 2023.

Bibliografia: f. 171-179.

Anexos: f. 180-206.

1. Semiárido. 2. Secas. 3. Pulso. 4. Filosfera. 5. Aridez. I. Título.

CDU: 57.063.3

Dedico esse trabalho a vida e memória do meu
amado pai, Cícero José da Silva.

AGRADECIMENTOS

Agradeço a minha família, o meu tesouro mais precioso, sem eles, nada até aqui seria possível.

Agradeço a vida do meu pai, Cícero, que infelizmente não pode ver seu filho pleitear o doutorado, mas que sempre foi o melhor exemplo de homem, pai e ser humano que eu pude ter. Agradeço por tudo que ele me ensinou e pela sorte e privilégio de ter tido um pai tão maravilhoso. A memória e o legado do meu pai viverão comigo.

Agradeço a minha mãe, Luzia, onde eu sempre encontrei um porto seguro. A mulher que lutou para que eu tivesse a possibilidade de estudar. Agradeço por ela ter me criado cercado de amor e por todos os sacrifícios que ela fez por mim e meus irmãos.

Agradeço aos meus irmãos que tanto amo, Eraldo e Luany, que compartilham comigo um laço tão especial e que são para mim motivo de orgulho e, eventualmente, de dor de cabeça.

Sou grato à minha avó, Maria (Dona Zezinha), que sempre me deu acalento e carinho no melhor estilo ‘voinha’.

Agradeço a minha amiga, companheira, namorada e esposa, Elizabeth, meu grande amor, que foi meu lastro esses quatro anos. Agradeço porque esse trabalho não seria possível sem ela. Agradeço por ela ter estado ao meu lado, me cuidando e incentivando. Agradeço a sorte do meu sorriso de tê-la como motivo.

Agradeço a minha orientadora, Melissa Landell, que me ensinou tanto e que ainda me ensina, mesmo após quase dez anos. Agradeço a essa mentora e orientadora dedicada. Agradeço pelas inúmeras oportunidades e pela paciência durante esse tempo.

Agradeço a minha sogra, Ana, aos meus cunhados, Dayse e Orlando. Todos foram tão gentis e me acolheram tão bem durante esses anos.

Agradeço aos meus irmãos de coração João, Stephannie, Gustavo e Maurício. Que me ajudaram nessa e em outras tantas empreitadas da minha vida.

Agradeço aos meus amigos e colegas de laboratório Aldeci, Vitória, Bruno, Victor e Júlio. Que me ajudaram diretamente na construção desse trabalho e compartilharam o dia a dia comigo nos últimos anos.

Agradeço profundamente a todos os meus professores e mestres que contribuíram na minha formação, desde os níveis mais básicos até a pós-graduação. Cada um deles merece toda a minha gratidão, respeito, valorização e consideração.

Agradeço a Universidade Federal de Alagoas (UFAL), uma universidade pública, gratuita e de qualidade, onde tive o privilégio de ser formado desde a graduação. Que apesar das constantes investidas de um governo anticiência, segue firme em seu papel de servir a sociedade alagoana. Agradeço a UFAL e aos governos que apoiaram a educação transformadora, de qualidade e acessível a todos. Agradeço porque eu, filho de uma professora e de um motorista, tive a oportunidade de buscar o mais alto título acadêmico graças a essa instituição. Agradeço e me orgulho por essa instituição buscar incluir o povo preto, pobre, indígena, LGBTQIA+, as pessoas portadoras de deficiência e tantos outros que são invisibilizados.

Agradeço aos meus colegas e professores do PPG-DIBICT, que compartilharam essa jornada comigo.

Agradeço aos membros da banca de avaliação continuada (BAC): Patrícia Valente, Jadson Bezerra e Gilberto Justino, por todo o aprendizado e contribuição ao longo do meu doutoramento.

Agradeço ao Sr. Alberto Nepomuceno Agra Jr. e ao seu finado pai Alberto Nepomuceno Agra pelo pioneirismo e dedicação em conservar a Caatinga alagoana, e a gentileza de autorizar as coletas na RPPN Tocaia.

Agradeço ao Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), a Fundação de Amparo à Pesquisa do Estado de Alagoas (FAPEAL) e a Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (Capes) pela concessão da bolsa e financiamento que permitiu o andamento deste trabalho.

O cientista planta tâmaras, a ciência as colhe.
(Reinterpretação autoral do ditado árabe)

RESUMO

Regiões secas (*Drylands*) são áreas com índices de aridez $\leq 0,65$. Nesse tipo de ambiente, as chuvas são escassas e ocorrem em pulsos (eventos raros de superdisponibilidade de recursos). Durante a seca, a demanda nutricional e metabólica é diminuída. Desta maneira, quanto maior o tempo de seca, maior o reservatório nutricional acumulado no ambiente. O tamanho do reservatório influencia a intensidade da resposta ao pulso de água, ou seja, existe uma ‘memória’ ambiental do evento de chuva anterior. Outro exemplo de ecossistema com frequente déficit de água é a filosfera (parte aérea das plantas). Esse ambiente intermedia a relação da planta com ambiente e é uma fonte megadiversa de microrganismos. A filosfera de bromélias, plantas tipicamente neotropicais, abrigam diversos grupos microbianos, incluindo leveduras. Entretanto, o conhecimento sobre leveduras em bromélias é disperso e ainda incipiente. Os objetivos do presente estudo foram: 1) Compilar, sintetizar e compreender a extensão do conhecimento sobre leveduras da filosfera de bromélias em escala global. 2) Verificar o efeito da chuva e da memória ambiental sobre a diversidade e estrutura taxonômica, filogenética e funcional das leveduras da filosfera de bromélias na Caatinga. 3) Caracterizar a possível nova espécie *Carlosrosaea* xxxxxxxx. Para tanto, a literatura sobre leveduras em bromélias foi sistematicamente revisada. Ademais, leveduras foram isoladas regularmente durante dois anos a partir de bromélias na Reserva Particular do Patrimônio Natural (RPPN) Tocaia, no município de Santana do Ipanema, Sertão alagoano. Na produção acadêmica das últimas três décadas foi encontrado registro de mais de 180 espécies de leveduras na filosfera de bromélias, distribuídas em quatro compartimentos: flores, frutos, folhas e tanques. Cerca de 70% das espécies ocorreram exclusivamente em um único compartimento e apenas 2% foram comuns a todos. Mais de 20 novas espécies de leveduras foram descritas em bromélias nesse período e ao menos 50 espécies mostraram algum potencial biotecnológico. Quanto ao efeito das chuvas, a diversidade-alfa não diferiu significativamente entre períodos sazonais. Entretanto, a composição taxonômica divergiu em mais de 70%, embora tenha se mantido funcionalmente estável. A chuva, seca e a memória ambiental tiveram pouca influência nas métricas de diversidade. No entanto, tiveram relação com a estrutura da comunidade. A análise das regiões ITS e D1/D2 do gene 26S rRNA de oito isolados da Caatinga indicou uma possível nova espécie de levedura com afinidade ao gênero *Carlosrosaea*. Diante disso, propõe-se uma nova espécie de *Carlosrosaea* para agrupar os isolados citados. Os resultados trazem informações sobre a diversidade e dinâmica das leveduras da filosfera de bromélias e como ela se relaciona com os pulsos de chuva, seca, sazonalidade e efeito de memória. Compreendendo o efeito da chuva e da seca na microbiota da filosfera, principalmente em regiões secas, pode estimar como a comunidade microbiana das folhas será afetada pelas mudanças climáticas que alteram os ciclos seco-úmido e, consequentemente, como isso se relaciona com o hospedeiro.

Palavras-chave: Caatinga, Regiões Secas, pulsos, filoplano, aridez.

ABSTRACT

Drylands are areas characterized by aridity indexes ≤ 0.65 . In this type of environment, rainfall is decisive for biological dynamics and occurs in pulses (rare events of overavailability of resources). During drought, nutritional and metabolic demand is decreased. Thus, the longer the dry period, the greater the nutritional reservoir accumulated in the environment. Reservoir size influences the strength of the response to the water pulse. That is, there is an environmental 'memory' of the previous rainfall event. Another example of an ecosystem with a frequent water deficit is the phyllosphere (the aerial part of plants). This environment is one of the largest sources of microorganisms on the planet. The phyllosphere of bromeliads, typically neotropical plants, harbor several microbial groups, including yeasts. However, knowledge about yeasts in bromeliads is scattered and still incipient. In this study our objectives were: 1) Compile, synthesize and understand the extent of knowledge about yeasts from the bromeliad phyllosphere. 2) To verify the effect of rain and environmental memory on the diversity and taxonomic, phylogenetic and functional structure of yeasts from the phyllosphere of bromeliads in the Caatinga. 3) Propose the description of the new species *Carlosrosaea* sp. nov. To this end, the literature on yeasts in bromeliads was systematically reviewed. Furthermore, yeasts were regularly isolated for two years from bromeliads in the Private Natural Heritage Reserve (RPPN) Tocaia, in the municipality of Santana do Ipanema, Alagoas. In the academic production of the last three decades, records of more than 180 species of yeasts were found in the phyllosphere of bromeliads, distributed in four compartments: flowers, fruits, leaves and tanks. About 70% of the species occurred exclusively in a single compartment and only 2% were common to all. More than 20 new yeast species were described in bromeliads during this period and at least 50 species showed some biotechnological potential. As for the effect of rainfall, alpha-diversity did not differ significantly between seasonal periods. However, the taxonomic composition differed by more than 70%, although it was functionally stable. Rain, drought and environmental memory had little influence on diversity metrics. However, they showed a relationship with the structure of the community. The analysis of the ITS and D1/D2 regions of the 26S rRNA gene of eight Caatinga isolates indicated a possible new yeast species with affinity to the genus *Carlosrosaea*. Therefore, a new species of *Carlosrosaea* is proposed to group the cited isolates. Our results bring information about the diversity and dynamics of yeasts in the bromeliad phyllosphere and how it relates to rainfall pulses, drought, seasonality and memory effect. Furthermore, we aggregate information about the tolerance of the phyllosphere yeast community to seasonal changes, from a taxonomic, phylogenetic and functional point of view. By understanding the effect of rain and drought on the phyllosphere microbiota, especially in dry regions, we can think about how the microbial community of the leaves will be affected by climate changes that alter the dry-wet cycles and, consequently, how this relates to the host.

Keywords: Caatinga, Drylands, pulses, phyllosphere, aridity.

SUMÁRIO

1.	APRESENTAÇÃO	13
2.	REVISÃO DA LITERATURA	16
2.1.	Regiões Secas, Florestas Sazonalmente Secas e Caatinga.....	16
2.2.	Ecologia de regiões secas.....	21
2.3.	Micro-organismos associados a plantas.....	26
2.4.	A relação da microbiota da filosfera com a água.....	31
2.5.	Leveduras.....	34
2.6.	Bromélias	39
3.	MÉTODOS	42
3.1.	Caracterização da área de estudo	42
3.2.	Coleta do material	42
3.3.	Processamento das amostras	43
3.4.	Obtenção e armazenamento dos isolados	44
3.5.	Análise de traços funcionais.....	44
3.5.1.	Fermentação de glicose	44
3.5.2.	Assimilação de fontes de carbono.....	45
3.5.3.	Produção de hidrolases extracelulares.....	45
3.5.4.	Produção de biossurfactantes/bioemulsificantes.....	47
3.6.	Identificação molecular dos isolados	47
3.6.1.	Extração de DNA genômico	47
3.6.2.	PCR, sequenciamento e análise das sequências	48
3.7.	Caracterização das possíveis espécies novas	49
3.7.1.	Assimilação de fontes de carbono e nitrogênio	49
3.7.2.	Formação de compostos amiloïdes extracelulares.....	50
3.7.4.	Testes de produção de urease e reação ao Diazônio Azul B (DBB)	51
3.7.5.	Testes de tolerância	51
3.8.	Caracterização micromorfológica	52
4.	CAPÍTULO 1 - Different plant compartments, different yeasts: the example of the bromeliad phyllosphere.....	53

5.	CAPÍTULO 2 - Diversity, rainfall pulses and memory effect: A look at yeast-plant system from Brazilian Tropical Semiarid Dryland	120
6.	CAPÍTULO 3 - <i>Carlosrosaea</i> xxxxxxx sp. nov., a new tremellomycetes yeast from Brazilian Seasonally Dry Tropical Forest (Caatinga).....	153
7.	DISCUSSÃO GERAL.....	167
8.	CONSIDERAÇÕES FINAIS	169
9.	REFERÊNCIAS.....	171
	ANEXOS	180

1. APRESENTAÇÃO

Regiões secas como a Caatinga cobrem mais de 40% da superfície continental no planeta. Essas regiões são frequentemente secundarizadas pelo Estado e pela ciência. Não é incomum que muitos representantes dessas regiões estejam subamostrados, sobretudo quando comparados aos seus equivalentes úmidos. A Caatinga não foge à regra e frequentemente recebe menos atenção comparada a seus pares úmidos como a Mata Atlântica e a Amazônia. Esse descaso tem componentes sociais e culturais, seja pelo ecossistema ocorrer no Nordeste do país, uma região historicamente desprivilegiada e negligenciada, ou pelo mito que foi construído no senso comum que a Caatinga é um ambiente com pouca biodiversidade.

Na maior parte do ano este ecossistema tem um aspecto seco, com troncos retorcidos e plantas sem folhas. Esse cenário que, durante a maior parte do tempo, diverge das paisagens verdes e exuberantes que habitam o imaginário popular quando se pensa em biodiversidade foi um lastro importante para a construção do mito entorno da Caatinga. Por outro lado, nas últimas décadas, um grande esforço tem sido empregado em compreender a Caatinga e sua biodiversidade. Regiões secas proveem diversos serviços ecossistêmicos fundamentais como a manutenção dos níveis globais de carbono atmosférico e do clima mundial. Em um cenário de emergência climática como o atual, esses serviços ecossistêmicos se tornam centrais no debate científico e no interesse público.

A disponibilidade de água, sobretudo em regiões secas, afeta todos os tipos de organismos, em todos os níveis de organização ecológica. Tanto macro quanto micro-organismos são afetados, porém, geralmente os organismos microscópicos em ambiente natural (não-clínico) são menos estudados que os macroscópicos. Micro-organismos prestam várias funções relevantes como fixação de carbono e nitrogênio, ciclagem de nutrientes e diversas outras funções que se dão no âmbito das associações mutualísticas com outros grupos, como a manutenção da saúde do hospedeiro e o sucesso reprodutivo. A insipiência dos estudos sobre micro-organismos gera uma

grande lacuna de conhecimento que diminui a compreensão do funcionamento do ecossistema de processos e funções importantes desempenhadas pela microbiota.

Diversos grupos microbianos têm sido encontrados em ambientes áridos, muitos deles com adaptações para tolerar forte radiação solar, estresse oxidativo e déficit hídrico. Entre os grupos estão bactérias, arqueas, microalgas, fungos filamentosos e leveduras. Além do solo, as plantas são substratos importantes para os micro-organismos, vários grupos são capazes de interagir positivamente com plantas produzindo hormônios vegetais, além de ter atividade antagônica a patógenos. A grande ocorrência de micro-organismos em plantas pode ser exemplificada com os fungos. Esse grupo é um dos mais diversos do mundo. Estima-se que existam entre 2,2 e 3,8 milhões de espécies, porém, conhece-se apenas cerca de 5% das espécies. Nos grandes bancos de dados micológicos como o *Mycobank*, a maioria dos registros de fungos concentra-se em substratos vegetais, sobretudo em folhas.

Fungos podem ser subdivididos segundo diversos critérios, um deles separa fungos em filamentosos e leveduras. Leveduras consistem em um grupo formado por várias linhagens de fungos que convergiram para um estado majoritariamente leveduriforme, ou seja, unicelular. Esse grupo de fungos unicelulares está distribuído em diversos ambientes e substratos como animais, plantas, solo e água. Possuem importância biotecnológica (ex. na produção de alimentos e bebidas fermentadas), clínica (causando fungemias como as espécies *Candida albicans* e *Cryptococcus neoformans*) e ambiental (sendo responsáveis por funções ecossistêmicas fundamentais para a manutenção da vida no planeta como a ciclagem de nutrientes).

Plantas tropicais são uma das maiores fonte de micro-organismos. Além disso, são um dos mais importantes, talvez o maior, substrato para descoberta de novas espécies de fungos. Entre os grupos de plantas tropicais mais diversos e endêmicos do Brasil está a família Bromeliaceae. Essa família se distribui entre ambientes úmidos como a Mata Atlântica até regiões semiáridas como a Caatinga. Bromélias são consideradas mosaicos complexos e diversos para leveduras, além de um modelo ecológico importante para regiões tropicais.

Diante da insipiência que ainda persiste nos estudos sobre microbiota ambiental, principalmente de leveduras, em regiões secas, este estudo objetivou: 1) Sintetizar o conhecimento sobre leveduras em bromélias, elucidando padrões e processos ecológicos, lacunas do conhecimento e potencial biotecnológico; 2) Avaliar a relação dos pulsos de chuvas e da sazonalidade nos padrões de diversidade funcional, taxonômica e filogenética de leveduras associadas à bromélias de uma região de Caatinga, no estado de Alagoas; 3) Caracterizar molecular e fisiologicamente uma possível espécie nova de leveduras associadas à bromélias, contribuindo para o conhecimento sobre a biodiversidade do bioma Caatinga.

2. REVISÃO DA LITERATURA


2.1. Regiões Secas, Florestas Sazonalmente Secas e Caatinga

Regiões secas não-polares (*drylands*) cobrem 41% da superfície continental do planeta, o equivalente a aproximadamente 6 bilhões de hectares, e são consideradas ecossistemas frágeis, com biodiversidade única. Esses ambientes fornecem importantes serviços ecossistêmicos no ciclo global do carbono, do nitrogênio, da água e na regulação do clima (HUANG et al., 2016; MAESTRE et al., 2016, 2021). Além disso, essas regiões abrigam 38% da população humana, 35% da biodiversidade global e 20% da diversidade de plantas (MAESTRE et al., 2021). De forma simples (e incompleta), pode-se dizer que regiões secas são áreas com baixa precipitação anual.

Existem vários critérios que podem ser empregados na classificação das regiões secas e seus subtipos. Classicamente, são divididas em hiperáridas, áridas e semiáridas, conforme o volume anual de chuvas (VAC), que não ultrapassa os 500 mm (NOY-MEIR, 1973). Nas regiões hiperáridas o VAC < 25 mm, nas áridas é 25 mm ≤ VAC < 250 mm, e nas semiáridas 250 mm ≤ VAC < 500 mm (HUANG et al., 2016). Essa definição é considerada incompleta por ter como base apenas o regime de chuva como critério delimitatório. Outra maneira que pode ser usada para identificar subtipos de regiões secas é a classificação climática de Köppen–Geiger que atualmente emprega características climáticas, chuva e temperatura para subdividir o clima do mundo em 31 categorias (KOTTEK et al., 2006). O clima árido (tipo B), segundo a classificação de Köppen–Geiger, pode ser subdividido em quatro: clima desértico quente (BWh), clima desértico frio (BWk), clima de estepe quente (BSh) e clima de estepe frio (BSk) (KOTTEK et al., 2006). Além disso, mesmo que os biomas não tenham limites tão fixos e precisos, sabe-se que os principais biomas que constituem as regiões secas são as Savanas, Pradarias e Desertos (HUANG et al., 2016; SAFRIEL et al., 2005). As diversas maneiras que podem ser usadas para classificar e definir as regiões secas contribuem para a compreensão desses ecossistemas por pontos de vista distintos. Sobrepondo essas visões pode-se construir uma compreensão mais completa desse tipo de ambiente.

Nas duas últimas décadas, o índice de aridez (IA) tem sido usado como uma boa ferramenta para classificação das regiões secas (HUANG et al., 2016). O IA é gerado pela razão entre a VAC média e demanda evaporativa anual média (expressa pela evapotranspiração potencial (PET) média anual), um valor de $IA < 1$ indica déficit de umidade. Segundo esse critério, regiões secas são áreas com $IA < 0,65$. É algo lógico imaginar que as regiões secas não são homogêneas e o IA permite classificá-las em quatro subtipos que diferem quanto a limitação de água, são eles: hiperáridas ($IA < 0,05$), áridas ($0,05 \leq IA < 0,2$), semiárida ($0,2 \leq IA < 0,5$), e subúmidas secas ($0,5 \leq IA < 0,65$), ocupando respectivamente, 6,6%, 10,6%, 15,2% e 8,7% da área do planeta (Figura 1) (HUANG et al., 2016; SAFRIEL et al., 2005).

Figura 1- Regiões secas em todo o mundo de acordo com índice de aridez (IA=VAC média/PET anual média).



Fonte: Modificado a partir de (MAESTRE et al., 2021).

Um fator que adiciona camadas de complexidade sobre ‘o que é’ uma região seca é que algumas dessas regiões podem receber grandes montantes de chuva durante parte do ano, um fato contraintuitivo quando pensa-se em regiões secas. Nesse contexto pode-se inserir as Florestas Tropicais Sazonalmente Secas (FTSS), esse tipo de bioma é classicamente reconhecido como uma floresta tropical com um peculiar regime de estações chuvosas e secas. Nas FTSS, a VAC varia entre 250-2000 mm e uma forte estação seca (meses com ≤ 100 mm de chuva) que pode durar seis meses ou mais. As FTSS podem representar mais de 40% de todas as florestas tropicais e subtropicais (ALLEN et al., 2017; BECKNELL; KISSING KUCEK; POWERS, 2012; PULLA et al., 2015). Quanto a esse bioma, a ciência mal arranhou a superfície do conhecimento, num período de 60 anos (entre 1945 e 2004) apenas 14% da produção científica confiável sobre Florestas Neotropicais tratavam de FTSS (PULLA et al., 2015).

A Caatinga constitui uma região semiárida e a maior FTSS da América do Sul, com área de ~1 milhão de km² cobrindo aproximadamente 10% do território brasileiro e 50% da região Nordeste (DA SILVA; LEAL; TABARELLI, 2017; MACHADO; LOPES, 2004; MORO et al., 2016; SANTOS et al., 2011). É formada por um mosaico de vegetação arbustivo-espinhosa decídua e manchas de floresta seca. A caatinga possui 135 unidades geoambientais e nove ecorregiões, além de ao menos 13 fitofisionomias (DA SILVA et al., 2017). Nesse ambiente, a pluviosidade é baixa (entre 240 e 1500mm por ano), e cerca de metade da área da Caatinga possui uma VAC média <750 mm e, em algumas áreas, <500 mm (LEAL et al., 2005; MORO et al., 2016; SANTOS et al., 2011). A estação chuvosa concentra-se em três meses consecutivos durante o ano e em alguns pontos 60% da VAC pode ocorrer em um único mês, o que gera um grande contraste na paisagem, como pode ser exemplificado na Figura 2 (MENEZES et al., 2012). Por outro lado, os períodos anuais de seca podem variar entre 7 e 11 meses. Além disso, os períodos de chuva podem variar entre os anos, gerando extensos períodos de seca contínua (LEAL et al., 2005; MACHADO; LOPES, 2004). Cerca de 28 milhões de pessoas vivem na Caatinga, sendo esta uma das regiões semiáridas mais populosas e biodiversas do mundo (DA SILVA; LEAL; TABARELLI, 2017).

Figura 2- Contraste paisagístico entre o período seco e chuvoso na Caatinga. Registro realizado na RPPN Tocaia, Sertão alagoano, município de Santana do Ipanema.



Fonte: Elaborado pelo autor (2022).

Por sua paisagem composta por plantas caducas durante grande parte do ano, longos períodos de seca, troncos secos e retorcidos e uma flora com muitos espinhos, construiu-se um mito no imaginário popular onde a Caatinga era um ambiente pobre em diversidade biológica. Desta maneira, esse domínio foi durante muito tempo negligenciado pelo Estado e pela ciência (LEAL et al., 2005). Nas últimas décadas

houve um esforço para desconstruir a imagem da Caatinga como um ambiente pobre, e o avanço dos estudos já contabilizam 3347 espécies de plantas, 548 de aves, 386 de peixes, 276 de formigas, 183 de mamíferos, 164 de fungos micorrízicos, 98 de anfíbios e 79 de répteis (DA SILVA et al., 2017; FERNANDES; CARDOSO; DE QUEIROZ, 2020; WINAGRASKI et al., 2019). Além disso, a taxa de endemismo é variável a depender do grupo em questão, sendo, por exemplo de 6% em mamíferos, 15,7% em plantas e 52,9% em peixes (DA SILVA et al., 2017; FERNANDES; CARDOSO; DE QUEIROZ, 2020).

Dentre os ‘Biomas’ brasileiros, a Caatinga é o menos estudado e com um número insuficiente de área protegidas, com apenas cerca de 7,4% da região com algum nível de proteção, entretanto boa parte não possui financiamento adequado (DA SILVA et al., 2017; LEAL et al., 2005; SANTOS et al., 2011). No ano de 2009, a Caatinga tinha uma perda de área superior a 45% e evidências que indicavam um aumento na velocidade do desmatamento, com a agropecuária se apresentando como uma das principais responsáveis (MMA, 2011). Menos de uma década depois a área da Caatinga modificada por ação humana é de cerca de 63,6% (DA SILVA et al., 2017) confirmando a tendência de 2009.

Na Caatinga, os períodos de seca são eventos normais que acontecem periodicamente, entretanto, secas longas historicamente causam prejuízos para as pessoas da região, gerando, por exemplo insegurança alimentar grave (MARENGO et al., 2011, 2018). Um dos fatores que contribuem para esses eventos de seca é o *El Niño*, um fenômeno climático-oceânico global caracterizado pelo aumento da temperatura da superfície do mar e alteração dos ventos alísios na região do Pacífico Equatorial. Esse processo afeta os regimes de chuva em regiões tropicais e de latitudes médias, porém não são capazes de explicar sozinhos os eventos de seca da Caatinga (KANE, 1989; MARENGO et al., 2011, 2018; MARENGO; TORRES; ALVES, 2016). Por outro lado, períodos chuvosos podem ser regulados entre outros fatores pelo fenômeno ‘contrário’, o *La niña*. Porém, os eventos de *La niña* estão ficando mais escassos e espaçados, enquanto as secas estão se tornando mais longas e severas (MARENGO et al., 2011; MARENGO; TORRES; ALVES, 2016).

As mudanças climáticas e o aquecimento global estão afetando todo o planeta, e na Caatinga a predição é que as mudanças aumentem o déficit de chuvas e consequentemente aumente a aridez, podendo levar a processos de desertificação e expansão de regiões áridas e semiáridas (HUANG et al., 2016; SAFRIEL et al., 2005). O aquecimento global que tem sido intensificado pela ação humana potencialmente pode gerar uma série de alterações no clima, e consequentemente, na biodiversidade, funções e serviços ecossistêmicos. Uma preocupante projeção indicava um aumento médio de 2 a 4 °C na temperatura e uma diminuição de 20% nas chuvas até 2100 (MARENGO et al., 2011; MARENGO; TORRES; ALVES, 2016). Entretanto, as atividades humanas têm acelerado esse processo e é possível que a marca de 4 °C de incremento na temperatura seja alcançada ainda na década de 2060 (MARENGO et al., 2018). O nordeste brasileiro é uma das regiões mais vulneráveis a mudanças climáticas, colocando em risco sua segurança hídrica, energética e alimentar (MARENGO et al., 2018). O bioma Caatinga é considerado um ambiente susceptível à aridificação e desertificação (cerca de 94% do semiárido nordestino é suscetível a desertificação). Ademais, já se observa em nível global uma tendência de expansão de áreas semiáridas (HUANG et al., 2016).

2.2. Ecologia de regiões secas

Diversos fatores bióticos e abióticos governam o funcionamento dos ecossistemas secos. Entre os abióticos, a quantidade de chuvas e a temperatura são centrais e alteram cadeias tróficas, ciclagem de nutrientes e padrões de coexistência de espécies (MAESTRE et al., 2016; NOY-MEIR, 1973; SCHWINNING et al., 2004). Nesses ecossistemas, a água pode estar direta ou indiretamente relacionada ao fluxo de energia, alterando o controle estomático, taxa de fotossíntese e respiração. Esses processos são intimamente relacionados à disponibilidade de água, sobretudo com insuficiência hídrica, condição que é dominante em regiões secas (COLLINS et al., 2014; NOY-MEIR, 1973). Além disso, a água também tem influência indireta em outras formas de transferência de matéria e energia como a herbivoria e carnivoria. Isso porque em ecossistemas áridos, a taxa de consumo de alimentos está muitas vezes

acoplada a disponibilidade de água no próprio alimento ou na água disponível para consumo (NOY-MEIR, 1973).

A disponibilidade de água é um fator limitante à vida, essa essencialidade é real e intuitivamente mais pronunciada em ambientes secos. Segundo Noy-meir (1973), os três atributos mais importantes para a compreensão do funcionamento dos ecossistemas secos são: i) a precipitação é escassa e, portanto, um fator dominante no controle dos processos biológicos. ii) Nesses ambientes a precipitação ocorre de forma ocasional e modesta. iii) A chuva nesse tipo de ambiente é variável e imprevisível. Ao indicar esses atributos centrais, Noy-Meir define desertos (e regiões secas) como ecossistemas controlados pela água e com chuvas escassas, reduzidas e imprevisíveis. Nesses ambientes, as chuvas geralmente não ultrapassam 2 mm, além disso, é estimado entre 10-50 dias chuvosos por ano, distribuídos entre 3-15 eventos de chuva, dos quais 5 ou 6 possuem magnitude para gerar efeitos biologicamente significativos (COLLINS et al., 2014; NOY-MEIR, 1973).

Ainda que a água não explique necessariamente todo o espectro de relações ecológicas em ambientes secos, inegavelmente ela afeta todos, seja de forma direta ou não (COLLINS et al., 2014). A relação entre funcionamento ecossistêmico em terras secas e a água vem sendo debatida há décadas, o prisma de conhecimento sobre esse tema indica que fatores diversos podem moldar a resposta do ambiente aos influxos de água (MAESTRE et al., 2016; SCHWINNING et al., 2004). Alguns destes fatores são: tipos funcionais de plantas, escala, tamanho e frequência das chuvas, limiares de resposta, textura e umidade prévia do solo (AUSTIN et al., 2004; CHESSON et al., 2004; MAESTRE et al., 2016; NOY-MEIR, 1973; OGLE; REYNOLDS, 2004; REYNOLDS et al., 2004). Por exemplo, solos com maior granulometria e/ou maior quantidade de matéria orgânica tendem a reter mais e por mais tempo a água, o que aumenta a chance dessa água ser usada por micro-organismos e plantas (MAESTRE et al., 2016; REYNOLDS et al., 2004; SCHWINNING et al., 2004). Por outro lado, solos mais arenosos permitem uma absorção mais rápida, desta forma, mesmo um grande volume de chuvas pode não ter efeito biológico efetivo (REYNOLDS et al., 2004).

Pode-se considerar que tudo que é consumido ou utilizado por um organismo, e durante esse processo deixa de estar disponível no meio, pode ser considerado um recurso (BEGON; TOWNSEND, 2020). Mudanças na disponibilidade dos recursos podem alterar o funcionamento dos ecossistemas em vários níveis de organização ecológica (SCHWINNING et al., 2004). No contexto da dinâmica de pulsos, pode-se entender um pulso como um evento efêmero e raro de superabundância de recursos (HOLT, 2008; YANG et al., 2008). Um pulso de recurso é um evento historicamente singular, ou seja, um evento de disponibilidade intensamente maior ao esperado para um determinado ambiente, não necessariamente promove relações adaptativas, mas sim respostas comportamentais e fisiológicas incomuns. Em suma, pulsos são fundamentalmente uma perturbação que gera um estado alternativo transiente (HOLT, 2008). Em resposta a essa repentina mudança da disponibilidade de recursos (pulso), as espécies podem alterar de forma intensa suas dinâmicas populacionais (ex. um pico de recrutamento de sementes ou de taxa metabólica microbiana). Esse uso do pulso de recurso pode ser direto ou indireto e tem potencial para afetar os processos em toda a comunidade e ecossistema em um efeito *bottom-up*, começando com a atividade microbiana e produção primária e, secundariamente, chegando aos níveis mais altos das cadeias. Além disso, o efeito *bottom-up* é seguido por um abrupto decaimento *top-down* (HOLT, 2008; YANG et al., 2008). Embora os pulsos de recursos sejam eventos de curta duração por definição, podem ser gatilhos importantes que iniciam processos poderosos e com efeitos ecológicos persistentes mesmo muito tempo depois do fim do pulso (AUSTIN et al., 2004; YANG et al., 2008).

Diversos ecossistemas respondem ao componente de pulsos, porém a disponibilidade de recurso acima do comum é o que caracteriza os pulsos dependendo da variação natural do ambiente em discussão (YANG et al., 2008). Em regiões secas, na maior parte do ano a situação é de déficit hídrico e, nesses ambientes, as chuvas ocorrem de forma escassa, reduzida e imprevisível. Portanto, pode-se considerar que as entradas de chuvas nesse sistema acontecem em ‘pulsos’ de curta duração (COLLINS et al., 2014; NOY-MEIR, 1973). Além dos efeitos diretos pelo acesso a água, os pulsos de chuva em ambientes secos podem gerar outros efeitos como à

translocação de nutrientes por percolação e escoamento, a disruptão da superfície do solo, afetando a disponibilidade de nutrientes, além da produção, deposição e decomposição de matéria orgânica (AUSTIN et al., 2004; COLLINS et al., 2014).

Pela natureza imprevisível das chuvas em regiões secas, esses ambientes apresentam flutuações abruptas entre seco-úmido em escala de dias ou até horas. Esse tipo de ciclo é comum em diversos ambientes e pode ocorrer em diversas escalas temporais, porém, tendem a ser mais abruptos em regiões secas altamente sazonais, como florestas tropicais decíduas. O ciclo seco-úmido afeta fortemente os processos microbianos e nutricionais do solo, incluindo mineralização de C e N, desnitrificação, volatilização de amônia e biomassa microbiana. Além disso, a maior umidade contribui em última instância para diminuição dos reservatórios de C e N (AUSTIN et al., 2004).

Os longos períodos de seca em *drylands* gera acúmulo de substratos minerais e orgânicos, pois o crescimento de plantas e micro-organismos é restrito durante esse período e há pouca demanda de nutrientes e menor taxa de decomposição e ciclagem (COLLINS et al., 2014). Por exemplo, o acúmulo de N ocorre majoritariamente no período seco, pois a difusão dos íons tende a ser restrita a meios aquosos e os ‘sumidouros’ de nitrogênio estão comprometidos nesse período (AUSTIN et al., 2004). Esse acúmulo leva a uma rápida e intensa taxa de mineralização e desnitrificação durante as fases iniciais do ciclo úmido. Desta forma, ciclos seco-úmido mais frequentes podem acentuar a perda de carbono e nitrogênio dos reservatórios do solo (AUSTIN et al., 2004; SCHWINNING et al., 2004).

Esse sistema de acúmulo e reservatórios podem ser diversos (ex. água, carbono e nitrogênio) e influencia as respostas ecossistêmica às novas precipitação, isso porque dependendo do tamanho do reservatório acumulado a resposta ambiental será diferente. Em certo nível, há uma “memória” ambiental dos eventos de precipitação anteriores. Essa memória, portanto, é uma das chaves para a compreensão da sensibilidade ambiental, sobretudo nos padrões intrazonais (SCHWINNING et al., 2004). Por exemplo, a duração de um período seco antes de um evento de chuva determina o tamanho do reservatório acumulado de matéria orgânica e N,

consequentemente, o tamanho da atividade de desnitrificação gerada pelo pulso de chuva (AUSTIN et al., 2004). Além disso, a duração da seca afeta também a prontidão fisiológica das plantas para o uso da água (OGLE; REYNOLDS, 2004; REYNOLDS et al., 2004; SCHWINNING et al., 2004). O efeito de memória torna difícil compreender a dinâmica ambiental considerando um único evento de chuva, ou numa escala de precipitação sazonal ou anual total (SCHWINNING et al., 2004). Isso porque a resposta a um pulso de chuva está relacionada não apenas a intensidade do pulso, mas também a distância temporal do último pulso ou a frequência dos ciclos seco-úmido.

Em ecossistemas que não experimentam um forte déficit hídrico, os processos de decomposição são predominantemente biológicos. Entretanto, em ecossistemas secos, principalmente nos períodos de baixa atividade microbiana, a decomposição da matéria orgânica pode ser mediada por fatores físico-químicos. Na superfície, a radiação solar e os ventos degradam, fragmentam e redistribuem nutrientes (MAESTRE et al., 2016). Essa maior contribuição de fatores físico-químicos na decomposição em regiões secas faz com que haja um potencial maior de decomposição do que de produção primária. Isso porque nesse contexto de limitação de água a produção primária é mais afetada que a decomposição (COLLINS et al., 2014). Ainda assim, os micro-organismos são agentes essenciais na dinâmica de ambientes secos. Micro-organismos do solo em latência (criptobióticos) podem responder a eventos de chuva pequenos e frequentes, que são insuficientes para gerar um efeito significativo em plantas, e assim podem dar início em processos de ciclagem e decomposição que aumentam a disponibilidade nutricional no solo (AUSTIN et al., 2004).

Em ecossistemas com limitação de água, a mineralização e imobilização de N por micro-organismos do solo tendem a ser regulados por três mecanismos: i) a relação C:N, ii) eficiência do uso de N pela microbiota (quantidade de N necessária para produzir uma unidade de biomassa microbiana C) e iii) a eficiência do crescimento microbiano (fração do C orgânico usado pelo micro-organismo para produzir biomassa) (AUSTIN et al., 2004). Dentre esses fatores, o que provavelmente mais impacta o balanço entre mineralização e imobilização de N é a relação C:N. Dependendo do

substrato a relação C:N varia substancialmente. Para a biomassa microbiana a relação é cerca de 4:1, em raízes fica entre 30:1 até 70:1 e em material lenhoso chega a 100:1 (AUSTIN et al., 2004). Desta maneira, espera-se que valores baixos de C:N elevem as taxas metabólicas microbianas e favoreçam a mineralização de N, enquanto valores elevados da relação desviam as rotas para a imobilização (AUSTIN et al., 2004).

Por sua vez, a eficiência do uso de N é essencialmente o inverso da relação C:N da biomassa microbiana e é determinada em grande parte pela estrutura da comunidade microbiana do solo (AUSTIN et al., 2004). As bactérias produzem biomassa com uma relação C:N mais baixa do que os fungos e assim imobilizam mais N por unidade de C assimilado, do que comunidades dominadas por fungos. Por outro lado, fungos são mais tolerantes a dessecação que bactérias, pois geralmente toleram menor atividade de água. Por isso, a abundância relativa de fungos tende a aumentar em comunidades de ambientes áridos e semiáridos durante o período seco. A resposta de mineralização ou imobilização durante os ciclos de umidade tendem a ser determinada pela taxa de crescimento relativo entre bactérias e fungos (AUSTIN et al., 2004). Como os solos das terras áridas são cronicamente secos, os fungos, em vez das bactérias, podem dominar os processos de decomposição e transformação de N, além de gerar associações simbióticas com produtores primários, por exemplo, micorrizas (COLLINS et al., 2014).

2.3. Micro-organismos associados às plantas

Plantas possuem associação com micro-organismos tanto em suas estruturas subterrâneas quanto nas aéreas (KOSKELLA, 2020; VORHOLT, 2012). A parte subterrânea, que compreende a interface solo-planta, é denominada rizosfera (BRINGEL; COUÉE, 2015; KOSKELLA, 2020). Por sua vez, ainda que exista certo grau de confusão na definição, a parte aérea das plantas, que compreende todas as estruturas da interface atmosfera-planta, é denominada de filosfera (KOSKELLA, 2020; LINDOW; BRANDL, 2003; VORHOLT, 2012). A filosfera inclui folhas (filoplano), frutos (carposfera), flores (antosfera, incluindo pólen e néctar), caule (caulosfera), e reservatórios de água (fitotelma) (BRINGEL; COUÉE, 2015; KOSKELLA, 2020;

LEVEAU, 2019; LIU; BRETELL; SINGH, 2020; VACHER et al., 2016; VORHOLT, 2012). Nos diferentes tecidos das plantas, os micro-organismos podem colonizar o interior (endofíticos) ou na superfície (epifíticos). Também é comum que a mesma espécie de micro-organismo possa ser encontrada em ambos os habitats (FONSECA; INÁCIO, 2006; VORHOLT, 2012). Além disso, pode-se agrupar os componentes da microbiota vegetal em transientes (micro-organismos efêmeros, que estão temporariamente associados) e residentes (micro-organismos muito frequentes, fortemente associados) (FONSECA; INÁCIO, 2006). Esses ambientes sustentam a maior diversidade de micro-organismos do planeta, com representantes de diversos grupos como bactérias, arqueas, vírus, fungos filamentosos, leveduras, microalgas, protozoários e nematoides (GOFFREDI; JANG; HAROON, 2015; KOSKELLA, 2020; LINDOW; BRANDL, 2003; THAPA; PRASANNA, 2018; VACHER et al., 2016; VORHOLT, 2012).

A maioria dos estudos sobre a microbiota das plantas se concentra na região da rizosfera (BRINGEL; COUÉE, 2015; KOSKELLA, 2020). Nessa região, a abundância microbiana é maior que na parte aérea das plantas (ANDREOTE; GUMIERE; DURRER, 2014; KOSKELLA, 2020; VORHOLT, 2012). Essa constatação sugere de forma errada que a microbiota da filosfera é de alguma maneira menos importante. Entretanto, a filosfera é a interface ecológica que permeia a relação entre o ambiente e a planta (KOSKELLA, 2020; VACHER et al., 2016). Diversos estudos têm enumerado contribuições essenciais da microbiota da filosfera em processos de defesa, patogenicidade, aquisição de nutrientes, crescimento, sucesso reprodutivo e evolução das plantas (KOSKELLA, 2020; LEVEAU, 2019; RODRIGUEZ et al., 2009; THAPA; PRASANNA, 2018). Por exemplo, a microbiota do néctar é capaz de alterar padrões de preferência de polinizadores, elevando ou diminuindo a frequência de visitação floral, e até mesmo alterando a preferência de diferentes guildas, o que tem impacto direto no sucesso reprodutivo e evolução vegetal (CANTO; HERRERA; RODRIGUEZ, 2017; SCHAEFFER et al., 2017). Além disso, a microbiota pode agir como antagonista a fitopatógenos, mediando a saúde da planta, ou facilitando a infecção por meio da produção de enzimas que degradam a cutícula da planta (BUCK, 2002; LEVEAU,

2019). Por exemplo, em experimento onde uma esterase produzida pela levedura *Pseudozyma antarctica* foi aspergida sobre plantas de tomate e infectada com o fitopatógeno *Botrytis cinerea*, a infecção foi mais grave (UEDA et al., 2018). Isso não indica que a levedura seja patogênica, mas que ela pode ser um potencial facilitador da infecção.

Estima-se que a filosfera represente 60% da biomassa de todos os organismos do planeta (KOSKELLA, 2020). Esse ambiente é extremo, dinâmico, heterogêneo e frequentemente oligotrófico (KOSKELLA, 2020; LEVEAU, 2019; VORHOLT, 2012). Além da elevada exposição à radiação solar e das oscilações bruscas de fatores de estresse como temperatura, oxidação, disponibilidade de nutrientes e água, outro ponto crítico é a camada de cutina, que forma a cutícula da planta (KOSKELLA, 2020; LEVEAU, 2019; MEYER; LEVEAU, 2012; VORHOLT, 2012). A cutícula da planta é uma camada lipídica formada por ácidos graxos de cadeia longa e chega a representar 15% do peso seco das folhas (BRINGEL; COUÉE, 2015). Essa camada protege as folhas e auxilia no controle hídrico, é também a primeira barreira para o estabelecimento dos micro-organismos da filosfera (LINDOW; BRANDL, 2003). A cutícula tem a capacidade de modelar a microbiota através de variações em sua composição química e arquitetura (e.g. rugosidade e tricomas), alterando padrões microbianos de colonização, fixação, composição e abundância (LEVEAU, 2019; LINDOW; BRANDL, 2003; REISBERG et al., 2013; VACHER et al., 2016; VORHOLT, 2012).

A cutícula da planta diminui a evaporação, retenção de água na superfície, lixiviação de metabólitos e difusão de nutrientes, gerando um ambiente adverso e com elevado estresse osmótico (BEATTIE; LINDOW, 1995a; BRINGEL; COUÉE, 2015; LINDOW; BRANDL, 2003; VACHER et al., 2016; VORHOLT, 2012). Por outro lado, a limitação da lixiviação também evita a perda precoce de nutrientes e colabora para manutenção destes nas regiões epifíticas (WHIPPS et al., 2008). Nesse sentido, para a microbiota filosférica, a cutícula pode ser interpretada ao mesmo tempo como uma vantagem e uma desvantagem. Pois, se por um lado gera estresse osmótico, por outro, preserva reservas nutricionais importantes.

Cada compartimento da filosfera (e.g. filoplano, antosfera, carposfera) possui uma microbiota distinta e particular (ABDELFATTAH et al., 2019; COLEMAN-DERR et al., 2016; FÉLIX et al., 2022; LEVEAU, 2019; LIU; HOWELL, 2021). O genótipo é um importante estruturador de comunidades microbianas associados a plantas (LEVEAU, 2019; SAPKOTA et al., 2015). Um mesmo indivíduo possui, por óbvio, um mesmo genótipo. Entretanto, esse genótipo é expresso de forma diferente em cada um dos compartimentos vegetais, o que pode explicar parte das diferenças entre as comunidades (LEVEAU, 2019). Entre os compartimentos da filosfera, o filoplano é o maior e um dos mais estudados (FONSECA; INÁCIO, 2006; KOSKELLA, 2020). A superfície das folhas do mundo soma uma área de aproximadamente 1 milhão de km², o equivalente ao dobro da superfície do planeta Terra (BRINGEL; COUÉE, 2015). É estimado que as folhas abriguem em média 10⁶-10⁷ células de bactérias por cm² e em valores totais cerca de 10²⁶ células (LINDOW; BRANDL, 2003; VORHOLT, 2012). Considerando que essa estimativa não considera células de fungos e demais micro-organismos, o filoplano é de fato um dos maiores habitats microbianos do mundo. Além disso, estima-se que 96% das bactérias da filosfera sejam comensais, portanto não possuem efeito sobre a saúde do hospedeiro, enquanto 2% são patogênicas e 2% promovem crescimento e possuem funções benéficas ao hospedeiro (LINDOW; LEVEAU, 2002).

A configuração microclimática adversa e com constantes flutuações da filosfera demanda aos micro-organismos transpor uma série de barreiras e gargalos ambientais e ecológicos para colonizar esse ambiente (BEATTIE; LINDOW, 1995a; LINDOW; BRANDL, 2003; THAPA; PRASANNA, 2018; VORHOLT, 2012). Micro-organismos da filosfera geralmente interagem com esse ambiente por meio de duas estratégias básicas. A primeira é a tolerância, que está relacionada a capacidade de suportar as condições ambientais extremas da filosfera, por exemplo, produzindo pigmentos fotoprotetivos e exopolissacarídeos (EPS) que auxiliam na tolerância à radiação UV e dessecação (BEATTIE; LINDOW, 1995a; THAPA; PRASANNA, 2018). A segunda é a estratégia de prevenção ou resguardo, que tem como cerne empregar mecanismos que evitem a direta exposição dos micro-organismos aos estressores

ambientais da filosfera. E essa pode ser, por exemplo, uma estratégia empregada por micro-organismos que crescem de forma endofítica (BEATTIE; LINDOW, 1995a; THAPA; PRASANNA, 2018). De certo, essas estratégias não são mutuamente excludentes, ou seja, os micro-organismos podem possuir as duas simultaneamente (BEATTIE; LINDOW, 1995a).

Segundo Kinkel (1997), a comunidade microbiana da filosfera é regulada por quatro processos populacionais; i) imigração, ii) emigração, iii) crescimento (geração) e iv) morte. Para Vellend (2010), os mecanismos que regulam as comunidades ecológicas são diversos, mas todos podem ser sintetizados em apenas quatro tipos de processos: seleção, deriva, especiação e dispersão. Nesse contexto, a seleção reflete diferenças na capacidade adaptativa (fitness); a deriva se refere as mudanças estocásticas nas abundâncias relativas das espécies em uma comunidade, a especiação se refere ao surgimento de novas espécies e dispersão trata do movimento dos organismos no ambiente (VACHER et al., 2016; VELLEND, 2010).

Um vasto número de fatores ambientais, ecológicos e evolutivos pode atuar sobre os processos que estruturam as comunidades da filosfera. Alguns dos fatores que alteram dispersão e colonização dos micro-organismos da filosfera envolvem as chuvas que promovem a remoção daqueles já presentes e trazem outros da atmosfera e do dossel adjacente; os ventos que podem transportar esporos e/ou micro-organismos na forma de bioaerosol e depositá-los sobre vegetais; animais (e.g. insetos e aves) que funcionam como vetores e transportam micro-organismos entre diferentes plantas e compartimentos. A microbiota pode ser passada por transferência vertical, onde a parental transmite a microbiota para as sementes (KOSKELLA, 2020; LEVEAU, 2019; VACHER et al., 2016; VORHOLT, 2012). A comunidade epífita é afetada pela sazonalidade, fatores climáticos, genótipo e ontogênese da planta (BRINGEL; COUÉE, 2015; FONSECA; INÁCIO, 2006; VACHER et al., 2016; VORHOLT, 2012). Além dos fatores externos, as interações bióticas entre os micro-organismos que regulam os padrões de coexistência também estruturam a microbiota (VACHER et al., 2016).

A principal fonte de nutrientes orgânicos para os micro-organismos da filosfera (sobretudo das folhas) são os exsudados disponibilizados pelas plantas, enquanto os nutrientes inorgânicos são geralmente obtidos do meio externo, por exemplo, pela deposição dos ventos (FONSECA; INÁCIO, 2006; KEMLER et al., 2017). Desta forma, a fisiologia da planta possui um papel vital na disponibilidade nutricional e, consequentemente, na estrutura da comunidade microbiana. Duas vias estão relacionadas ao transporte de carboidratos nas plantas, a via simplástica e a apoplástica. Na via simplástica, o movimento de carboidratos se dá entre os citoplasmas célula-a-célula. Por outro lado, a via apoplástica, depende do movimento da sacarose nos espaços extracelulares, o apoplasto. Nesse tipo de transporte, alguns açúcares presentes no apoplasto não são carregados para o floema e difundem-se através da cutícula, um mecanismo essencial para a disponibilização de nutrientes para o filoplano. Esse mecanismo não se restringe a disponibilização de açúcares e pode transportar outros nutrientes orgânicos e inorgânicos (VACHER et al., 2016). Além disso, enquanto a composição de carboidratos no interior das folhas é dominada por sacarose, na superfície os açucares menores são mais frequentes como a glicose e frutose (VACHER et al., 2016).

2.4. A relação da microbiota da filosfera com a água

A filosfera é a porta da frente na interação planta-ambiente, é ela que recebe em primeira mão os efeitos do ambiente externo. A microbiota da filosfera, portanto, recebe esses estressores juntamente com o hospedeiro. Uma das condições mais desafiadoras e mais comuns que os micro-organismos enfrentam na filosfera é o estresse por déficit hídrico. Mesmo em ambientes mais úmidos, a cutícula é uma superfície hidrofóbica e com baixa molhabilidade (capacidade de um líquido em manter contato com uma superfície sólida), e quanto maior a cerosidade, maior é a dificuldade para a permanência da água nas regiões epifíticas como as folhas, gerando um ambiente com pouca água disponível e com grande pressão osmótica pelo acúmulo de metabólitos. Portanto, mecanismos de osmoproteção são vitais para a sobrevivência na filosfera, podem incluir acúmulo e/ou síntese de solutos que invertem a direção da pressão osmótica para dentro das células (BEATTIE; LINDOW, 1995b; THAPA;


PRASANNA, 2018). A trealose, por exemplo, é um osmorregulador importante e comumente usado como estratégia de sobrevivência pela bactéria *Pseudomonas syringae* (THAPA; PRASANNA, 2018).

Alguns fatores podem alterar a disponibilidade de água na filosfera, tais como o envelhecimento do hospedeiro e o regime de chuvas (VACHER et al., 2016). O envelhecimento vegetal tende a alterar características físico-químicas na cutícula, que passa a ter maior molhabilidade e permeabilidade, aumentando a adesão da água e a disponibilidade de nutrientes (OSO et al., 2021; VACHER et al., 2016). Por outro lado, o efeito da chuva nas comunidades da filosfera ainda não está elucidado, enquanto alguns estudos sugerem um incremento na abundância (ALLARD et al., 2020), outros, sugerem um decréscimo (ALLARD et al., 2020; KINKEL, 1997). Há ainda estudos que indicam a ausência de efeito da chuva na comunidade da filosfera (STONE; JACKSON, 2021). É possível que a frequência e intensidade das chuvas também tenham relação com a resposta microbiana (KINKEL, 1997). Sabe-se que as chuvas alteram as taxas de colonização e dispersão, promovendo lavagem das folhas e retirada de micro-organismos, ao mesmo tempo que transferem para a planta outros micro-organismos da atmosfera (LEVEAU, 2019; MORRIS, 2001; VACHER et al., 2016).

Pela natureza oligotrófica e adversa da filosfera, esse ambiente tende a selecionar micro-organismos capazes de ‘remodelá-lo’ para acessar os recursos. Por exemplo, estreitando a cutícula, produzindo hormônios, enzimas, exopolissacarídeos (EPS) e biossurfactantes (BEATTIE; LINDOW, 1995b; LEVEAU, 2019). Algumas dessas estratégias são capazes de aumentar e/ou facilitar o acesso à água e a tolerância a seca do micro-organismo. Por exemplo, o fitopatógeno *P. syringae* pode aumentar a molhabilidade e a disponibilidade local de água na folha liberando biossurfactantes (KOSKELLA, 2020). Devido à natureza hidrofóbica da cutícula, o aumento da molhabilidade desse ambiente pode permitir a solubilização e difusão de nutrientes à medida que permite uma maior adesão da água (Figura 3) (LINDOW; BRANDL, 2003). Folhas muito cerasas possuem pouca molhabilidade e mesmo se houver água disponível através da umidade ou da chuva, dificilmente se formará um filme de água. A formação desse filme de água, além de promover disponibilidade

ídrica para a comunidade microbiana na folha, altera rapidamente o pH e redistribui nutrientes na região epífítica (MORRIS, 2001).

Figura 3 - Esquema indicando como a diminuição da tensão superficial e aumento da molhabilidade, que pode ser mediado por biossurfactantes, permite um maior contato da água com superfícies hidrofóbicas como as folhas.



Fonte: Elaborado pelo autor com imagens de He et al.(2021).

Biossurfactantes são moléculas quimicamente diversas produzidas por vários grupos microbianos que tem como principal característica ser anfipáticas, ou seja, possuem uma região polar e uma apolar. Essas moléculas possuem numerosas atividades como: diminuição de tensão superficial, aumento da molhabilidade em superfícies hidrofóbicas, emulsificação entre líquidos imiscíveis e solubilização de hidrocarbonetos (BEATTIE; LINDOW, 1995b; OSO et al., 2021; THAPA; PRASANNA, 2018; ZEISLER-DIEHL; BARTHLOTT; SCHREIBER, 2020). A produção de enzimas que degradam a cutícula e de biossurfactantes tem sido proposta como um mecanismo que pode aumentar a permeabilidade nas folhas e facilitar a movimentação de bactérias no filoplano (DOAN; LEVEAU, 2015; LEVEAU, 2019; LINDOW; BRANDL, 2003; OSO et al., 2021). Assim, a produção de biossurfactantes pode ser uma ferramenta importante

para que micro-organismos epifíticos alterem seu habitat para explorá-lo com maior eficiência (LEVEAU, 2019; LINDOW; BRANDL, 2003).

Por sua vez, a produção de EPS, formação de uma matriz extracelular e biofilme podem auxiliar a comunidade microbiana da filosfera, conferindo, entre outros atributos, tolerância a dessecação. Muitas vezes os EPSs que formam a matriz são higroscópicos (capazes de reter água), conferindo vantagem num ecossistema com déficit hídrico. Além disso, a matriz gerada pode dar gênese a um biofilme microbiano, que é uma estratégia onde populações ou comunidades microbianas formam um complexo e estruturado agregado celular, capaz de responder de forma coordenada a pressões ambientais (OSO et al., 2021; THAPA; PRASANNA, 2018; ZEISLER-DIEHL; BARTHLOTT; SCHREIBER, 2020).

2.5. Leveduras

Leveduras constituem um grupo artificial formado por várias linhagens de fungos com convergência na regulação de mecanismos moleculares que permitem a manutenção de um estado unicelular (NAGY et al., 2014). Apesar de serem predominantemente unicelulares, algumas espécies alternam entre fase unicelular e filamentosa, podendo transitar entre fases intermediárias através da formação de pseudohifas e aglomerados celulares, sendo conhecidas como *yeast-like*, fungos dimórficos ou fungos semelhantes a leveduras (BASTIDAS; HEITMAN, 2009; BOEKHOUT et al., 2011). Assim como nos demais fungos, a parede celular das leveduras também é formada por polissacarídeos estruturais microfibrilares, representados principalmente por quitinas e β -glicanas, responsáveis por conferir rigidez e resistência à parede. Outros compostos como β homo- e hétero-polissacarídeos, proteínas, pequenos lipídeos e sais inorgânicos também são encontrados na parede e atuam como elementos de sedimentação (ORTIZ-CASTELLANOS; RUIZ-HERRERA, 2015).

Fungos podem ser monomórficos ou dimórficos, apresentando transições morfológicas reguladas por fatores moleculares e diferentes gatilhos ambientais (BASTIDAS; HEITMAN, 2009; NAGY et al., 2014). Os mecanismos internos

e externos que desencadeiam essas mudanças morfológicas são diversos e podem apresentar ligação com a fase do ciclo de vida do organismo, fatores nutricionais, evolutivos e com estresses ambientais (ex. temperatura) (BASTIDAS; HEITMAN, 2009; NAGY et al., 2014). A transição entre estes estágios morfológicos pode estar associada a processos patogênicos em algumas espécies, como em *Candida albicans*, que é capaz de mudar (*shift*) para um estado filamentoso que beneficia a adesão às mucosas do hospedeiro (BASTIDAS; HEITMAN, 2009; COOPER, 2011). A linha que separa um estado morfológico de outro (unicelular, dimórfico ou filamentoso) é um terreno cinzento e ainda sem consenso. Provavelmente a morfologia é um continuum, desta forma a nomenclatura estática e categórica utilizada para denominá-los possui caráter utilitário e pedagógico. Além disso, considerando que todos os fungos (ou ao menos a maioria das linhagens) possuem as ferramentas moleculares necessários para a gênese e manutenção do estado unicelular (NAGY et al., 2014), é provável que espécies que hoje são consideradas estritamente filamentosas possam ser encontradas no futuro em um estado leveduriforme e vice-versa.

A principal forma de reprodução das leveduras é assexuada, via brotamentos, fissão ou produção de esporos ejetáveis (balistosporos). Porém, podem apresentar estágios de reprodução sexual sem a produção de corpos de frutificação (KURTZMAN, 2011a, 2011b). Em leveduras a poliploidia é um fenômeno comum (BOEKHOUT et al., 2011; ROBERTS; OLIVER, 2011), o aumento no número da ploidia pode estar relacionado a uma resposta a pressões ambientais. A poliploidia permite que o organismo tenha um maior número de genes disponíveis em seu “arsenal”, que por sua vez, pode influenciar seu *fitness* (VOORDECKERS et al., 2015). Outro ponto é que a poliploidia tem nuances biotecnológicas, por exemplo, muitas linhagens poliploides têm sido aplicadas em processos fermentativos por serem superiores em desempenho e mais tolerantes a estresses do processo industrial como valores extremos de temperatura e pH (SELMECKI et al., 2015).

Filogeneticamente, leveduras são agrupadas em dois filos do reino Fungi: Ascomycota e Basidiomycota (sub-reino Dikarya) (KURTZMAN, 2011a, 2011b). O filo Ascomycota possui três subfilos: Pezizomycotina, Taphrinomycotina e

Saccharomycotina (KURTZMAN, 2011a, 2011b). A sinapomorfia que une o filo dos ascomicetos é a formação de esporos meióticos em ascos, conhecidos como ascósporos (SCHOCH et al., 2009). O filo Basidiomycota também possui três subfilos, são eles: Pucciniomycotina, Agaricomycotina e Ustilaginomycotina (KURTZMAN, 2011a, 2011b). A sinapomorfia que une o grupo são esporos meióticos ‘nus’ (basidiósporos), sem asco, que se originam de uma estrutura em forma de haste (basídio) (BOEKHOUT et al., 2011).

As leveduras são organismos heterotróficos e as enzimas extracelulares possuem papel fundamental em sua nutrição. São elas que hidrolisam macromoléculas e deixam os nutrientes disponíveis para que a célula possa absorvê-los, e desta forma, direta ou indiretamente mediam os processos de decomposição (ORTIZ-CASTELLANOS; RUIZ-HERRERA, 2015). Leveduras são importantes decompósitos nos ecossistemas e um dos primeiros grupos a colonizar substratos ricos em nutrientes (FONSECA; INÁCIO, 2006; GANTER; MORAIS; ROSA, 2017; STARMER; LACHANCE, 2011). Esses organismos estão distribuídos ao redor do planeta, presentes em múltiplos habitats e nichos, associados a plantas, animais e até mesmo a outros fungos. Estão presente em substratos como água doce e salgada, solo e atmosfera, em ambientes naturais e urbanos, úmidos e desérticos (BOEKHOUT et al., 2022; ROSA; PÉTER, 2006; STARMER; LACHANCE, 2011).

Leveduras são capazes de formar biofilme e, por exemplo, alterar a textura do solo, contribuindo consideravelmente para a biomassa microbiana destes. Muitas leveduras são capazes de produzir fatores que estimulam o crescimento das plantas e antagonizar com fitopatógenos (BOTH, 2011; BRINGEL; COUÉE, 2015; MOLLER; LERM; BOTH, 2016). As várias interações ecológicas que leveduras podem participar incluem amensalismo, competição, predação, mutualismo e até mesmo como parasitas de outros fungos (BOTH, 2011). Um bom exemplo é que alguns líquens podem ser constituídos de associação mutualísticas entre macrofungos, leveduras e algas (SPRIBILLE et al., 2016). Em relação ao padrão biogeográfico de leveduras, Tedersoo et al. (2014) demonstraram que em nível global, leveduras de solo diminuem sua

diversidade à medida que se afastam dos trópicos, seguindo o padrão geral de distribuição de espécies, que se aplica também para macro-organismos.

Várias espécies de *Papiliotrema* (anteriormente classificadas como *Cryptococcus*) podem produzir cápsulas polissacarídicas que conferem resistência à dessecação (FONSECA; INÁCIO, 2006; KEMLER et al., 2017). Além disso, características como a produção de pigmentos podem gerar uma vantagem competitiva em um ambiente exposto a diversos estresses oxidativos, como a superfície das folhas. Estes pigmentos são diversos, mas são muitas vezes carotenóides que são amplamente encontrados em membros de Sporidiobolales (por exemplo, *Rhodotorula* e *Sporobolomyces*), ou melanina, produzida por leveduras negras como *Aureobasidium*, *Hortaea* e *Valentiella* (BEZERRA et al., 2022; FONSECA; INÁCIO, 2006; KEMLER et al., 2017). Algumas espécies de *Sporobolomyces* e *Bullera* produzem balistosporos, esporos ejetáveis com função de dispersão (FONSECA; INÁCIO, 2006; KEMLER et al., 2017). Leveduras basidiomicetos são geralmente mais generalistas e capazes de assimilar diversas fontes de carbono (politróficas), tendo vantagem em ambientes oligotróficos. Portanto, são frequentes e dominantes na filosfera, principalmente associados diretamente à cutícula (FÉLIX et al., 2022; FONSECA; INÁCIO, 2006; HAGLER et al., 1993; KEMLER et al., 2017).

Leveduras da filosfera tendem a ocorrer em folhas em uma abundância média de 10^3 a 10^5 CFU/g, relativamente baixo comparado a bactéria (GOUKA; RAAIJMAKERS; CORDOVEZ, 2022). Fatores com a disponibilidade de nutrientes, e genótipo do hospedeiro podem afetar a abundância de leveduras da filosfera (GOUKA; RAAIJMAKERS; CORDOVEZ, 2022). A montagem da comunidade de leveduras da filosfera envolve uma gama de fatores estocásticos e determinísticos. Por exemplo, a colonização possui facetas estocásticas e ocorre principalmente através de vetores como a chuva, os ventos e animais (majoritariamente insetos e aves). Por outro lado, a seleção envolve processos determinísticos relacionados a espécies do hospedeiro e ao fitness das espécies ou linhagens de levedura colonizadora (ANDREWS et al., 1987; BLACKWELL, 2017; KINKEL, 1997; MITTELBACH et al., 2015; VACHER et al., 2016; VELLEND, 2010). A abundância de leveduras da filosfera pode ser afetada por diversos

fatores. Alguns estudos sugerem que a sazonalidade e a senescência do hospedeiro incrementam essa abundância (FONSECA; INÁCIO, 2006; GLUSHAKOVA; CHERNOV, 2004, 2007; GOUKA; RAAIJMAKERS; CORDOVEZ, 2022).

Leveduras da filosfera apresentam uma ampla diversidade de interações com o hospedeiro e com outros micro-organismos. Na interação com o hospedeiro, as leveduras podem estimular o crescimento vegetal produzindo hormônios como o ácido indolacético e elicitando as defesas da planta através das vias do ácido jasmônico, ácido salicílico e etileno (GOUKA; RAAIJMAKERS; CORDOVEZ, 2022). Em interações com micro-organismos a produção de toxinas-*killer* pode oferecer vantagem na competição sobre outras leveduras (GOUKA; RAAIJMAKERS; CORDOVEZ, 2022). Leveduras possuem a capacidade se acessar e assimilar uma gama de compostos orgânicos, bem como, de tolerar e degradar compostos tóxicos (micotoxinas e antifúngicos), essas habilidades conferem vantagens competitivas por espaço e nutriente sobre outros grupos de micro-organismos como bactérias e fungos filamentosos (GOUKA; RAAIJMAKERS; CORDOVEZ, 2022).

Há uma grande lacuna em relação a riqueza de fungos no mundo. Uma estimativa recente é de que existam entre 2,2 e 3,8 milhões de espécies de fungos, porém o número de espécies corretamente descritas é de aproximadamente 120.000 (de 3,15 a 5,45% do total estimado) (HAWKSORTH; LÜCKING, 2017). Em relação a leveduras, a estimativa é de cerca de 200.000 espécies (BOEKHOUT et al., 2022) e a diversidade de leveduras atualmente descrita está entre 2.200 e 2.300 espécies (BOEKHOUT et al., 2022; YURKOV et al., 2021), pouco mais de 1% da riqueza total estimada.

Alguns dos maiores reservatórios de novidades taxonômicas de fungos filamentosos e leveduras são plantas vasculares (sobretudo na América do Sul), insetos e solo (BOEKHOUT et al., 2022; HAWKSORTH; LÜCKING, 2017). Nas últimas décadas houve aumento no número de espécies de leveduras descritas. Entre os anos de 1952-1984 a taxa de descrição era de 10 espécies por ano, entretanto, nos últimos anos esse número tem estado acima de 60 espécies (BOEKHOUT et al., 2022). Além

disso, Li et al. (2020), descreveram mais de 100 espécies em um único estudo, o que indica o avanço dessa área. Alguns dos ecossistemas que continuam subamostrados são os ambientes áridos e gélidos, porém, esses parecem promissores para a descoberta de novas espécies (BOEKHOUT et al., 2022).


2.6. Bromélias

Bromeliaceae é uma família de plantas quase exclusivamente neotropical reconhecida, entre outros fatores, por sua grande diversidade, alto endemismo e por fornecer diversos serviços ecossistêmicos de apoio à biodiversidade e regulação hídrica (BENZING, 2000; LADINO et al., 2019). A família é composta por 3500 espécies, 50 gêneros e 8 subfamílias. Bromélias possuem distribuição por diversos países Sul e Centro-americanos, chegando a ocorrer até o Sudeste da América do Norte. Apenas uma espécie é encontrada fora das américas, a espécie *Pitcarnia feliciana*, que ocorre na Costa Oeste da África (BENZING, 2000; MANETTI; DEIAPORTE; LAVERDE, 2009). A espécie de bromélia mais conhecida e com maior importância econômica é *Ananas comosus* (abacaxi) que é a única espécie da família cultivada extensivamente para alimentação (MANETTI; DEIAPORTE; LAVERDE, 2009). Bromélias atraem o interesse científico devido às suas associações com vários grupos de macro e micro-organismos, que variam de invertebrados e anfíbios a vírus, bactérias e leveduras como ilustrado na Figura 4 (GOFFREDI; JANG; HAROON, 2015; LANDELL et al., 2015; LEROY et al., 2016, 2017).

Cerca de metade dos gêneros de bromélias e mais da metade das subfamílias contém espécies capazes de formar tanques de água no centro da roseta, uma estrutura chamada fitotelma (LADINO et al., 2019; MALES; GRIFFITHS, 2017). Esse tanque de água contribui, por exemplo, para a reprodução de diversas espécies animais, tanto vertebrados como anfíbios, quanto invertebrados como artrópodes (BENZING, 2000; LADINO et al., 2019). Dependendo da espécie e das condições ambientais, o fitotelma pode acumular de 0,015 a 45 L de água em um único indivíduo (ZOTZ et al., 2020), e cerca de 50.000 L em um hectare (LADINO et al., 2019). A capacidade de formar o fitotelma faz com que bromélias, sob determinadas condições,

consigam alterar a dinâmica hídrica local (LADINO et al., 2019). Bromélias podem funcionar como um ecossistema insular em microescala (FRANK; LOUNIBOS, 1987), e talvez também como refúgio em épocas de seca, principalmente para organismos aquáticos ou fortemente dependentes da água.

Figura 4- Ilustração de um corte longitudinal em uma bromélia capaz de formar tanque (fitotelma).



Fonte: (BOND, 1975).

Inovações evolutivas, estratégias ecofisiológicas, além de diversos mecanismos de captação de água, explicam o amplo espectro de ocorrência das bromélias. O grande espectro ambiental de bromélias vai desde o seu hábito que pode ser epífito, terrestre ou saxícola (rupícolas) e sua distribuição, que ocorre de 0 a 4000 metros de altitude, de florestas seca a úmidas e de ambientes tropicais a temperados (BENZING, 2000; LADINO et al., 2019; MALES; GRIFFITHS, 2017). As espécies exclusivamente epífitas possuem adaptações fisiológicas e morfológicas que permitem sobreviver sem contato com o solo, absorvendo água e nutrientes apenas através de tricomas nas folhas (BASÍLIO et al., 2015; BENZING, 2000). Nestas espécies, com destaque para as do gênero *Tillandsia*, as raízes perdem total ou parcialmente a função de absorção e passam a ter função de fixação ao substrato.

Traços importantes que contribuíram para o sucesso evolutivo das bromélias são: os tricomas absorventes das folhas, o surgimento do metabolismo fotossintético tipo CAM (Metabolismo ácido das crassuláceas), a capacidade de retenção de água no tanque e a suculência (BENZING, 2000; MALES; GRIFFITHS, 2017). Essas características são importantes para a tolerância à seca e, possivelmente, para a colonização de ambientes xerófilos e de altitude. Males e Griffiths (2017) combinaram a filogenia das bromélias com cinco tipos funcionais, são eles: C₃ terrestres, C₃ tanque-epífitas, CAM atmosférica-epífitas, CAM terrestres e CAM tanque-epífitas. Entre esses tipos funcionais, foram observadas variações morfológicas, ecológicas e fisiológicas. Por exemplo, existem diferenças em características relacionadas à tolerância à seca, como massa de água por unidade de área e potencial osmótico em pleno turgor (MALES; GRIFFITHS, 2017). Além disso, esses grupos funcionais são indicadores indiretos de maior ou menor dependência da raiz para absorção de água. Em termos simples, as epífitas atmosféricas CAM são as mais adaptadas para absorção através de tricomas foliares, enquanto plantas C₃ terrestre são as mais dependentes da absorção das raízes (BENZING, 2000; MALES; GRIFFITHS, 2017).

As inflorescências de bromélias constituem um habitat complexo constituído por vários compartimentos como sépalas, pétalas e brácteas florais. Essas estruturas variam muito em toda a família em formas e cores, além do néctar que varia em

composição e concentração de açúcar (BENZING, 2000; KRÖMER et al., 2008). A família Bromeliaceae possui um amplo espectro de polinizadores, desde insetos, aves, morcegos e até roedores, mas é em grande parte polinizada por beija-flores (SIQUEIRA-FILHO; LEME, 2006). Algumas bromélias têm inflorescências discretas, mas a grande maioria tem inflorescências coloridas, grandes e vistosas, além de produzir grandes quantidades de néctar, uma característica incomum em angiospermas e que explica sua associação predominante com polinizadores vertebrados (BENZING, 2000; BERNARDELLO, 1992; WOLOWSKI; FREITAS, 2015).

3. MÉTODOS

3.1. Caracterização da área de estudo

As amostras de folhas foram coletadas de bromélias da espécie *Bromelia laciniosa* Mart. Ex Schult. & Schult.f (Bromeliaceae, Bromelioideae), uma espécie endêmica da Caatinga (FERREIRA; FABRICANTE; SIQUEIRA-FILHO, 2015). É usada na medicina popular e suas folhas, flores e frutos utilizados no tratamento de cólicas infantis, diarreia, febre, icterícia, caspa e hepatite (DE ALBUQUERQUE et al., 2007). As coletas foram realizadas ao longo de um transecto de 5 × 200 metros na Reserva Particular do Patrimônio Natural Tocaia (RPPN Tocaia), no município de Santana do Ipanema, Alagoas-Brasil, numa área de Caatinga (9°23'08.9"S e 37 °15'22.8"W). Nessa região, as chuvas ocorrem de forma irregular, com volume anual de 500 a 700 mm. A RPPN Tocaia possui uma área de aproximadamente 20 hectares e altitude máxima de cerca de 400 metros, é composta principalmente por uma fisionomia arbórea. O clima predominante é do tipo semiárido Bsh (Köppen).

3.2. Coleta do material

Foram realizadas 12 coletas. Destas, seis no período seco (outubro, novembro e dezembro de 2020 e de 2021) e seis no período chuvoso (maio, junho e julho de 2021 e de 2022). Em cada coleta, cinco indivíduos de bromélias adultas e aparentemente saudáveis foram amostrados aleatoriamente ao longo do transecto

previamente estabelecido. De cada indivíduo de bromélia coletado foram retiradas 3 folhas escolhidas ao acaso, totalizando 60 indivíduos e 180 folhas coletadas durante as 12 coletas. As folhas foram armazenadas em sacos plásticos estéreis e transportadas em temperatura ambiente (27 ± 2 °C) por aproximadamente 4 h até o Laboratório de Diversidade Molecular da UFAL. No laboratório, as folhas foram armazenadas a 10-15 °C e o processamento foi realizado em no máximo 24 h após a coleta do material em campo.

3.3. Processamento das amostras

O processamento das amostras seguiu metodologia modificada a partir de Landell, Mautone e Valente (2006). As folhas passaram por uma lavagem inicial com água destilada estéril para remoção de possíveis contaminantes como poeira e outros materiais exógenos. Em seguida, contemplando as 3 folhas coletadas de cada planta, fragmentos foliares entre 5 e 20 cm² foram cortados aleatoriamente até se obter uma área total de 120 cm². Sequencialmente a esse procedimento, a massa dos 120 cm² de folha foi aferida para cada amostra. Os fragmentos das folhas foram alocados em um frasco Erlenmeyer de volume igual a 250 mL contendo 75 mL de água destilada estéril. Em seguida esse frasco foi agitado por 30 min a 180 RPM e temperatura ambiente. Após a agitação, o produto de lavagem foi recolhido em um frasco estéril e os fragmentos de folhas preservados no frasco Erlenmeyer. Após retirar a água da primeira lavagem, adicionou-se 75 mL de solução de Tween 80 0,5% e o material foi novamente agitado sob o mesmo tempo, rotação e temperatura. O produto da segunda lavagem foi então agregado ao da primeira, formando assim um *pool*. Uma fração de 100 µL deste *pool* foi semeada em duplicata em placas de Petri contendo meio Ágar YM modificado (0,3% extrato de levedura, 0,3% extrato de malte, 0,5% peptona bacteriológica, 1% glicose, 2% ágar, 0,04% cloranfenicol, pH 4,0) nas concentrações de 10⁰ e 10⁻¹. As placas foram incubadas a 25-28 °C e observadas diariamente durante 10 dias para o isolamento das leveduras.

3.4. Obtenção e armazenamento dos isolados

Colônias representando morfotipos distintos de leveduras foram delimitadas considerando características como cor, margem, textura, forma e elevação da colônia. Representantes desses morfotipos foram isolados em meio Ágar YEPD (2% glicose, 1% peptona bacteriológica, 0,5% extrato de levedura e 2% ágar). Para verificar se as colônias isoladas eram realmente de leveduras, utilizou-se microscopia óptica de luz nos aumentos de 40 e 100x. Todas os isolados de levedura encontrados durante os 10 dias de observação foram preservadas em 1) meio Ágar GYMP (2% de glicose, 2% de extrato de malte, 0,5% de extrato de levedura, 0,2% de fosfato de sódio monobásico e 2% de ágar), 2) caldo GYMP modificado (1% de glicose, 1% de extrato de malte, 0,25% de extrato de levedura e 0,1% de fosfato de sódio monobásico) e 3) caldo GYMP contendo volume final de 30% de glicerol. Os meios 1 e 2 foram armazenados entre 5 e 15 °C e o meio 3 entre -10 e -20 °C.

3.5. Análise de traços funcionais

Traços metabólicos como a capacidade de fermentar glicose, assimilar diferentes fontes de carbono, produzir enzimas extracelulares e surfactantes/emulsificantes foram avaliadas para se criar um perfil funcional para cada isolado.

3.5.1. Fermentação de glicose

A capacidade dos isolados de leveduras fermentar glicose foi avaliada utilizando tubos de ensaio com meio básico para fermentação de glicose (0,75% de peptona bacteriológica, 0,45% de extrato de levedura e 2% glicose) (KURTZMAN et al., 2011). Tubos de *Durham* invertidos foram posicionados dentro dos tubos de ensaio, os isolados que fermentaram glicose produziram gás carbônico que se acumulou no interior dos tubos de *Durham*, indicando resultado positivo para o teste. Esse teste foi graduado em grau 1, 2 ou 3 dependendo de quanto gás se armazenou no tubo, sendo que no grau 3 o tubo estava totalmente preenchido por gás. O teste durou 21 dias, com leituras diárias nos primeiros cinco dias, e semanais a partir do sétimo dia.

3.5.2. Assimilação de fontes de carbono

O teste de assimilação de fontes de carbono foi aplicado para determinar a capacidade das leveduras utilizarem diferentes fontes de carbono no crescimento. Para o inóculo, células de leveduras foram diluídas em 2 mL de água destilada estéril até atingir uma concentração aproximada de 10^5 células/mL, o que equivale ao grau 1 do cartão de Wickerham. Após isso, foram incubadas em temperatura ambiente (27 ± 2 °C) entre 24-48 h para que suas reservas fossem exauridas, diminuindo a chance de um falso positivo. O perfil de assimilação foi avaliado semanalmente, durante 21 dias incubados a 22-25 °C seguindo Kurtzman et al. (2011). As fontes de carbono utilizadas foram celobiose, D-arabinose, galactose, glicose, glicerol, inulina, rafinose, ramnose e xilose. Os meios de cultura para os testes de assimilação das fontes de carbono consistiam em 0,67% de *Yeast Nitrogen Base* (YNB), 2% de ágar ultrapuro Difco® e 0,5% da fonte de carbono especificada (exceto, Raffinose 1%).

3.5.3. Produção de hidrolases extracelulares

A capacidade dos isolados produzirem seis enzimas hidrolíticas extracelulares foi avaliada: amilase, caseinase, celulase, esterase, lipase e pectinase. O diâmetro dos halos de hidrólise e das colônias de cada isolado foram medidos para avaliar a produção de enzimas extracelulares através da atividade enzimática (pz), aplicada a equação modificada do $pz = \left| \left(\frac{Dc}{Dc+Dh} \right) - 1 \right|$. Onde pz representa a atividade enzimática, Dh e Dc são, respectivamente, o diâmetro do halo de hidrólise e o diâmetro da colônia. A padronização do inóculo para avaliação da produção enzimática foi realizada utilizando a turbidez. As células dos isolados foram diluídas em 2 mL de água destilada estéril até atingir uma concentração aproximada de 10^5 células/mL (i.e., grau 1 do cartão de Wickerham). Em seguida, o inóculo foi aplicado em forma de ponto no meio de cultura utilizando o carimbo replicador (método de *replica plating*). As placas de petri contendo os meios para cada enzima foram analisadas após um período de 7-10 dias de incubação a 25-28 °C.

A atividade amilolítica foi avaliada usando meio de cultura Ágar amido modificado (0,5% amido solúvel, 0,5% peptona bacteriológica, 0,5% extrato de

levedura, 0,05% sulfato de magnésio, 0,001% sulfato de ferro, 0,001% cloreto de sódio, 1,5% ágar). Após o período de incubação, uma solução de lugol a 1% foi adicionada à placa para visualização do halo de hidrólise. O isolado com atividade amilolítica apresentou um halo que não é corado por lugol e permanece claro (BUZZINI; MARTINI, 2002; CARRASCO et al., 2012; MAUTONE et al., 2010).

A atividade celulolítica foi avaliada usando o meio Ágar CMC modificado (0,5% carboximetilcelulose (CMC), 0,1% nitrato de sódio, 0,1% fosfato monobásico de potássio, 0,1% cloreto de potássio, 0,05% sulfato de magnésio, 0,05% extrato de levedura, 0,1% glicose e 1,7% ágar). Para visualização do resultado, adicionou-se solução de vermelho Congo 0,1% às placas, que foram deixadas em repouso com a solução por 40 min. Em seguida, as placas foram lavadas com solução de NaCl 1M. Em caso de resultado positivo para hidrólise de CMC, a formação de um halo alaranjado foi observada (BUZZINI; MARTINI, 2002; CARRASCO et al., 2012).

A atividade de esterase foi avaliada em meio Ágar Tween 80 modificado (2,5% tween 80, 0,1% glicose, 1% peptona bacteriológica, 0,5% cloreto de sódio, 0,1% cloreto de cálcio e 2% ágar). Após o período de incubação, os isolados que hidrolisaram o substrato apresentaram um halo formado por um precipitado esbranquiçado (BUZZINI; MARTINI, 2002; CARRASCO et al., 2012).

A atividade da protease foi avaliada em meio Ágar Caseína modificado (1% caseína, 0,5% glicose e 2% ágar pH 7,0). Após o período de incubação, os isolados produtores de caseinase apresentaram um halo esbranquiçado (BUZZINI; MARTINI, 2002; CARRASCO et al., 2012).

A atividade da pectinase foi avaliada usando o meio Ágar Pectina modificado (0,67% de base de nitrogênio de levedura (YNB), 1% de pectina cítrica, 1% de glicose e 1,8% de ágar, pH 7,0). Após o período de incubação, solução de brometo de hexadeciltrimetilamônio (CTAB) a 1% foi adicionada à placa. Os isolados produtores apresentaram um halo mais translúcido ao redor da colônia (BUZZINI; MARTINI, 2002; CARRASCO et al., 2012).

3.5.4. Produção de surfactantes/emulsificantes

A atividade tensoativa foi avaliada por meio do teste de Parafilm® M. Os isolados foram cultivados durante 48 h em caldo YEPD em temperatura ambiente (27 ± 2 °C), em seguida 20 µL do cultivo foi adicionado em triplicata sobre fragmentos de parafilme M e mantidos em repouso por 1 min. Após esse tempo, as gotas formadas pela aplicação do cultivo na superfície hidrofóbica do parafilme tiveram seus diâmetros aferidos. Além disso, o formato da gota foi categorizado conforme seu espalhamento. Foram utilizados três controles: caldo YEPD estéril, água destilada estéril (ambos controles negativos) e solução de SDS 1% (controle positivo). Apresentar um diâmetro maior e uma gota mais plana são indícios que a tensão superficial da gota foi diminuída, indicando atividade tensoativa.

Para verificar a atividade bioemulsificante, foi usado o índice de emulsão com estabilidade após 24 h (IE_{24}). Os isolados foram cultivados em 2 mL de caldo YEPD por 48 h, à temperatura ambiente (27 ± 2 °C). Após o crescimento, 2 mL de querosene foi adicionada ao cultivo e agitada com auxílio de um vórtex por 2 min e mantida em repouso por 24 h. Em seguida foram aferidos a altura total do cultivo e a altura da emulsão em centímetros. O IE_{24} foi obtido a partir da equação $IE_{24} = \left(\frac{ALT_{emulsão}}{ALT_{total}} \right) \times 100$, ou seja, a proporção da altura da emulsão em relação à altura total do cultivo, em porcentagem.

3.6. Identificação molecular dos isolados

3.6.1. Extração de DNA genômico

Para a identificação molecular dos isolados, o DNA genômico foi extraído utilizando o protocolo para preparação em pequena escala de DNA de leveduras modificado, originalmente proposto por Sambrook e Russel (2001). Para tal, inicialmente as culturas puras dos isolados foram inoculadas em caldo YEPD durante 2-3 dias a 22-25°C. Após, as células foram precipitadas em tubos de microcentrifuga de 1,5 mL a 10000 x g durante 5 min. Após a centrifugação, o sobrenadante foi descartado e adicionou-se 1 mL de água destilada estéril e homogeneizado. Em seguida, as células

foram novamente centrifugadas sob mesma rotação e tempo. Ao final, o sobrenadante foi novamente descartado.

Foi adicionado ao precipitado de células obtidos no passo anterior 500 µL de solução de lise (0,15 M de NaCl, 50 mM de Tris-HCl pH 8, 10 mM de ácido etilenodiamino tetra-acético (EDTA) pH 8 e 2% de dodecil sulfato de sódio-SDS) e homogeneizado por agitação em vortex. Em seguida, os tubos contendo o homogeneizado foram incubados em banho-maria a 65°C por 1 h. Logo após retirar os tubos, adicionou-se imediatamente 200 µL de acetato de potássio 5M pH 4.8, a reação foi homogeneizada por 30 s e incubada em banho de gelo por 30 min. Em seguida, os tubos foram centrifugados a 14000 x g por 5 min e o sobrenadante foi transferido para outro tubo de microcentrífuga. O tubo foi novamente centrifugado a 14000 x g por 5 min e o sobrenadante foi transferido para outro tubo sem utilizar micropipeta. Segundo, adicionou-se 600µl de isopropanol absoluto conservado a -20 °C. Os tubos foram gentilmente agitados por inversão durante 5 min e mantidos a -20 °C de um dia para outro. Os tubos foram então centrifugados a 14000 x g durante 20 min e o sobrenadante foi descartado. Um volume de 500 µl de etanol 70% conservado a -20 °C foi adicionado e os tubos foram novamente centrifugados a 14000 x g por 5 min. Ao término o sobrenadante foi descartado e o precipitado passou por secagem a temperatura ambiente (27 ± 2 °C) e foi posteriormente solubilizado em 50 µL de tampão Tris-EDTA (TE) numa proporção de 1 para 1, pH 7,4. A qualidade da extração de DNA foi aferida posteriormente em gel de agarose (0,8%) diluída em tampão TAE 1X (Solução mãe 50X = 242 g TRIS base, 57,1 mL ácido acético glacial e 100 mL de solução de EDTA 0,5 M pH 8,0, em 1 L de água destilada) (COBBAN et al., 2016).

3.6.2. PCR, sequenciamento e análise das sequências

As regiões D1/D2 do gene 26S LSU do rRNA (~600pb) foram amplificadas pela técnica de reação em cadeia da polimerase (PCR) utilizando os oligonucleotídeos iniciadores NL-1 (5'- GCA TATC AAT AAG CGG AGG AAA AG -3') e NL-4 (5'- GG TCC GTG TTT CAA GAC GG -3') (FELL et al., 2000; KURTZMAN; ROBNETT, 1998). Para amplificar a região do espaçador interno transcrita (*Internal Transcrit Spacer - ITS*) do rRNA foram utilizados os oligonucleotídeos iniciadores ITS-1 (5'-TCC GTA GGT GAA

CCT GCG G-3') e ITS-4 (5'-TCC TCC GCT TAT TGA TAT GC-3') (FELL et al., 2000; SCHOCH et al., 2012). Os parâmetros da reação foram estabelecidos seguindo Landell et al. (2010) e o produto da reação foi observado em uma eletroforese usando gel de agarose (1%) diluído em tampão TAE 1X. Os produtos (25 µL) foram purificados através de uma nova precipitação utilizando isopropanol absoluto (20 µL) e acetato de sódio 3M (5 µL). O material foi centrifugado durante 20 min a 13.000 G, o sobrenadante foi descartado e substituído por etanol 70% (50 µL), centrifugado novamente e mais uma vez o sobrenadante foi descartado. Após a secagem do precipitado, o material foi suspenso em 15 µL de água ultrapura. Para o sequenciamento foi utilizado o método de Sanger através do protocolo do Laboratório de Genética Molecular Humana da Universidade Federal de Pernambuco em um sistema de sequenciamento automatizado ABI 3130 *Genetic Analyzer* utilizando o polímero *BigDye* v3.1 e POP7 (*Life Technologies*). Os consensos das sequências foram gerados no software *Staden Package* (STADEN et al., 2000) e MEGA X. Em seguida, as sequências obtidas foram comparadas com outras depositadas no banco de dados do *GenBank* no site do *National Center for Biotechnology Information* (NCBI) utilizando a ferramenta *Basic Local Alignment Search Tool* (BLAST) e o algoritmo BLASTn (ALTSCHUL et al., 1997).

3.7. Caracterização das possíveis espécies novas

Para a caracterização das possíveis espécies novas de leveduras foi empregado o modelo de taxonomia polifásica, incluindo caracterização e análises moleculares e fenotípicas. A informação molecular foi obtida através do sequenciamento das regiões D1/D2 e ITS do rRNA (*vide* secção anterior). A caracterização fenotípica seguiu os protocolos descritos por Kurtzman et al. (2011).

3.7.1. Assimilação de fontes de carbono e nitrogênio

O processo para obtenção do padrão de assimilação de fontes de C e N dos isolados de espécies não-descritas encontradas seguiu protocolo descrito na seção 3.5.2. O meio base para as fontes de nitrogênio consistia em 1,17% de *Yeast Carbon Base* (YCB) e 2% de ágar ultrapuro Difco®, a concentração de cada fonte de nitrogênio é indicada no Quadro 1, bem como a lista de todas as fontes utilizadas.

Quadro 1- Lista de Fontes de carbono e de nitrogênio

Fontes de carbono		Fontes de nitrogênio
Glicose*	L-arabinitol	
Ácido galacturônico ^a	N-acetilglicosamina (NAG)	Peptona*,d
Ácido succínico ^a	Maltose	
Amido	Manitol	Cadaverina ^c
Celobiose	Melezitose	
Citrato	Melibiose	Creatina ^d
D-arabinose	Rafinose	
L-arabinose	Ramnose	Creatinina
Dulcitol	Ribitol	
Eritritol	Ribose	Etilamina ^c
Galactose	Sacarose ^b	Lisina ^e
Glicerol	Salicina	
Glucanato	Sorbitol	Nitrato ^d
Inositol	Trealose	
Inulina	Tween 20	
Lactato	Tween 80	Nitrito ^c
Lactose	Xilose	

* Controle positivo

^a O pH foi ajustado para 7,0 antes da esterilização.

^b A fonte de carbono foi adicionada após a esterilização para evitar degradação.

^c Avaliada através do método de *pour plate* (concentração final da fonte 0,064%).

^d Concentração final da fonte 0,078%.

^e Concentração final da fonte 0,056%.

3.7.2. Formação de compostos amiloïdes extracelulares

A capacidade dos isolados formarem compostos amiloïdes extracelulares foi verificada após 21 dias de crescimento em 22-25 °C em meio básico para glicose (0,67% de YNB, 2% de ágar e 0,5% glicose). Após esse período, uma solução de lugol 1% foi adicionada sobre as colônias. Os isolados que produziram compostos amiloïdes formaram colônias com coloração azul e verde escuros (KURTZMAN et al., 2011).

3.7.4. Testes de produção de urease e reação ao Diazônio Azul B (DBB)

A avaliação da produção de urease e da capacidade do isolado de reagir ao corante DBB foram utilizados como indicadores presuntivos de afinidade aos filos Ascomycota ou Basidiomycota (KURTZMAN et al., 2011). Para o teste foi usado ágar ureia inclinado em tubos de ensaio (1,17% YCB, 1% de ureia, 2% de ágar e fucsina ácida), a ureia previamente esterilizada foi adicionada apenas após a esterilização pois é termossensível. Colônias puras foram semeadas no meio e observadas diariamente ao longo de três dias. Os isolados que foram capazes de produzir urease, hidrolisam a ureia produzindo amônia que, por sua vez, basifica o meio de cultura e neutralizaram a fucsina ácida, fazendo o meio perder a coloração rosa-arroxeadas e ficar esbranquiçado. Após os três dias de leitura do teste de urease, os tubos foram incubados a 60 °C durante 16 h. Após esse período, 1 mL de solução de DBB diluído em Tris-HCl 1 M pH 7,0 foi adicionada aos tubos. Os isolados foram considerados positivos para a reação quando a colônia apresentou uma coloração avermelhada em contato com a solução e, portanto, foram considerados com afinidade basidiomicética.

3.7.5. Testes de tolerância

3.7.5.1. Tolerância osmótica

Para verificar a tolerância osmótica, os isolados foram semeados e Ágar YEPD modificado contendo 50% de glicose (2% de ágar, 50% de glicose, 1% de peptona, 0,5% de extrato de levedura). O meio foi autoclavado durante 10 min a 110 °C. Além deste, YEPD modificado contendo 5 e 10% de NaCl também foi usado para verificar a tolerância osmótica dos isolados (KURTZMAN et al., 2011).

3.7.5.2. Ácido acético

Foi verificada a tolerância dos isolados de crescer em meio contendo 1% de ácido acético. Para tal, foi usado meio YEPD modificado contendo 1% de ácido acético glacial (KURTZMAN et al., 2011). O ácido só foi adicionado após esterilização do meio de cultura base.

3.7.5.3. Resistência a cicloheximida

A capacidade dos isolados em tolerar crescimento em meio contendo ciclohexamida foi verificada em meio básico para glicose (0,67% de YNB e 0,5% de glicose) com concentração final de 0,1 e 0,01% de cicloheximida (KURTZMAN et al., 2011). A cicloheximida foi acrescida no meio após a esterilização pois é termossensível. A leitura foi realizada após 7 dias de incubação a 22-25 °C.

3.8. Caracterização micromorfológica

Os isolados foram caracterizados morfologicamente seguindo Kurtzman et al. (2011). Para verificar a possível produção de pseudohifas, hifas verdadeiras e/ou estruturas sexuais, os isolados foram semeados em Ágar Batata Dextrose (BDA), Ágar Fubá, Ágar Extrato de Malte e Ágar YEPD. Os isolados foram incubados a 22-25 °C e observados semanalmente por 21 dias.

Publicado em *Yeast Journal (ecoYeast reviews)* DOI: 10.1002/yea.3804

4. CAPÍTULO 1 - Different plant compartments, different yeasts: the example of the bromeliad phyllosphere

Ciro Ramon Félix^{1,2}, Bruno Emanuel da Silva Nascimento¹, Patrícia Valente³, Melissa Fontes Landell^{1*}

¹ Universidade Federal de Alagoas, Instituto de Ciências Biológicas e da Saúde, Maceió – AL, Brazil.

² Programa de Pós-graduação em Diversidade Biológica e Conservação nos Trópicos, Universidade Federal de Alagoas, Maceió – AL, Brazil.

³ Universidade Federal do Rio Grande do Sul, Departamento de Microbiologia, Imunologia e Parasitologia, Porto Alegre – RS, Brazil.

***Correspondence:** Melissa Fontes Landell, Setor de Genética - ICBS, Universidade Federal de Alagoas, Av. Lourival Melo Mota, s/n, Tabuleiro dos Martins, CEP: 57072-900, Maceió - AL, Brasil, Tel. (+55) 82-32141995

(e-mail: melissa.landell@gmail.com)

ORCID of the authors:

Ciro R. Félix - 0000-0001-5087-8872

Bruno Emanuel da Silva Nascimento - 0000-0001-6690-0595

Patrícia Valente - 0000-0001-5374-0991

Melissa Fontes Landell - 0000-0001-6848-0803

Take Away:


- Bromeliads are a great model for studies of yeast-plant interactions.
- More than 180 yeast species have already been recorded in bromeliads.
- Bromeliads compartments have distinct yeast communities.
- Bromeliad subfamilies and functional types were inconclusive in yeast structuring.
- Yeasts from bromeliads can find industrial application in biotechnology.

Abstract

The plant phyllosphere is one of the largest sources of microorganisms, including yeasts. In bromeliads, the knowledge of yeasts is dispersed and still incipient. To understand the extent of our knowledge on the subject, this review proposes to compile and synthesize existing knowledge, elucidating possible patterns, biotechnological and taxonomic potentials, bringing to light new knowledge, and identifying information gaps. For such, we systematically review scientific production on yeasts in bromeliads using various databases. The results indicated that the plant compartments flowers, fruits, leaves, and water tank (phytotelma) have been studied when focusing on the yeast community in the bromeliad phyllosphere. More than 180 species of yeasts and yeast-like fungi were recorded from the phyllosphere, 70% were exclusively found in one of these four compartments and only 2% were shared among all. In addition, most of the community had a low frequency of occurrence, and approximately half of the species had a single record. Variables such as bromeliad subfamilies and functional types, as well as plant compartments, were statistically significant, though inconclusive and with low explanatory power. At least 50 yeast species with some biotechnological potentials have been isolated from bromeliads. More than 90% of these species were able to produce extracellular enzymes. In addition, other biotechnological applications have also been recorded. Moreover, new species have been described, though yeasts were only exploited in approximately 1% of the existing bromeliads species, which highlights that there is still much to be explored. Nevertheless, it appears that we are still far from recovering the completeness of the diversity of yeasts in this host. Furthermore, bromeliads proved to be a good ecological model for prospecting new yeasts and for studies on the interaction between plants and yeasts. In addition, the yeast community diverged among plant compartments, establishing bromeliads as a microbiologically complex and heterogeneous mosaic.

Keywords: leaf, flower, fruit, phytotelma, systematic review and Bromeliaceae.

Graphical abstract



1. INTRODUCTION

There are variations on the phyllosphere concept, nonetheless, in general, it is the entire interface between plants and the atmosphere, including leaves (phylloplane), fruits (carposphere), flowers (anthosphere), stems (caulosphere), and waterscape (phytotelma) (Bringel and Couée, 2015; Vacher et al., 2016). The phyllosphere is an extreme, dynamic, and heterogeneous environment, often oligotrophic. Besides the traditional exposure and sudden fluctuations of stress factors, such as temperature, solar radiation, oxidation, the availability of nutrients and water, another critical point is the cutin layer, which forms the plant's cuticle. The plant's cuticle protects the leaves and assists in water control. It also decreases the evaporation and leaching of metabolites, limits the availability of nutrients, and creates an extreme and adverse environment (Bringel and Couée, 2015; Vorholt, 2012). Nowadays, it is well established that plants are microbiologically structured environments, rich in diversity, and ecologically complex (Redford et al., 2010; Vorholt, 2012). Each plant compartment (e.g. leaves, flowers, fruits, etc.) can and often do have distinct microbial communities that can vary in space, time, and ontogenetically (Abdelfattah et al., 2019; Fonseca and Inácio, 2006; Liu and Howell, 2021).

Bromeliaceae is an almost exclusively neotropical plant family recognized, among other factors, for its great diversity (approximately 3,500 species, 50 genera and 8 subfamilies), high endemism and for providing several ecosystem services in support of biodiversity and water regulation (Benzing, 2000; Ladino et al., 2019). Different than other members of angiosperms, bromeliads are mostly pollinated by vertebrates, mainly hummingbirds, but there is a long list of possible pollinators which involve groups such as bats, rodents and bees (Benzing, 2000; Siqueira-Filho and Leme, 2006; Tschapka and Von Helversen, 2007). Bromeliad inflorescences are mostly large and colorful, produce large volumes of nectar, with these traits reinforcing their relationship with vertebrate pollinators (Benzing, 2000; Siqueira-Filho and Leme, 2006; Wolowski and Freitas, 2015). Furthermore, they present a wide spectrum of colonized environments, lifestyles and ecophysiological types, occupying environments ranging from tropical to temperate climate regions, from 0 to 4,000 m of altitude and from rainy to dry forests

(Benzing, 2000; Crayn et al., 2015; Ladino et al., 2019; Males and Griffiths, 2017). Another important element about Bromeliaceae is that several species can accumulate water in a structure called phytotelma, a region in the center of the rosette. This water tank contributes, for example, to the reproduction of several mosquito species, including some vectors of human diseases such as *Aedes aegypti*, *A. albopictus*, *Culex* sp. and *Haemagogus* sp. (Benzing, 2000; Ladino et al., 2019). Among the species of mosquitoes associated with bromeliads, a minority is a vector of certain human diseases (less than 6%) (Ladino et al., 2019). Bromeliads function as a microscale island ecosystem (Frank and Lounibos, 1987), and perhaps also as a refuge in dry times, mainly, but not exclusively, for aquatic or heavily water-dependent organisms.

The microbiota associated with bromeliads has been studied for over a century (Frank and Lounibos, 1987), and the associated microbial groups include bacteria, archaea, viruses, filamentous fungi, yeasts, basal fungi (Chytridiomycota) and several other microeukaryotes (Goffredi et al., 2015; Leroy et al., 2016; Louca et al., 2017; Morais et al., 2020). In addition, vascular plants are one of the greatest sources of fungal diversity (Hawksworth and Lücking, 2017). For example, in the MycoBank database, the substrate with the highest number of records are leaves. These associated communities are modulated and/or regulated by a complex network of physicochemical, climatic, ecological, geographic factors and host-related factors such as senescence and genotype (Fonseca and Inácio, 2006; Vacher et al., 2016; Vorholt, 2012; Whipps et al., 2008).

Yeasts are a group formed by several fungal isolates with a convergent regulatory evolution for the maintenance of the unicellular phenotype (Nagy et al., 2014). These organisms are known to be among the first to establish themselves in nutrient-rich environments (Fonseca and Inácio, 2006; Ganter et al., 2017; Starmer and Lachance, 2011) and to occupy a wide spectrum of habitats distributed across the planet, whether associated with plants and animals, water, soil or the atmosphere (Rosa and Péter, 2006; Starmer and Lachance, 2011). Yeasts have been used for millennia in beverage and food fermentation processes, with the most well-known species being *Saccharomyces cerevisiae* - the “baker’s yeast” (Rosa and Péter, 2006). Although some

yeast species are human pathogens, such as *Candida albicans* and *Cryptococcus neoformans* (Rosa and Péter, 2006), they represent a small portion of this group. For instance, several yeast types associated with plants present antagonistic activity to pathogens and are capable of producing plant hormones that stimulate plant growth (Buck, 2002; Limtong and Koowadjanakul, 2012; Marques et al., 2021). It has recently been estimated that there are currently approximately 1,600 described yeast species (Vu et al., 2016). However, this number may probably exceed 2300 species (Yurkov et al., 2021).

For about 30 years several generations of scientist have been studying yeasts associated with bromeliads. During this period, most studies have focused on yeast diversity and on the description of new species. Thus, through a systematic review we collected almost three decades of data on yeasts in bromeliads, we analyzed the diversity, frequency and structure of yeasts in bromeliads in different plant compartments (leaves, flowers, fruits and tank water) and considered other variables such as the bromeliad subfamily and functional type. Moreover, to understand the extent of our knowledge on the subject, we compile and synthesize existing knowledge, besides bringing to light new knowledge. In addition, we synthesized knowledge on new species and the biotechnological potential of bromeliad yeasts.

2. METHODS

2.1. Search in databases and data collection

We explored four databases to compile the bibliographic material used in this study: PubMed (<https://pubmed.ncbi.nlm.nih.gov/>), Scielo (<https://www.scielo.org/>), Scopus (<https://www.scopus.com/>) and Web of Science (<https://www.webofscience.com/>). The search terms (yeast* OR micro-fung* OR microfung* OR "unicellular fung*") AND (bromelia* OR pineapple OR ananas) were applied in a standardized way, searching for titles, abstracts and keywords on papers throughout the months of September and October 2021. The results were downloaded into bibtex files and concatenated into a single file. The online tool texmed

(<https://www.bioinformatics.org/texmed/>) was used to access the bibtex file from PubMed. In addition, manual searches were carried out on Google Scholar (<https://scholar.google.com/>).

2.2. Bibliography screening: inclusion and exclusion criteria

The papers files were imported into the R statistical software (R Team, 2016) using the revtools package (Westgate, 2019), duplicate references were removed. Following the reading of abstracts or complete material, the next step involved verifying whether the references met the following criteria: (1) Only studies focused on yeasts associated with bromeliads, whether in the fields of biodiversity, ecology, taxonomy, or application in biotechnological processes. However, articles that only used yeast counts as an indicator of food (pineapple fruit) contamination were excluded. (2) Non-English language papers were disregarded. (3) Only papers published in scientific journals were considered. Sources such as publications in congresses, symposia, or books were not incorporated. (4) Papers focused only on yeasts from industrialized juices and wines were also excluded. (5) All studies should be available online.

2.3. Research questions

- What is the community of yeasts associated with bromeliads, how is it structured among the plant compartments and what are the potential causal actors that shape yeast communities?
- What new yeast species associated with bromeliads have been described?
- What are the main biotechnological properties found in yeast from bromeliads?

2.4. Data extraction, tabulation, and synthesis

The units used were defined considering the scope of the study on a sample of bromeliad species, the compartments and sampling sites, as outlined in Figure 1. Thus, our work unit consisted of each Delimited Unit of Analysis (DUA). The DUAs were used to incorporate part of the variation in various information layers of the analyzed papers (e.g. hosts, substrates and sampling sites). The studies available in the literature have often different methodologies and vary in numbers of samples, hosts, compartments and sampling sites, DUAs are a way to standardize the analysis, although

there are biases involved. The papers were grouped into four categories according to their work focus (each study could be placed in more than one category):

- 1.** Diversity and ecology - for articles with a focus on describing the local diversity of yeasts, which may or may not relate to ecological issues.
- 2.** Taxonomy - papers focusing on the description of new species or taxonomic groups.
- 3.** Biotechnology - studies that sought to apply yeasts in some biotechnological process and/or prospect some product or service of these isolates that could have some biotechnological application.
- 4.** Others- when the paper did not fit into any of the previously mentioned categories.




Fig. 1- Schematic drawing exemplifying how the units used were delimited. The delimited units of analysis (DUAs) would be related to the number of bromeliad species, compartments, and sampling sites from each paper.

To analyze the community of yeasts in bromeliads, the following information was collected from the papers: the lists of species, host name, compartment type (flowers, fruits, leaves or water tank), yeast identification method (molecular or phenotypic), culture medium isolation, collection site and biotechnological products or services produced/provided by yeasts. Yeast species with dubious, inconclusive, or incomplete identification were disregarded (except for biotechnology studies, where the identification at the genus level was also computed). The names of yeast species were revised using the MycoBank database (<https://www.mycobank.org/>) and specialist bibliography, e.g. (Kurtzman, 2011; Liu et al., 2015; Wang et al., 2016, 2015). From the correct names of yeast species, other taxonomic levels were assigned (Phylum, Subphylum, Class, Order, Family and Genus). The taxonomically correct names of bromeliads were verified on the Global Biodiversity Information Facility-GBIF (<https://www.gbif.org/>) and the subfamilies assigned through consultation on the Bromeliad Society International- BSI (<https://www.bsi.org/new/taxonomy/>). Furthermore, functional types were assigned to bromeliads in accordance with (Males and Griffiths, 2017).

2.5. Statistical analysis

Alpha-diversity indices and estimators (Simpson's index, Dominance index and Chao1) and Whittaker beta-diversity were calculated in the PAST software (Hammer et al., 2001), using the frequency of occurrence of species in DUAs (see the previous topic) in bromeliad compartments. The frequency of occurrence was calculated using the expression $F_{oc}=(f\times 100)/D$, where ' F_{oc} ' is the frequency of occurrence of the group in question (e.g. species, genus, etc), 'f' is the count of the occurrence of the group in the DUAs (in total or in a specific compartment), and 'D' is the total number of the DUAs considered (for the entire phylosphere D=164, for flowers D=21, fruits D=18, leaves D=100 and water tanks D=25). An analysis of variance (ANOVA) was performed to verify whether there was a significant difference among the alpha-diversity indicators cited among the plant compartments. In addition, the normality of the data was verified

by applying the Shapiro-Wilk normality test, while Tukey's multiple comparisons test was applied afterward to verify the pair-by-pair difference between the compartments, with all these analyses being performed in the R software (R Team, 2016). The collector curve was also plotted using the frequency of species in the DUAs in the different compartments. The rarefaction and extrapolation (by Chao1) methods were used with 100 bootstrap pseudo-replications. This analysis and the design of the figure were carried out in R software using the packages iNEXT (Hsieh et al., 2016) and ggplot2 (Wilke, 2016).

The BETADISPER analysis, 'Multivariate homogeneity of groups dispersions (variances)', in R software was carried out to verify the similarity of the sample units dispersion between functional types (CAM tank-epiphytes, CAM atmospheric epiphyte, CAM terrestrial and C₃ tank-epiphytes), subfamilies (Bromelioideae, Tillandsioideae, and Pitcairnioideae), compartment types (flowers, fruits, leaves, and water tank) and identification methods (phenotypic or molecular). When a variable with more than two levels was tested, we used the permutation test of the homogeneity of multivariate dispersions (PERMDISP) to verify the paired significance between the groups. In this regard, the Vegan package (Dixon, 2003) was used to generate a Principal Coordinate Analysis (PCoA) applying Jaccard distance and a Multivariate Permutation Analysis (PERMANOVA), employing with 999 permutations to verify the probability of the evaluated groups (e.g. functional types) to be found at random. These data were plotted using the ggplot2 package and the difference between the centroid distances was verified with an ANOVA and Tukey's test. For the BETADISPER, PERMANOVA and PERMDISP, only the DUAs from the 20 papers in the category Diversity and ecology were used. This specific data was used as adding studies from the other categories could bias the heterogeneity of the scope of the papers.

3. RESULTS

A total of 665 papers were identified from all databases. Following the removal of duplicate papers, 383 unique references remained. After applying all filters

and criteria, 43 studies published between 1993 and 2021 were selected to be reviewed (Table 1 and Supplementary Figure 1). Twenty papers were grouped under the category Diversity and ecology, nineteen in the category Taxonomy, fourteen in the category Biotechnology, and four in the category Other. From all studies, 164 Delimited Units of Analysis (DUAs) were established (see method).

3.1. Bromeliad species and yeasts

Forty-four bromeliad species identified were recorded, most within the subfamily Bromelioideae (70%), followed by Tillandsioideae (30%). Representatives of Pitcairnioideae were identified, but none with a conclusive identification, being named only as *Dyckia* sp. and *Encholirium* spp. Furthermore, four functional types were recorded in these bromeliads: C₃ tank-epiphytes (15%), CAM atmospheric epiphyte (15%), CAM tank-epiphytes (46%) and CAM terrestrial (24%). The bromeliads with the highest yeast richness recorded were *Vriesea minarum* L.B. Sm. (n = 46 species), *Ananas comosus* (L.) Merr. (n = 44) and *Bromelia karatas* L. (n = 41).

3.2. Yeasts from bromeliad phyllosphere

From all studies in the sample, 181 yeasts and yeast-like species from four bromeliad compartments were mentioned: flowers, fruits, leaves, and water tank (Table 2). These species were grouped into 88 genera, 36 families, 20 orders, 8 classes and 5 subphyla of Basidiomycota and Ascomycota (Tables 2 and 3). Approximately 13% (n = 24) of the species found belong to the genus *Candida*, followed by *Papiliotrema* (n = 9; 5%), *Rhodotorula* (n = 9; 5%), and *Hannaella* (n = 8; 4%). Other prominent groups are the families Saccharomycetaceae (n = 27; 15%), Sporidiobolaceae (n = 16; 9%) and Ustilaginaceae (n = 12; 7%), which together group 30% of the species. Similarly, about 80% of all yeast species recorded are inserted in four orders, as follows: Saccharomycetales (n = 69; 38%), Tremellales (n = 46; 25%), Sporidiobolales (n = 16; 9%) and Ustilaginales (n = 14; 8%). Two-thirds of the species belong to Saccharomycotina and Agaricomycotina subphyla. Overall, the Basidiomycota phylum had more species than the Ascomycota (58 and 42%, respectively).

Most of the yeast species showed a low occurrence, as 46% were singletons (species occurring in only one DUA) and 65% were singletons or doubletons. Furthermore, about 97% of yeast species were present in less than 10% of the DUAs. The only species with a frequency $\geq 10\%$ were *Papiliotrema laurentii* (13%), *Papiliotrema flavescens* (12%), *Meyerozyma guilliermondii* (12%), *Candida intermedia* (11%), *Pseudozyma hubeiensis* (11%) and *Aureobasidium pullulans* (10%) (Figure 2A). Combined, at least one of these five species occurred in more than a third of all units analyzed. Regarding the most frequent genera, only eight of the 88 registered had a frequency of occurrence greater than 10%: *Papiliotrema* (34%), *Candida* (21%), *Aureobasidium* (15%), *Meyerozyma* (15%), *Rhodotorula* (15%), *Carlosrosaea* (12%), *Hannaella* (12%) and *Pseudozyma* (10%). At least one of these genera occurred in 73% of all sampled units (Figure 2B).

In addition, 52% of the species occurred in only one bromeliad species, while only ten yeast species had a frequency of occurrence higher or equal than 20% on 44 bromeliad species: *P. laurentii* (39% occurrence), *C. intermedia* (36%), *P. flavescens* (31%), *Ps. hubeiensis* (30%), *Naganishia albida* (27%), *Papiliotrema leoncinii* (25%), *A. pullulans* (23%), *Hannaella pagnoccae* (23%), *M. guilliermondii* (25%) and *Carlosrosaea hohenbergiae* (20%). Combined, at least one of these species occurred in more than 90% of the bromeliad species recorded.

Table 1- List of articles inserted and analyzed in the review.


Paper title	Defined category	Reference
Yeast and coliform bacteria of water accumulated in bromeliads of mangrove and sand dune ecosystems of southeast Brazil	Diversity	Hagler et al., 1993
A preliminary note on yeast communities of bromeliad-tank waters of Rio de Janeiro, Brazil	Diversity	Araujo et al., 1998
The influence of air pollution on the phyllosphere microflora composition of <i>Tillandsia</i> leaves (Bromeliaceae)	Biotechnology and Diversity	Brighigna et al., 2000
Microbial antagonists control postharvest black rot of pineapple fruit	Biotechnology and Diversity	Reyes et al., 2004
<i>Candida bromeliacearum</i> sp. nov. and <i>Candida ubatubensis</i> sp. nov., two yeast species isolated from the water tanks of <i>Canistropsis seidelii</i> (Bromeliaceae)	Taxonomy	Ruivo et al., 2005
Biodiversity of yeasts associated to bromeliads in Itapuã Park, Viamão/RS	Diversity	Landell et al., 2006
<i>Farysizyma</i> gen. nov., an anamorphic genus in the Ustilaginales to accommodate three novel epiphytic basidiomycetous yeast species from America, Europe and Asia	Taxonomy	Inácio et al., 2008
<i>Cryptococcus bromelialarum</i> sp. nov., an orange-coloured basidiomycetous yeast isolated from bromeliads in Brazil	Taxonomy	Landell et al., 2009
Autochthonous yeasts associated with mature pineapple fruits, freshly crushed juice and their ferments; and the chemical changes during natural fermentation	Biotechnology and Diversity	Chanprasartsuk et al., 2010
Comparative analysis of fungal growth in commercially and laboratory prepared fruit juices – Using orange and pineapple as a case study	Diversity	Udota and Urua, 2010
First report of bacterium and yeasts associated with pineapple fruit collapse in Espírito Santo State, Brazil	Diversity	Korres et al., 2010
Taxonomic structure of the yeasts and lactic acid bacteria microbiota of pineapple (<i>Ananas comosus</i> L. Merr.) and use of autochthonous starters for minimally processing	Diversity and Other	Di Cagno et al., 2010
<i>Candida aechmeae</i> sp. nov. and <i>Candida vriesiae</i> sp. nov., novel yeast species isolated from the phylloplane of bromeliads in Southern Brazil	Taxonomy	Landell et al., 2010
<i>Candida krusei</i> and <i>Kloeckera apis</i> inhibit the causal agent of pineapple fusariosis, <i>Fusarium guttiforme</i>	Biotechnology and Diversity	Korres et al., 2011
Pineapple wine fermentation with yeasts isolated from fruit as single and mixed starter cultures	Biotechnology	Chanprasartsuk et al., 2012
<i>Bandaniozyma</i> gen. nov., a genus of fermentative and non-fermentative Tremellaceous yeast species	Taxonomy	Valente et al., 2012
<i>Kazachstania bromeliacearum</i> sp. nov., a yeast species from water tanks of bromeliads	Taxonomy	Araujo et al., 2012
Comparison of methods for identification of yeasts isolated during spontaneous fermentation of freshly crushed pineapple juices	Diversity and Other	Chanprasartsuk et al., 2013

<i>Kazachstania rupicola</i> sp. nov., a yeast species isolated from water tanks of a bromeliad in Brazil	Taxonomy	Safar et al., 2013
<i>Hagleromyces</i> gen. nov., a yeast genus in the Saccharomycetaceae, and description of <i>Hagleromyces aurorensis</i> sp. nov., isolated from water tanks of bromeliads	Taxonomy	Sousa et al., 2014
<i>Hannaella pagnoccae</i> sp. nov., a tremellaceous yeast species isolated from plants and soil	Taxonomy	Landell et al., 2014
The diversity and extracellular enzymatic activities of yeasts isolated from water tanks of <i>Vriesea minarum</i> , an endangered bromeliad species in Brazil, and the description of <i>Occultifur brasiliensis</i> f.a., sp. nov	Biotechnology, Diversity and Taxonomy	Gomes et al., 2015
Enzymatic activity and susceptibility to antifungal agents of brazilian environmental isolates of <i>Hortaea werneckii</i>	Diversity and Other	Formoso et al., 2015
Phenotypic and molecular diversity of <i>Meyerozyma guilliermondii</i> isolates isolated from food and other environmental niches, hints for an incipient speciation	Other	Corte et al., 2015
<i>Bullera vrieseae</i> sp. nov., a tremellaceous yeast species isolated from bromeliads	Taxonomy	Landell et al., 2015
<i>Kockovaella libkindii</i> sp. nov., a yeast species isolated from water tanks of bromeliad	Taxonomy	Gomes et al., 2016
<i>Papiliotrema leoncinii</i> sp. nov. and <i>Papiliotrema miconiae</i> sp. nov., two tremellaceous yeast species from Brazil	Taxonomy	Pagani et al., 2016
Isolation of <i>Saccharomyces cerevisiae</i> from pineapple and orange and study of metal's effectiveness on ethanol production	Biotechnology	Nasir et al., 2017
Pineapple (<i>Ananas comosus</i> L. Merr.) wine production in Angola: Characterisation of volatile aroma compounds and yeast native flora	Biotechnology and Diversity	Dellacassa et al., 2017
<i>Carlosrosaea hohenbergiae</i> sp. nov. and <i>Carlosrosaea aechmeae</i> sp. nov., two tremellaceous yeasts isolated from bromeliads in north-eastern Brazil	Taxonomy	Félix et al., 2017
<i>Occultifur plantarum</i> f.a., sp. nov., a novel cystobasidiomycetous yeast species	Taxonomy	Khunnamwong et al., 2017
<i>Pattersoniomycetes tillandsiae</i> gen. et comb. nov.: linking sexual and asexual morphs of the only known smut fungus associated with Bromeliaceae	Taxonomy	Piątek et al., 2017
Probiotic potential of yeasts isolated from pineapple and their use in the elaboration of potentially functional fermented beverages	Biotechnology and Diversity	Amorim et al., 2018
Fungal Planet description sheets: 716–784	Taxonomy	Crous et al., 2018
Fungal Planet description sheets: 868–950	Taxonomy	Crous et al., 2019
The natural lipolytic yeast <i>Candida</i> sp. Rmutsb-27 isolated from pineapple for treatment of cooking oil contaminated wastewater	Biotechnology	Tangsombatvichit et al., 2020
Richness and biotechnological potential of the yeast community associated with the bromeliad phylloplane in the Brazilian Neotropical Forest	Biotechnology and Diversity	Navarro et al., 2020

Yeast in plant phytotelmata: Is there a core community in different localities of rupestrian savannas of Brazil?	Biotechnology and Diversity	Morais et al., 2020
Changes of quality of minimally-processed pineapple (<i>Ananas comosus</i> , var. queen victoria) during cold storage: Fungi in the leading role	Diversity	Leneveu-jenvrin et al., 2020
<i>Vishniacozyma alagoana</i> sp. nov. a tremellomycetes yeast associated with plants from dry and rainfall tropical forests	Taxonomy	Félix et al., 2020
Plant growth promoting traits of yeasts isolated from the tank bromeliad <i>Vriesea minarum</i> L.B. Smith and the effectiveness of <i>Carlosrosaea vrieseae</i> for promoting bromeliad growth	Biotechnology	Marques et al., 2021
Plant endophytic yeasts <i>Pichia fermentans</i> and <i>Meyerozyma caribbica</i> improve growth, biochemical composition, haematological parameters and morphology of internal organs of premature <i>Barbonyxus gonionotus</i>	Biotechnology and Diversity	Islam et al., 2021
Behind the nectar: the yeast community in bromeliads inflorescences after the exudate removal	Diversity	Félix et al., 2021

Table 2- Number of yeast taxa found at taxonomic levels in each compartment of bromeliads.

	Species	Genera	Family	Order	Class	Subphyla	Phyla
Total bromeliads	181	88	36	20	8	5	2
Flowers	28	22	13	7	6	5	2
Fruits	38	23	12	7	6	4	2
Leaves	84	44	23	13	7	5	2
Tank water	111	61	27	16	8	5	2



3.3. Yeasts from bromeliad compartments

The diversity and composition of yeast species varied among bromeliad compartments. For example, regarding diversity (estimated with Simpson's index), total richness (estimated with Chao 1) and dominance, leaves and water tanks were significantly more diverse, richer and with communities less dominant than flowers and fruits ($p<0.05$). On the other hand, these same variables did not vary significantly between flowers and fruits and between leaves and water tanks ($p>0.05$), as shown in Figure 3A-C. However, the number of singletons (species occurring in only one unity) found in each compartment was proportionally similar to water tanks (57%), leaves (52%), fruits and flowers (both 50%).




Fig. 3- Boxplots showing Simpson's alpha diversity index (A), total richness estimated by Chao-1 (B) and the dominance of yeast communities (C) among the bromeliad phyllosphere compartments.

Furthermore, more than 70% ($n = 129$) of the species were recorded exclusively in a single compartment (Figure 4B). Only four species were shared among all compartments: *A. pullulans*, *Debaryomyces hansenii*, *M. guilliermondii* and *Rhodotorula glutinis*. In terms of the community composition, the most distinct

compartments were fruits and leaves, which shared only eight species and had a Whittaker beta-diversity of 0.85 (Figure 4A). Observing the compartments separately, it is possible to observe that the most exclusive species, in terms of the richness of each compartment, was water tank (62%), followed by leaves (51%), fruit (32%) and flower (18%). The collector curve corroborates the estimates of Chao 1 and indicates that the diversity is even greater than that found, mainly in water tanks and leaves (Figure 4C and Supplementary Figure 2).




Fig. 4- (A), Whittaker's beta-diversity between the different compartments. (B), the Venn diagram illustrates the sharing of species among the different compartments of the bromeliad phyllosphere. (C), Extrapolation rarefaction curve. The curve was elaborated with 1000 pseudoreplica bootstrap and the extrapolation was performed with the Chao1 estimator.

Moreover, depending on the plant compartment analyzed, the species most frequently found may vary. For example, the most frequent yeasts on flowers were *C. intermedia* (38%) and *P. flavesiensis* (33%), on fruits were *M. guilliermondii* (56%), *Hanseniaspora uvarum* (50%) and *Pichia kudriavzevii* (44%), while on leaves they were *P. laurentii* (14%), *Ps. hubeiensis* (11%), *Ca. hohenbergiae* (11%) and *P. leoncinii* (10%) and on water tanks they were *C. intermedia* (36%), *Rhodotorula mucilaginosa* (32%), *P. laurentii* (28%) and *N. albida* (28%).

3.3.1. Yeasts in bromeliad flowers

The only exclusively non-singleton species found in flowers was *Pseudohyphozyma bogoriensis*. Moreover, only three genera have more than one yeast species: *Papiliotrema* ($n = 4$ species; 14%), *Sporobolomyces* ($n = 3$; 11%) and *Aureobasidium* ($n = 2$; 7%), and approximately 86% of the reported genera were unispecific. The Sporidiobolaceae family was the most represented ($n = 5$ species, 18%). As for orders, classes and other hierarchical groups, no major differences were found in the entire phyllosphere. However, in flowers, the subphylum Pucciniomycotina stands out ($n = 8$; 29%), mainly when compared with the class of Microbotryomycetes ($n = 7$; 25%) containing the genus *Sporobolomyces*, *Colacogloea*, *Pseudohyphozyma*, *Rhodosporidiobolus*, and *Rhodotorula*. The phylum Basidiomycota recorded most of the species, accounting for 19 species (69%).

3.3.2. Yeasts in bromeliad fruits

Candida sorboxylosa, *Pi. kudriavzevii*, *Pichia fermentans* and *Saccharomyces cerevisiae* were exclusively isolated from fruits and presented a relatively high frequency in this compartment. Furthermore, *Candida* ($n = 4$ species; 11%), *Rhodotorula* ($n = 4$; 11%), *Pichia* ($n = 4$; 11%), *Hanseniaspora* ($n = 3$; 8%), *Starmerella* ($n = 2$; 6%) and *Meyerozyma* ($n = 2$; 6%) were the genera with more than one reported species. Different than the other compartments, many species of the Saccharomycetaceae family ($n = 14$; almost 40% of all species in this compartment), as well as the order Saccharomycetales ($n = 26$ species; 72%) were present in fruits. This is consequently reflected in the prevalence of species in the subphylum Saccharomycotina and the

Ascomycota phylum, respectively 72 and 78%. Furthermore, among all compartments analyzed in this study, fruits were the only ones without any representative of the Ustilaginomycotina subphylum.

3.3.3. Yeasts in bromeliad leaves

Several species were frequently and exclusively found in leaves: *Farysia itapuensis*, *Genolevuria bromelialarum*, *Papiliotrema siamense*, *Queiroziella brasiliensis*, *Sporobolomyces salmonicolor* and *Symmetrospora suhii*. Moreover, the five most representative genera were *Papiliotrema* ($n = 7$; 9%), *Hannaella* ($n = 6$; 7.5%), *Candida* ($n = 5$; 6%), *Rhodotorula* ($n = 4$; 5%) and *Aureobasidium* ($n = 4$; 5%). Together, these five genera represented a third of the total species in leaves. The families with highest number of species were Sporidiobolaceae ($n = 8$; 10%) and Trimorphomycetaceae ($n = 5$; 6%). However, in terms of order, Tremellales had 31 species (almost 40% of the total), followed by Saccharomycetales ($n = 12$; 15%) and Sporidiobolales ($n = 8$; 10%). The three richest classes account for 70% of yeasts in leaves: Tremellomycetes ($n = 34$; 43%), Saccharomycetes ($n = 12$; 15%) and Microbotryomycetes ($n = 10$; 12%). The subphyla Agaricomycotina and Pucciniomycotina grouped two thirds of the yeast species. Among all compartments, leaves had the highest prevalence of the phylum Basidiomycota, representing 79% of the species.

3.3.4. Yeasts in bromeliad water tank

Bromeliad water tank was the compartment with the greatest richness. At least five species that were not singletons or doubletons were found exclusively in water tanks: *Kazachstania bromeliacearum*, *Kwoniella heveanensis*, *Occultifur brasiliensis*, *Papiliotrema rajasthanensis*, and *Starmerella magnoliae*. The genus *Kwoniella* was found exclusively in water tanks. In addition, *Candida* and *Rhodotorula* were the richest genera ($n = 21$ species; 12% and $n = 8$ species; 7%). Twenty-seven families are registered in this compartment, being Saccharomycetaceae ($n = 18$; 16%), Ustilaginaceae ($n = 11$; 10%) and Sporidiobolaceae ($n = 10$; 9%). Combined, these families compose more than a third of the total yeast species. Almost two-thirds of the species belong to the orders Saccharomycetales ($n = 49$; 43%) and Tremellales ($n = 24$;

21%). This is also true for the classes Saccharomycetes ($n = 49$; 43%) and Tremellomycetes ($n = 29$; 25%) and for the subphyla Saccharomycotina ($n = 49$; 43%) and Agaricomycotina ($n = 29$; 25%). Basidiomycota was the richest phylum, comprising 54% of the species.

3.4. Yeast community structure in bromeliads

3.4.1. Definition of units

From the 20 articles in the category Diversity and ecology, 103 DUAs were defined considering bromeliad species/compartment/paper/sampled site (Figure 1). In this data group, 171 yeast species were recorded. Among the DUAs, 20 corresponded to flowers (19%), 16 to fruits (16%), 54 to leaves (52%), and 13 to water tanks (13%). In 28 units, classical methods (phenotypic) for the identification of yeasts and other molecular methods were used. Our analyses (PERMANOVA) indicated that the methods (phenotypic or molecular) are data classes that in fact exert some explanatory power on the data set, and that the groups formed have little chance of being found by chance ($p=0.001$). However, the low explanatory power shows a low correlation between these classes and the community structure ($R^2=0.06$). The PERMDISP result was also significant ($p=0.004$), indicating a heterogeneous dispersion between groups formed by samples with different identification methods. The group formed by samples identified with molecular methods had higher variance, probably an effect of cryptic species that are a methodological artifact of phenotypic methods. Therefore, the analyses were prepared with all units, without discriminating the types of identification methods, but also with the data of each identification method separately (Table 4).

3.4.2. Community structure

The results found herein were not conclusive, as the statistical significance of the classes varied depending on the data set evaluated (Table 4). Furthermore, even when the result showed statistical significance, the low explanatory values of R^2 indicate a low correlation between the variables and community structure. Bromeliad compartments and functional types were the categories with the highest average explanatory power (although the values were low). One of the reasons for the low

explanatory power can be the large variance in some classes, for example in leaves and in CAM tank-epiphytes. Comparing the classes of each category pair-by-pair (without discriminating the types of identification method), the three subfamilies registered diverged significantly from each other (p -value was always ≤ 0.003). However, regarding the functional types, only CAM tank-epiphyte bromeliads diverged significantly from C₃ tank-epiphytes ($p=0.023$) and from CAM atmospheric epiphyte ($p=0.002$). In addition, CAM atmospheric epiphyte bromeliads diverged marginally from CAM terrestrial plants ($p=0.056$). Nonetheless, in compartments, the leaves were the only class that diverged significantly from flowers ($p=0.01$) and fruits ($p=0.003$). When comparing the distance of the centroids of each type of compartment (flowers, fruits, leaves and water tanks) to evaluate internal heterogeneity, leaves were the compartment with the highest variance (Figure 5). The difference in the variance of the groups was significant ($p<0.0001$) and when compared pair-by-pair, leaves diverged significantly from fruits ($p=0.0001$), water tanks ($p=0.0009$) and showed a marginally significant difference in relation to flowers ($p=0.064$).




Fig. 5- (A) Boxplot of the distances of the points on the PCoA in relation to the centroid (variance) of the compartments and bromeliads. (B) Principal Coordinate Analysis (PCoA) applying Jaccard distance from the yeast community on bromeliads, the points refer to the Delimited Units of Analysis (DUAs) used in the analysis, and the central point of each class is the centroid.

3.5. New yeast species in bromeliads

A total of 22 new yeast species described were recorded (7 ascomycetes and 15 basidiomycetes) associated with 30 species of bromeliads (Supplementary Table 1). The genera with the highest number of new species described are *Candida* (n = 4) and *Carlosrosaea* (n = 3). In addition, Tremellales and Saccharomycetales are the most recorded orders, presenting 10 and 7 new species, respectively. The new species described are grouped into 4 subphyla, Agaricomycotina (n = 10 species), Saccharomycotina (n = 7), Pucciniomycotina (n = 3) and Ustilaginomycotina (n = 2).

The new species were described associated with leaves (n = 10), water tanks (n = 8), with some new species found in both substrates (n = 3). Only the species *Ca. hohenbergiae* was described as associated with leaves and flowers. The bromeliad species with the highest number of new yeast species described were *V. minarum* L.B.Sm. (n = 6), *Vriesea friburgensis* Mez (n = 4) and *Werauhia gigantea* (Mart. ex Schult. & Schult.f.) J.R.Grant (n = 4); both belong to the subfamily Tillandsioideae. *P. leoncini* was the species associated with the highest number of bromeliad species (10 species).

3.6. Bromeliad yeasts in biotechnological processes

In the analyses, at least 50 species of yeasts were found isolated from bromeliads capable of being used in biotechnological processes in the industrial, agricultural, and/or environmental sectors (Supplementary Table 2). The industrial potential of yeasts isolated from bromeliads is mainly the production of extracellular enzymes such as amylase, cellulase, esterase, pectinase, protease, and xylanase. More than 90% of the listed species with some biotechnological potentials were able to produce at least one of these enzymes. The enzymes produced by the largest number of yeast species were respectively protease, pectinase and xylanase. However, other biotechnological applications were also registered, such as the probiotic activity of *Meyerozyma caribbica*, the potential in the production of biodiesel from *Candida* sp., and the production of fermented beverages by *Hanseniaspora guilliermondii*, *Hanseniaspora*

opuntiae, *H. uvarum*, *M. caribbica*, *M. guilliermondii*, *Saccharomyces cerevisiae-like* and *Saccharomycodes ludwigii*. Among agricultural and related applications are the promotion of plant growth by *Carlosrosaea vriesae* and the production of traits related to plant growth, such as the production of indoleacetic acid (IAA), siderophores and the ability to solubilize phosphate. In addition, yeasts from bromeliads have also shown their value as a potential source of animal nutrition, such as *M. caribbica* and *S. ludwigii*, which were used as feed in fish farming. In the environmental sector, *A. pullulans*, *Candida* spp., *Cryptococcus* spp. and *Sporobolomyces* spp. were used as potential air quality indicators. In addition, *Candida* sp. showed potential in bioremediation, specifically related to the treatment of oil-contaminated environments.

Table 3- Yeasts taxa, hosts and compartments recorded in the review.

Yeast species	Host species	Flower	Fruit	Leaves	Water tank	Reference
Ascomycota (Phyla)						
Pezizomycotina (Subphyla)						
<i>Aureobasidium leucospermi</i>	<i>Ananas comosus</i> (L.) Merr.		x			Chanprasartsuk et al., 2012
	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira		x			Navarro et al., 2020
<i>Aureobasidium melanogenum</i>	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira		x			Navarro et al., 2020
	<i>Tillandsia caput-medusae</i> É.Morren		x			Brighigna et al 2000
	<i>Tillandsia schiedeana</i> Steud.		x			Brighigna et al 2000
	<i>Ananas comosus</i> (L.) Merr.		x			Chanprasartsuk et al., 2010; 2013
<i>Aureobasidium pullulans</i>	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira	x				Félix et al., 2021
	<i>Aechmea tomentosa</i> Mez	x				Félix et al., 2021
	<i>Vriesea minarum</i> L.B.Sm.			x		Gomes et al., 2015
	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.			x		Hagler et al., 1993
	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.			x		Hagler et al., 1993
	<i>Bromelia karatas</i> L.			x		Moraes et al., 2020
	<i>Billbergia</i> sp.		x			Navarro et al., 2020
	<i>Canistrum improcerum</i> Leme & J.A.Siqueira		x			Navarro et al., 2020
<i>Aureobasidium thailandense</i>	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira	x				Félix et al., 2021
	<i>Aechmea tomentosa</i> Mez	x				Félix et al., 2021
	<i>Billbergia</i> sp.	x				Félix et al., 2021
	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira		x			Navarro et al., 2020
	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira		x			Navarro et al., 2020

	<i>Aechmea tomentosa</i> Mez		x	Navarro et al., 2020
	<i>Aechmea werdermannii</i> Harms		x	Navarro et al., 2020
	<i>Ananas comosus</i> (L.) Merr.		x	Navarro et al., 2020
	<i>Billbergia</i> sp.		x	Navarro et al., 2020
	<i>Hohenbergia stellata</i> Schult. & Schult.f.		x	Navarro et al., 2020
<i>Endosporium aviarium</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Morais et al., 2020
<i>Exophiala placitae</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Morais et al., 2020
<i>Hortaea werneckii</i>	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira	x		Formoso et al., 2015; Navarro et al., 2020
<i>Talaromyces amestolkiae</i>	<i>Ananas comosus</i> (L.) Merr.	x		Leneveu-jenvrin et al., 2020
Saccharomycotina (Subphyla)				
<i>Candida aechmeae</i>	<i>Aechmea recurvata</i> (Klotzsch) L.B.Sm.		x	Landell et al., 2010
	<i>Billbergia nutans</i> H.Wendl.		x	Landell et al., 2010
<i>Candida albicans</i>	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Ananas comosus</i> (L.) Merr.	x		Udota and Urura, 2010
<i>Candida bromeliacearum</i>	<i>Canistropsis seidelii</i> (L.B.Sm. & Reitz) Leme		x	Ruivo et al., 2005
<i>Candida buenavistaensis</i>	<i>Encholirium</i> sp.		x	Morais et al., 2020
<i>Candida diddensiae</i>	<i>Werauhia gigantea</i> (Mart. ex Schult. & Schult.f.) J.R.Grant		x	Landell et al., 2006
<i>Candida duobushaemulonis</i>	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
<i>Candida glabrata</i>	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Encholirium</i> sp.		x	Morais et al., 2020
<i>Candida glucosophila</i>	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998
<i>Candida heveicola</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Morais et al., 2020
<i>Candida intermedia</i>	<i>Aechmea nudicaulis</i> (L.) Griseb.		x	Araujo et al., 1998
	<i>Nidularium procerum</i> Lindm.		x	Araujo et al., 1998
	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998
	<i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm.		x	Araujo et al., 1998

	<i>Aechmea costantinii</i> (Mez) L.B.Sm.	x		Félix et al., 2021
	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira	x		Félix et al., 2021
	<i>Aechmea fulgens</i> Brongn.	x		Félix et al., 2021
	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira	x		Félix et al., 2021
	<i>Billbergia</i> sp.	x		Félix et al., 2021
	<i>Canistrum alagoanum</i> Leme & J.A.Siqueira	x		Félix et al., 2021
	<i>Canistrum aurantiacum</i> É.Morren	x		Félix et al., 2021
	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Encholirium</i> sp.		x	Morais et al., 2020
	<i>Araeococcus chlorocarpus</i> (Wawra) Leme & J.A.Siqueira	x		Navarro et al., 2020
<i>Candida jaroonii</i>	<i>Canistrum aurantiacum</i> É.Morren	x		Navarro et al., 2020
<i>Candida leandreae</i>	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
<i>Candida melibiosica</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
<i>Candida membranifaciens</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
	<i>Ananas comosus</i> (L.) Merr.	x		Nasir et al., 2017
	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2013
<i>Candida nivariensis</i>	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Encholirium</i> sp.		x	Morais et al., 2020
<i>Candida orthopsilosis</i>	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Encholirium</i> sp.		x	Morais et al., 2020
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Encholirium</i> sp.		x	Morais et al., 2020
<i>Candida parapsilosis</i>	<i>Araeococcus chlorocarpus</i> (Wawra) Leme & J.A.Siqueira	x		Navarro et al., 2020
	<i>Canistrum alagoanum</i> Leme & J.A.Siqueira	x	x	Navarro et al., 2020

	<i>Hohenbergia ridleyi</i> (Baker) Mez		x	Navarro et al., 2020
<i>Candida pseudointermedia</i>	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Encholirium</i> sp.		x	Morais et al., 2020
<i>Candida sinolaborantium</i>	<i>Aechmea fulgens</i> Brongn.		x	Navarro et al., 2020
<i>Candida soli</i>	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
<i>Candida sorboxylosa</i>	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2010
	<i>Aechmea nudicaulis</i> (L.) Griseb.		x	Araujo et al., 1998
	<i>Nidularium procerum</i> Lindm.		x	Araujo et al., 1998
<i>Candida tropicalis</i>	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998
	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2010; 2013; Udoa and Urúa, 2010
<i>Candida ubatubensis</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015; Morais et al., 2020
	<i>Canistropsis seidelii</i> (L.B.Sm. & Reitz) Leme		x	Ruivo et al., 2005
<i>Candida vrieseae</i>	<i>Werauhia gigantea</i> (Mart. ex Schult. & Schult.f.) J.R.Grant		x	Landell et al., 2010
<i>Clavispora lusitaniae</i>	<i>Ananas comosus</i> (L.) Merr.	x		Amorim et al 2018
<i>Cyberlindnera saturnus</i>	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998
	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.		x	Araujo et al., 1998
	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998
	<i>Aechmea recurvata</i> (Klotzsch) L.B.Sm.	x		Landell et al., 2006
<i>Debaryomyces hansenii</i>	<i>Billbergia nutans</i> H.Wendl.		x	Landell et al., 2006
	<i>Tillandsia gardneri</i> Lindl.	x		Landell et al., 2006
	<i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm.		x	Landell et al., 2006
	<i>Ananas comosus</i> (L.) Merr.	x		Udoa and Urúa, 2010
<i>Dipodascus albidus</i>	<i>Aechmea recurvata</i> (Klotzsch) L.B.Sm.	x		Landell et al., 2006
<i>Diutina rugosa</i>	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.		x	Araujo et al., 1998
	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998

<i>Galactomyces candidus</i>	<i>Ananas comosus</i> (L.) Merr.	x		Udota and Urua, 2010
<i>Hagleromyces aurorensis</i>	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Bromelia karatas</i> L.		x	Sousa et al., 2014
<i>Hanseniaspora guilliermondii</i>	<i>Ananas comosus</i> (L.) Merr.	x		Dellacassa et al., 2017
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
<i>Hanseniaspora opuntiae</i>	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2010; 2013; Dellacassa et al., 2017
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
<i>Hanseniaspora uvarum</i>	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2010; 2013; Dellacassa et al., 2017
	<i>Aechmea fulgens</i> Brongn.	x		Félix et al., 2021
	<i>Cryptanthus dianae</i> Leme	x		Félix et al., 2021
	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
	<i>Ananas comosus</i> (L.) Merr.	x		Udota and Urua, 2010
<i>Kazachstania africana</i>	<i>Aechmea nudicaulis</i> (L.) Griseb.		x	Araujo et al., 1998
<i>Kazachstania bromeliacearum</i>	<i>Aechmea nudicaulis</i> (L.) Griseb.		x	Araujo et al., 2012
	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.		x	Araujo et al., 2012
	<i>Nidularium procerum</i> Lindm.		x	Araujo et al., 2012
	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Araujo et al., 2012
	<i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm.		x	Araujo et al., 2012
<i>Kazachstania rupicola</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Safar et al., 2013; Gomes et al., 2015
<i>Kloeckera apiculata</i> var. <i>apis</i>	<i>Ananas comosus</i> (L.) Merr.	x		Korres et al., 2011
<i>Kluyveromyces aestuarii</i>	<i>Aechmea nudicaulis</i> (L.) Griseb.		x	Araujo et al., 1998
<i>Kluyveromyces marxianus</i>	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
	<i>Ananas comosus</i> (L.) Merr.	x		Udota and Urua, 2010
<i>Kodamaea ohmeri</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
<i>Lachancea thermotolerans</i>	<i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm.		x	Araujo et al., 1998

<i>Metschnikowia fructicola</i>	<i>Tillandsia geminiflora</i> Brongn.		x	Landell et al., 2006
<i>Metschnikowia hawaiiensis</i>	<i>Tillandsia gardneri</i> Lindl.		x	Landell et al., 2006
<i>Metschnikowia koreensis</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
<i>Meyerozyma caribbica</i>	<i>Ananas comosus</i> (L.) Merr.	x		Amorim et al 2018; Leneveu-jenvrin et al., 2020; Islam et al., 2021
	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Encholirium</i> sp.		x	Morais et al., 2020
<i>Meyerozyma guilliermondii</i>	<i>Ananas comosus</i> (L.) Merr.	x		Reyes et al., 2004; Chanprasartsuk et al., 2010; 2013; Dellacassa et al., 2017; Di Cagno et al., 2010
	<i>Aechmea muricata</i> (Arruda) L.B.Sm.	x		Félix et al., 2021
	<i>Aechmea werdermannii</i> Harms	x		Félix et al., 2021
	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Encholirium</i> sp.		x	Morais et al., 2020
	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira	x		Navarro et al., 2020
	<i>Aechmea fulgens</i> Brongn.		x	Navarro et al., 2020
	<i>Billbergia</i> sp.		x	Navarro et al., 2020
<i>Pichia fermentans</i>	<i>Canistrum alagoanum</i> Leme & J.A.Siqueira	x		Navarro et al., 2020
	<i>Ananas comosus</i> (L.) Merr.	x		Udota and Urua, 2010; Chanprasartsuk et al. 2013; Islam et al., 2021
<i>Pichia kudriavzevii</i>	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2010; 2013; Korres et al., 2011
<i>Pichia membranifaciens</i>	<i>Aechmea nudicaulis</i> (L.) Griseb.		x	Araujo et al., 1998
	<i>Nidularium procerum</i> Lindm.		x	Araujo et al., 1998
	<i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm.		x	Araujo et al., 1998

	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2013
	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
<i>Pichia occidentalis</i>	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2010
<i>Priceomyces melissophilus</i>	<i>Tillandsia gardneri</i> Lindl.		x	Landell et al., 2006
<i>Saccharomyces cerevisiae</i>	<i>Ananas comosus</i> (L.) Merr.	x		Udota and Urúa, 2010; Leneveu-jenvrin et al., 2020
<i>Saccharomyces ludwigii</i>	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2010; 2013
<i>Saturnispora silvae</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
<i>Schwanniomyces etchellsii</i>	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998
	<i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm.		x	Araujo et al., 1998
<i>Schwanniomyces occidentalis</i>	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.		x	Araujo et al., 1998
<i>Schwanniomyces vanrijiae</i>	<i>Nidularium procerum</i> Lindm.		x	Araujo et al., 1998
	<i>Tillandsia gardneri</i> Lindl.		x	Landell et al., 2006
<i>Starmerella apicola</i>	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2010
<i>Starmerella magnoliae</i>	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998
	<i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm.		x	Araujo et al., 1998
<i>Starmerella stellata</i>	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2013
<i>Torulaspora delbrueckii</i>	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira	x		Félix et al., 2021
	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.		x	Hagler et al., 1993
	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Ananas comosus</i> (L.) Merr.	x		Udota and Urúa, 2010
<i>Wickerhamiella jalapaonensis</i>	<i>Aechmea fulgens</i> Brongn.	x		Félix et al., 2021
<i>Wickerhamiella sorbophila</i>	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
<i>Wickerhamomyces anomalus</i>	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998
	<i>Ananas comosus</i> (L.) Merr.	x		Dellacassa et al., 2017
<i>Yamadazyma mexicana</i>	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998

<i>Yarrowia lipolytica</i>	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998
	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2010; 2013
<i>Zygocharamyces bailii</i>	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2010; 2013
	<i>Tillandsia gardneri</i> Lindl.	x		Landell et al., 2006
Basidiomycota (Phyla)				
Agaricomycotina (Subphyla)				
<i>Atelosaccharomyces hodeloi</i>	<i>Nidularium procerum</i> Lindm.		x	Araujo et al., 1998
	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
<i>Bulleribasidium variabile</i>	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998
<i>Bulleromyces albus</i>	<i>Bromelia antiacantha</i> Bertol.		x	Landell et al., 2006
	<i>Tillandsia geminiflora</i> Brongn.		x	Landell et al., 2006
<i>Carcinomyces nordestinensis</i>	<i>Bromelia antiacantha</i> Bertol.		x	Crous et al., 2019
<i>Carlosrosaea aechmeae</i>	<i>Aechmea costantinii</i> (Mez) L.B.Sm.		x	Félix et al., 2017; Navarro et al., 2020
	<i>Aechmea fulgens</i> Brongn.	x	x	Félix et al., 2017, 2021; Navarro et al., 2020
	<i>Canistrum alagoanum</i> Leme & J.A.Siqueira	x		Félix et al., 2017, 2021
	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira		x	Félix et al., 2017; Navarro et al., 2020
	<i>Hohenbergia ramageana</i> Mez		x	Félix et al., 2017
	<i>Tillandsia</i> sp.		x	Félix et al., 2017; Navarro et al., 2020
	<i>Canistrum aurantiacum</i> É.Morren		x	Navarro et al., 2020
	<i>Hohenbergia ridleyi</i> (Baker) Mez		x	Navarro et al., 2020
	<i>Hohenbergia stellata</i> Schult. & Schult.f.		x	Navarro et al., 2020
	<i>Tillandsia chapeuensis</i> Rauh		x	Navarro et al., 2020
<i>Carlosrosaea hohenbergiae</i>	<i>Tillandsia gardneri</i> Lindl.		x	Landell et al., 2015
	<i>Vriesea friburgensis</i> Mez		x	Landell et al., 2015
	<i>Vriesea minarum</i> L.B.Sm.		x	Landell et al., 2015

<i>Cystofilobasidium macerans</i>	<i>Neoregelia cruenta</i> (Graham) L.B.Sm. <i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993 Hagler et al., 1993
<i>Dimennazyma cisti-albidi</i>	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira	x		Félix et al., 2021
	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.		x	Hagler et al., 1993
	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
<i>Dioszegia hungarica</i>	<i>Vriesea friburgensis</i> Mez <i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm. <i>Werauhia gigantea</i> (Mart. ex Schult. & Schult.f.) J.R.Grant		x x x	Landell et al., 2006 Landell et al., 2006 Landell et al., 2006
<i>Fellomyces borneensis</i>	<i>Aechmea werdermannii</i> Harms <i>Neoregelia gigas</i> Leme & L.Kollmann		x x	Navarro et al., 2020 Navarro et al., 2020
<i>Fellomyces fuzhouensis</i>	<i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm.		x	Landell et al., 2006
<i>Fellomyces penicillatus</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
<i>Fellomyces polyborus</i>	<i>Dyckia</i> sp.		x	Landell et al., 2006
	<i>Tillandsia gardneri</i> Lindl.		x	Landell et al., 2009
<i>Genolevuria bromelialarum</i>	<i>Vriesea friburgensis</i> Mez <i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm.		x x	Landell et al., 2009 Landell et al., 2009
<i>Goffeauzyma gastrica</i>	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
<i>Hannaella kunmingensis</i>	<i>Hohenbergia stellata</i> Schult. & Schult.f.		x	Navarro et al., 2020
	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
	<i>Tillandsia gardneri</i> Lindl.	x		Landell et al., 2006
	<i>Tillandsia geminiflora</i> Brongn.		x	Landell et al., 2006
<i>Hannaella luteola</i>	<i>Bromelia karatas</i> L. <i>Encholirium</i> sp.		x x	Morais et al., 2020 Morais et al., 2020
	<i>Aechmea fulgens</i> Brongn.		x	Navarro et al., 2020
	<i>Neoregelia gigas</i> Leme & L.Kollmann		x	Navarro et al., 2020
<i>Hannaella oryzae</i>	<i>Araeococcus chlorocarpus</i> (Wawra) Leme & J.A.Siqueira		x	Navarro et al., 2020

	<i>Vriesea minarum</i> L.B.Sm.		✗	✗	Landell et al., 2014; Gomes et al., 2015
	<i>Bromelia karatas</i> L.		✗		Landell et al., 2014; Morais et al., 2020
	<i>Encholirium</i> sp.		✗		Landell et al., 2014; Morais et al., 2020
	<i>Tillandsia geminiflora</i> Brongn.		✗		Landell et al., 2014
<i>Hannaella pagnoccae</i>	<i>Werauhia gigantea</i> (Mart. ex Schult. & Schult.f.) J.R.Grant		✗		Landell et al., 2014
	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira		✗		Navarro et al., 2020
	<i>Aechmea muricata</i> (Arruda) L.B.Sm.		✗		Navarro et al., 2020
	<i>Araeococcus chlorocarpus</i> (Wawra) Leme & J.A.Siqueira		✗		Navarro et al., 2020
	<i>Hohenbergia stellata</i> Schult. & Schult.f.		✗		Navarro et al., 2020
<i>Hannaella phetchabunensis</i>	<i>Tillandsia</i> sp.		✗		Navarro et al., 2020
<i>Hannaella siamensis</i>	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira		✗		Navarro et al., 2020
<i>Hannaella sinensis</i>	<i>Vriesea minarum</i> L.B.Sm.			✗	Gomes et al., 2015
<i>Hannaella zaeae</i>	<i>Bromelia karatas</i> L.			✗	Morais et al., 2020
	<i>Encholirium</i> sp.			✗	Morais et al., 2020
<i>Kockovaella libkindii</i>	<i>Vriesea minarum</i> L.B.Sm.			✗	Gomes et al., 2016
<i>Kockovaella sacchari</i>	<i>Neoregelia gigas</i> Leme & L.Kollmann		✗		Navarro et al., 2020
<i>Kwoniella dendrophila</i>	<i>Bromelia karatas</i> L.			✗	Morais et al., 2020
	<i>Bromelia karatas</i> L.			✗	Morais et al., 2020
<i>Kwoniella heveanensis</i>	<i>Encholirium</i> sp.			✗	Morais et al., 2020
	<i>Vriesea minarum</i> L.B.Sm.			✗	Morais et al., 2020
	<i>Bromelia karatas</i> L.			✗	Morais et al., 2020
<i>Kwoniella mangroviensis</i>	<i>Encholirium</i> sp.			✗	Morais et al., 2020
<i>Naganishia albida</i>	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.			✗	Araujo et al., 1998
	<i>Nidularium procerum</i> Lindm.			✗	Araujo et al., 1998
	<i>Quesnelia arvensis</i> (Vell.) Mez			✗	Araujo et al., 1998

	<i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm.		x	Araujo et al., 1998
	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.		x	Hagler et al., 1993
	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
	<i>Aechmea recurvata</i> (Klotzsch) L.B.Sm.	x		Landell et al., 2006
	<i>Bromelia antiacantha</i> Bertol.	x		Landell et al., 2006
	<i>Tillandsia gardneri</i> Lindl.	x		Landell et al., 2006
	<i>Tillandsia geminiflora</i> Brongn.	x		Landell et al., 2006
	<i>Vriesea friburgensis</i> Mez	x		Landell et al., 2006
	<i>Werauhia gigantea</i> (Mart. ex Schult. & Schult.f.) J.R.Grant	x		Landell et al., 2006
	<i>Ananas comosus</i> (L.) Merr.	x		Reyes et al., 2004
<i>Papiliotrema flavescentia</i>	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira	x	x	Navarro et al., 2020; Félix et al., 2021
	<i>Aechmea muricata</i> (Arruda) L.B.Sm.	x		Félix et al., 2021
	<i>Araeococcus chlorocarpus</i> (Wawra) Leme & J.A.Siqueira	x	x	Navarro et al., 2020; Félix et al., 2021
	<i>Billbergia</i> sp.	x		Félix et al., 2021
	<i>Canistrum aurantiacum</i> É.Morren	x		Félix et al., 2021
	<i>Cryptanthus dianae</i> Leme	x		Félix et al., 2021
	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015; Morais et al., 2020
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Encholirium</i> sp.		x	Morais et al., 2020
	<i>Aechmea costantinii</i> (Mez) L.B.Sm.	x		Navarro et al., 2020
	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira	x		Navarro et al., 2020
	<i>Aechmea werdermannii</i> Harms	x		Navarro et al., 2020
	<i>Canistrum improcerum</i> Leme & J.A.Siqueira	x		Navarro et al., 2020
<i>Papiliotrema laurentii</i>	<i>Hohenbergia stellata</i> Schult. & Schult.f.	x		Navarro et al., 2020
	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.		x	Hagler et al., 1993; Araujo et al., 1998

<i>Papiliotrema leoncinii</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015; Morais et al., 2020
	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Gomes et al., 2015
	<i>Bromelia antiacantha</i> Bertol.	x		Landell et al., 2006
	<i>Tillandsia crocata</i> (É.Morren) N.E.Br.	x	x	Landell et al., 2006
	<i>Tillandsia gardneri</i> Lindl.		x	Landell et al., 2006
	<i>Tillandsia geminiflora</i> Brongn.		x	Landell et al., 2006
	<i>Weruahia gigantea</i> (Mart. ex Schult. & Schult.f.) J.R.Grant		x	Landell et al., 2006
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Encholirium</i> sp.		x	Morais et al., 2020
	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira		x	Navarro et al., 2020
	<i>Aechmea fulgens</i> Brongn.		x	Navarro et al., 2020
	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira		x	Navarro et al., 2020
	<i>Ananas comosus</i> (L.) Merr.		x	Navarro et al., 2020
	<i>Canistrum alagoanum</i> Leme & J.A.Siqueira		x	Navarro et al., 2020
	<i>Canistrum aurantiacum</i> É.Morren		x	Navarro et al., 2020
	<i>Tillandsia chapeuensis</i> Rauh		x	Navarro et al., 2020
	<i>Aechmea fulgens</i> Brongn.		x	Pagani et al., 2016
	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira		x	Pagani et al., 2016
	<i>Aechmea recurvata</i> (Klotzsch) L.B.Sm.		x	Pagani et al., 2016
	<i>Bromelia antiacantha</i> Bertol.		x	Pagani et al., 2016
	<i>Bromelia</i> sp.		x	Pagani et al., 2016
	<i>Canistrum alagoanum</i> Leme & J.A.Siqueira		x	Pagani et al., 2016
	<i>Canistrum aurantiacum</i> É.Morren		x	Pagani et al., 2016
	<i>Tillandsia crocata</i> (É.Morren) N.E.Br.		x	Pagani et al., 2016
	<i>Tillandsia gardneri</i> Lindl.		x	Pagani et al., 2016
	<i>Vriesea friburgensis</i> Mez		x	Pagani et al., 2016

	<i>Werauhia gigantea</i> (Mart. ex Schult. & Schult.f.) J.R.Grant		x	Pagani et al., 2016
<i>Papiliotrema mangalensis</i>	<i>Aechmea fulgens</i> Brongn.		x	Navarro et al., 2020
	<i>Cryptanthus dianae</i> Leme		x	Navarro et al., 2020
<i>Papiliotrema miconiae</i>	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira	x		Félix et al., 2021
	<i>Canistrum alagoanum</i> Leme & J.A.Siqueira	x		Félix et al., 2021
	<i>Canistrum aurantiacum</i> É.Morren	x		Félix et al., 2021
	<i>Aechmea werdermannii</i> Harms		x	Navarro et al., 2020
<i>Papiliotrema nemorosa</i>	<i>Aechmea costantinii</i> (Mez) L.B.Sm.	x		Félix et al., 2021
	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira	x		Félix et al., 2021
	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015; Morais et al., 2020
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Encholirium</i> sp.		x	Morais et al., 2020
	<i>Cryptanthus zonatus</i> (Vis.) Beer	x		Navarro et al., 2020
	<i>Tillandsia chapeuensis</i> Rauh		x	Navarro et al., 2020
<i>Papiliotrema rajasthanensis</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Encholirium</i> sp.		x	Morais et al., 2020
<i>Papiliotrema siamense</i>	<i>Ananas comosus</i> (L.) Merr.		x	Navarro et al., 2020
	<i>Cryptanthus dianae</i> Leme		x	Navarro et al., 2020
	<i>Tillandsia kegeliana</i> Mez		x	Navarro et al., 2020
<i>Papiliotrema terrestris</i>	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Encholirium</i> sp.		x	Morais et al., 2020
<i>Phaeotremella foliacea</i>	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998
<i>Piskurozyma taiwanensis</i>	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira	x		Navarro et al., 2020
<i>Rhynchogastrema complexa</i>	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Ananas comosus</i> (L.) Merr.		x	Valente et al., 2012

	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.		x	Valente et al., 2012
	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
<i>Saitozyma flava</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Morais et al., 2020
	<i>Bromelia karatas</i> L.		x	Navarro et al., 2020
	<i>Cryptanthus dianae</i> Leme	x		Félix et al., 2021
	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
<i>Saitozyma podzolica</i>	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira		x	Navarro et al., 2020
	<i>Aechmea muricata</i> (Arruda) L.B.Sm.		x	Navarro et al., 2020
	<i>Aechmea werdermannii</i> Harms		x	Navarro et al., 2020
	<i>Ananas comosus</i> (L.) Merr.		x	Navarro et al., 2020
	<i>Araeococcus chlorocarpus</i> (Wawra) Leme & J.A.Siqueira		x	Navarro et al., 2020
<i>Solicoccozyma aeria</i>	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
<i>Tremella fuciformis</i>	<i>Cryptanthus dianae</i> Leme		x	Navarro et al., 2020
	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2010; 2013
<i>Tremella globispora</i>	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Vriesea minarum</i> L.B.Sm.		x	Morais et al., 2020
<i>Trichosporon bergelli</i>	<i>Aechmea nudicaulis</i> (L.) Griseb.		x	Araujo et al., 1998
<i>Vanrija humicola</i>	<i>Vriesea friburgensis</i> Mez		x	Landell et al., 2006
	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira	x		Félix et al., 2021
	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira		x	Félix et al., 2020; Navarro et al., 2020
<i>Vishniacozyma alagoana</i>	<i>Bromelia antiacantha</i> Bertol.		x	Félix et al., 2020
	<i>Hohenbergia stellata</i> Schult. & Schult.f.		x	Félix et al., 2020; Navarro et al., 2020
	<i>Aechmea fulgens</i> Brongn.		x	Navarro et al., 2020
<i>Vishniacozyma taibaiensis</i>	<i>Billbergia</i> sp.		x	Navarro et al., 2020

	<i>Cryptanthus dianae</i> Leme		x	Navarro et al., 2020
<i>Vonarxula javanica</i>	<i>Billbergia</i> sp.		x	Navarro et al., 2020
Pucciniomycotina (Subphyla)				
<i>Colacogloea diffluens</i>	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira	x		Félix et al., 2021
	<i>Tillandsia chapeuensis</i> Rauh		x	Navarro et al., 2020
<i>Curvibasidium nothofagi</i>	<i>Encholirium</i> sp.		x	Morais et al., 2020
<i>Cyrenella elegans</i>	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira		x	Navarro et al., 2020
	<i>Aechmea tomentosa</i> Mez		x	Navarro et al., 2020
<i>Cystobasidium calyptogenae</i>	<i>Encholirium</i> sp.		x	Morais et al., 2020
<i>Cystobasidium laryngis</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998
	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
<i>Cystobasidium minutum</i>	<i>Tillandsia gardneri</i> Lindl.		x	Landell et al., 2006
	<i>Vriesea friburgensis</i> Mez		x	Landell et al., 2006
	<i>Werauhia gigantea</i> (Mart. ex Schult. & Schult.f.) J.R.Grant		x	Landell et al., 2006
	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2010; 2013
<i>Erythrobasidium hasegawianum</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
	<i>Encholirium</i> sp.		x	Morais et al., 2020
<i>Microbotryozyma collariae</i>	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
<i>Occultifur brasiliensis</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
<i>Occultifur externus</i>	<i>Encholirium</i> sp.		x	Morais et al., 2020
	<i>Vriesea minarum</i> L.B.Sm.		x	Morais et al., 2020
	<i>Tillandsia chapeuensis</i> Rauh		x	Navarro et al., 2020
<i>Occultifur plantarum</i>	<i>Bromelia karatas</i> L.	x		Félix et al., 2021

	<i>Canistrum aurantiacum</i> É.Morren	x		Félix et al., 2021
	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.		x	Khunnamwong et al., 2017
	<i>Aechmea werdermannii</i> Harms		x	Navarro et al., 2020
<i>Pseudohyphozyma bogoriensis</i>	<i>Aechmea fulgens</i> Brongn.	x		Félix et al., 2021
	<i>Araeococcus chlorocarpus</i> (Wawra) Leme & J.A.Siqueira	x		Félix et al., 2021
<i>Queiroziella brasiliensis</i>	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira		x	Crous et al., 2018
	<i>Tillandsia geminiflora</i> Brongn.		x	Crous et al., 2018
	<i>Werauhia gigantea</i> (Mart. ex Schult. & Schult.f.) J.R.Grant		x	Crous et al., 2018
	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira		x	Navarro et al., 2020
<i>Rhodosporidiobolus poonsookiae</i>	<i>Cryptanthus dianae</i> Leme		x	Navarro et al., 2020
<i>Rhodosporidiobolus ruineniae</i>	<i>Aechmea muricata</i> (Arruda) L.B.Sm.	x		Félix et al., 2021
	<i>Cryptanthus dianae</i> Leme	x		Félix et al., 2021
	<i>Encholirium</i> sp.		x	Morais et al., 2020
<i>Rhodosporidium diobovatum</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015
<i>Rhodotorula aurantiaca</i>	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998
	<i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm.		x	Araujo et al., 1998
	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.		x	Hagler et al., 1993
	<i>Aechmea recurvata</i> (Klotzsch) L.B.Sm.		x	Landell et al., 2006
	<i>Tillandsia gardneri</i> Lindl.		x	Landell et al., 2006
	<i>Tillandsia geminiflora</i> Brongn.		x	Landell et al., 2006
	<i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm.		x	Landell et al., 2006
	<i>Werauhia gigantea</i> (Mart. ex Schult. & Schult.f.) J.R.Grant		x	Landell et al., 2006
	<i>Ananas comosus</i> (L.) Merr.	x		Reyes et al., 2004
<i>Rhodotorula babjevae</i>	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.		x	Hagler et al., 1993
	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993

	<i>Aechmea nudicaulis</i> (L.) Griseb.		x	Araujo et al., 1998
	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.		x	Hagler et al., 1993; Araujo et al., 1998
	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
<i>Rhodotorula glutinis</i>	<i>Tillandsia crocata</i> (É.Morren) N.E.Br.	x		Landell et al., 2006
	<i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm.		x	Landell et al., 2006
	<i>Ananas comosus</i> (L.) Merr.		x	Reyes et al., 2004
<i>Rhodotorula graminis</i>	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.		x	Hagler et al., 1993
<i>Rhodotorula lactosa</i>	<i>Tillandsia geminiflora</i> Brongn.		x	Landell et al., 2006
	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.		x	Araujo et al., 1998
	<i>Nidularium procerum</i> Lindm.		x	Araujo et al., 1998
	<i>Quesnelia arvensis</i> (Vell.) Mez		x	Araujo et al., 1998
<i>Rhodotorula mucilaginosa</i>	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2010; 2013; Leneveu-jenvrin et al., 2020
	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015; Morais et al., 2020
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Encholirium</i> sp.		x	Morais et al., 2020
<i>Rhodotorula paludigena</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Morais et al., 2020
<i>Rhodotorula taiwanensis</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Morais et al., 2020
	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira	x		Navarro et al., 2020
<i>Rhodotorula toruloides</i>	<i>Ananas comosus</i> (L.) Merr.	x		Chanprasartsuk et al., 2013
	<i>Encholirium</i> sp.		x	Morais et al., 2020
<i>Sporobolomyces carnicolor</i>	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira	x		Félix et al., 2021
	<i>Bromelia antiacantha</i> Bertol.		x	Landell et al., 2006
<i>Sporobolomyces roseus</i>	<i>Tillandsia gardneri</i> Lindl.		x	Landell et al., 2006
	<i>Tillandsia geminiflora</i> Brongn.	x		Landell et al., 2006
	<i>Tillandsia stricta</i> Sol. ex Ker Gawl.		x	Landell et al., 2006

	<i>Werauhia gigantea</i> (Mart. ex Schult. & Schult.f.) J.R.Grant		x	Landell et al., 2006
	<i>Ananas comosus</i> (L.) Merr.	x		Udota and Urua, 2010
<i>Sporobolomyces salmonicolor</i>	<i>Bromelia antiacantha</i> Bertol.		x	Landell et al., 2006
	<i>Tillandsia geminiflora</i> Brongn.		x	Landell et al., 2006
	<i>Vriesea friburgensis</i> Mez		x	Landell et al., 2006
	<i>Werauhia gigantea</i> (Mart. ex Schult. & Schult.f.) J.R.Grant		x	Landell et al., 2006
<i>Sporobolomyces shibatanus</i>	<i>Bromelia antiacantha</i> Bertol.		x	Landell et al., 2006
	<i>Tillandsia crocata</i> (É.Morren) N.E.Br.	x		Landell et al., 2006
	<i>Tillandsia geminiflora</i> Brongn.		x	Landell et al., 2006
	<i>Vriesea friburgensis</i> Mez		x	Landell et al., 2006
<i>Symmetrospora marina</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Morais et al., 2020
	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira		x	Navarro et al., 2020
	<i>Aechmea werdermannii</i> Harms		x	Navarro et al., 2020
	<i>Hohenbergia stellata</i> Schult. & Schult.f.		x	Navarro et al., 2020
<i>Symmetrospora suhii</i>	<i>Aechmea cephaloides</i> J.A.Siqueira & Leme		x	Navarro et al., 2020
	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira		x	Navarro et al., 2020
	<i>Canistrum aurantiacum</i> É.Morren		x	Navarro et al., 2020
	<i>Cryptanthus burle-marxii</i> Leme		x	Navarro et al., 2020
<i>Symmetrospora symmetricus</i>	<i>Hohenbergia ridleyi</i> (Baker) Mez		x	Navarro et al., 2020
<i>Yunzhangia sonckii</i>	<i>Vriesea friburgensis</i> Mez		x	Landell et al., 2006
Ustilaginomycotina (Subphyla)				
<i>Anomalomyces panici</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015; Morais et al., 2020
<i>Anomalomyces yakirrae</i>	<i>Vriesea minarum</i> L.B.Sm.		x	Morais et al., 2020
<i>Anthracocystis elionuri</i>	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
<i>Anthracocystis everhartii</i>	<i>Bromelia karatas</i> L.		x	Morais et al., 2020

<i>Cintractia samanensis</i>	<i>Vriesea minarum</i> L.B.Sm.			x	Morais et al., 2020
	<i>Dyckia</i> sp.		x		Inácio et al., 2008
	<i>Tillandsia gardneri</i> Lindl.		x		Inácio et al., 2008
<i>Farysia itapuensis</i>	<i>Tillandsia geminiflora</i> Brongn.		x		Inácio et al., 2008
	<i>Vriesea friburgensis</i> Mez		x		Inácio et al., 2008
	<i>Vriesea procera</i> (Mart. ex Schult. & Schult.f.) Wittm.		x		Inácio et al., 2008
<i>Gjaerumia minor</i>	<i>Vriesea minarum</i> L.B.Sm.			x	Morais et al., 2020
<i>Jaminaea rosea</i>	<i>Cryptanthus zonatus</i> (Vis.) Beer		x		Navarro et al., 2020
<i>Kalmanozyma brasiliensis</i>	<i>Encholirium</i> sp.			x	Morais et al., 2020
<i>Langdonia jejuensis</i>	<i>Vriesea minarum</i> L.B.Sm.			x	Morais et al., 2020
<i>Meira miltonrushii</i>	<i>Aechmea werdermannii</i> Harms		x		Navarro et al., 2020
	<i>Hohenbergia ridleyi</i> (Baker) Mez		x		Navarro et al., 2020
<i>Meira nashicola</i>	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira		x		Navarro et al., 2020
<i>Microstroma bacarum</i>	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.			x	Araujo et al., 1998
	<i>Tillandsia crocata</i> (É.Morren) N.E.Br.		x		Landell et al., 2006
	<i>Neoregelia cruenta</i> (Graham) L.B.Sm.			x	Hagler et al., 1993
<i>Moesziomyces antarcticus</i>	<i>Quesnelia quesneliana</i> (Brongn.) L.B.Sm.			x	Hagler et al., 1993
	<i>Hohenbergia stellata</i> Schult. & Schult.f.		x		Navarro et al., 2020
	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira	x	x		Félix et al., 2021; Navarro et al., 2020
	<i>Vriesea minarum</i> L.B.Sm.			x	Gomes et al., 2015
<i>Moesziomyces aphidis</i>	<i>Araeococcus chlorocarpus</i> (Wawra) Leme & J.A.Siqueira		x		Navarro et al., 2020
	<i>Bromelia karatas</i> L.		x		Navarro et al., 2020
	<i>Canistrum alagoanum</i> Leme & J.A.Siqueira		x		Navarro et al., 2020
	<i>Hohenbergia stellata</i> Schult. & Schult.f.		x		Navarro et al., 2020
<i>Pattersoniomyces tillandsiae</i>	<i>Canistrum improcerum</i> Leme & J.A.Siqueira		x		Piątek et al., 2017; Navarro et al., 2020

	<i>Tillandsia flabellata</i> Baker		x	Piątek et al., 2017
	<i>Tillandsia leiboldiana</i> Schleidl.		x	Piątek et al., 2017
	<i>Vriesea minarum</i> L.B.Sm.		x	Piątek et al., 2017
<i>Pseudozyma hubeiensis</i>	<i>Araeococcus chlorocarpus</i> (Wawra) Leme & J.A.Siqueira	x	x	Navarro et al., 2020; Félix et al., 2021
	<i>Cryptanthus dianae</i> Leme	x	x	Navarro et al., 2020; Félix et al., 2021
	<i>Vriesea minarum</i> L.B.Sm.		x	Gomes et al., 2015; Morais et al., 2020
	<i>Bromelia karatas</i> L.		x	Morais et al., 2020
	<i>Encholirium</i> sp.		x	Morais et al., 2020
	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira		x	Navarro et al., 2020
	<i>Aechmea werdermannii</i> Harms		x	Navarro et al., 2020
	<i>Ananas comosus</i> (L.) Merr.		x	Navarro et al., 2020
	<i>Billbergia</i> sp.		x	Navarro et al., 2020
	<i>Canistrum alagoanum</i> Leme & J.A.Siqueira		x	Navarro et al., 2020
	<i>Canistrum improcerum</i> Leme & J.A.Siqueira		x	Navarro et al., 2020
	<i>Hohenbergia stellata</i> Schult. & Schult.f.		x	Navarro et al., 2020
	<i>Tillandsia chapeuensis</i> Rauh		x	Navarro et al., 2020
	<i>Tillandsia kegeliana</i> Mez		x	Navarro et al., 2020
<i>Sympodiomyopsis paphiopedili</i>	<i>Aechmea leptantha</i> (Harms) Leme & J.A.Siqueira		x	Navarro et al., 2020
<i>Sympodiomyopsis yantaiensis</i>	<i>Aechmea froesii</i> (L.B.Sm.) Leme & J.A.Siqueira		x	Navarro et al., 2020
<i>Ustilago sparsa</i>	<i>Bromelia karatas</i> L.		x	Morais et al., 2020

4. DISCUSSION

4.1. Diversity

It is not the first time that bromeliads are mentioned as an environment rich in yeast diversity, e.g. (Morais et al., 2020; Navarro et al., 2020). However, these results present a more comprehensive and precise view of the size of this richness, as well as how it is distributed among the different compartments of the phyllosphere. The richest taxonomic groups (species number) in the bromeliad phyllosphere are groups commonly found in other plants, especially in leaves (Fonseca and Inácio, 2006), such as genera *Candida*, *Papiliotrema*, *Rhodotorula*, *Hannaella*, and the orders *Saccharomycetales*, *Tremellales*, *Sporidiobolales* and *Ustilaginales*. Regarding the frequency of occurrence of the genera, *Papiliotrema*, *Candida*, *Aureobasidium*, *Meyerozyma*, *Rhodotorula*, *Carlosrosaea*, *Hannaella* and *Pseudozyma* were the most frequent in bromeliads. At least one of these genera was found in about 70% of the analyzed units. The classically dominant genera in the phyllosphere are *Aureobasidium*, *Cryptococcus*, *Rhodotorula* and *Sporobolomyces* (Fonseca and Inácio, 2006; Vacher et al., 2016). Even today, it is not uncommon to mention *Cryptococcus* as one of the most frequent genera in plants (Vacher et al., 2016). Nevertheless, most species previously grouped in this genus have been reclassified and are distributed in other genera, such as *Kwoniella*, *Naganishia*, *Papiliotrema*, *Saitozyma* and *Vishniacozyma* (Liu et al., 2015). At least 17 species previously grouped in the genus *Cryptococcus* were transferred to *Papiliotrema* (Liu et al., 2015). Including species known to be frequent in plant substrates such as *P. flavesrens*, *P. laurentii* and *P. nemorosus* (Fonseca and Inácio, 2006). We reported genera that are conventionally found in the phyllosphere, though often not very expressive. For instance, ascomycete yeasts tend to be less frequent in the phyllosphere than basidiomycetes (Fonseca and Inácio, 2006). Nonetheless, in bromeliad flowers (Félix et al., 2021), fruits (Chanprasartsuk et al., 2010), leaves (Landell et al., 2006) and water tanks (Araujo et al., 1998; Morais et al., 2020), the most frequent groups found include the genus *Candida*, but also *Meyerozyma* and *Hanseniaspora*. Still, the genus *Pseudozyma* has already been frequently recorded in bromeliad water tanks in the Brazilian savanna (Morais et al., 2020). It stood out also as one of the most common genera in bromeliads in Northeastern Brazil, both in leaves and in flowers (Félix et al., 2021; Navarro et al., 2020). Nevertheless,

Pseudozyma was the most frequent in other Poales plants (e.g. rice and maize) (Nasanit et al., 2016, 2015). Another recurrent genus in bromeliads is *Carlosrosaea*, which currently has five described species, all associated with plant substrates. To date, three of them are exclusively associated with bromeliads: *Ca. vrieseae*, *Ca. aechmeae* and *Ca. hohenbergiae* (Félix et al., 2021).

There is evidence to suggest that the phyllosphere community is more resilient and tolerates disturbances generated by rough stresses, such as rainfall. In fact, it is more influenced by long-term events, for example, seasonality (Stone and Jackson, 2021). On the other hand, there are also works that point to more immediate changes in the phyllosphere community, mainly on fruit surfaces, in response to disruptive events (Allard et al., 2020). However, this pattern is not absolute and may vary depending on the plant species and compartment studied (Allard et al., 2020). The aerial part of plants is a dynamic environment mainly regulated by four population processes: i) immigration, ii) emigration, iii) growth (generation), and iv) death (Fonseca and Inácio, 2006; Kinkel, 1997). Specifically in bromeliads, the microbial community can vary taxonomically and be functionally stable, even at small spatial scales (Louca et al., 2017). Most species recorded in the bromeliad phyllosphere had a low frequency of occurrence: 65% had at most two occurrences in the analyzed units, 52% occurred in only one host species, while 70% occurred in a single compartment of the phyllosphere (flowers, fruits, leaves or water tank). Only five yeast species (3% of the total) had a frequency of occurrence greater than 10%: *P. laurentii*, *P. flavesrens*, *M. guilliermondii*, *C. intermedia*, *Ps. hubeiensis* and *A. pullulans*. The literature corroborates the low frequency of occurrence of yeasts in the phyllosphere, with most species found presenting a frequency of occurrence lower than 10%. Depending on the study, most may have a frequency lower than 1% (Glushakova et al., 2014; Glushakova and Chernov, 2010, 2007; Limtong and Kaewwichian, 2015; Nasanit et al., 2016). In bromeliads, the low frequency of occurrence and the large number of singletons have already been recorded in flowers, leaves and water tanks (Félix et al., 2021; Gomes et al., 2015; Morais et al., 2020; Navarro et al., 2020). This highlights that the richness of the phyllosphere tends to be mostly originated from rare groups. For example, the contribution of rare bacteria to the alpha and beta-diversity of the *Phragmites australis* phyllosphere is significantly greater than the contribution of abundant ones

(Zhou et al., 2019). If the intense dynamic of the phyllosphere accelerates the population processes that regulate it, the taxonomic shift of the phyllosphere can explain the low frequency of occurrence of the species, as a large part of the community members would be frequently changing. Nonetheless, this does mean that yeasts from the bromeliad phyllosphere are mostly transient. On the contrary, it shows that the community is dynamic and in a frequent turnover. Recent results point to a microbiota-sharing relationship among phyllosphere compartments and air, with leaves being the largest source of microorganisms. The air in this relationship could be compared to a microorganism 'stock' fed mainly by the plant compartments. Thus, the microbiota originating from the plant compartments is available and can, in a second moment, recolonize these compartments (Abdelfattah et al., 2019).

4.2. Yeasts in the different bromeliad compartments

Fonseca and Inácio (2006) mention that the phyllosphere yeast community tends to be more similar among leaves, intact fruits and flowers (when nectar is disregarded), especially concerning species composition and the prevalence of basidiomycetes yeasts. This prediction arises from the idea that these habitats are analogous as they share a similar configuration, since the interaction between the epiphytic microbiota and the plant cuticle is necessary in all these compartments. Our result partly confirms this statement, as among the compartments analyzed, the most similar communities were flowers and leaves; the only ones in which it is mandatory that yeasts interact directly with the cuticle. For instance, in fruits and water tanks, the microbial communities can interact with the plant without necessarily having any direct contact with the cuticle, as the tanks create an aquatic environment and used fruit samples are mainly dismantled, as discussed ahead. Regarding the diversity, leaves and water tanks were significantly more diverse than fruits and flowers. Furthermore, the composition was very distinct: 70% of all yeast species were exclusive to one of the compartments and only 2% were shared among all. The community structure followed the pattern and diverged between compartments, but inconclusively. However, it could be safely observed that leaves were the most heterogeneous compartment compared to the others. Using independent methods of cultivation, similar standards have been recently recorded, with few fungal OTUs shared among grape vineyards, fruit and flowers. In some

studies, the proportion of shared OTUs is as low as 0.3%, reinforcing how these compartments can be distinct (Abdelfattah et al., 2019; Liu and Howell, 2021).

4.3. Flowers

Flowers had the lowest richness of yeast species among the compartments, as well as the lowest proportion of exclusive species (18%). Moreover, they were the compartment with a community similarly related to leaves. It is important to emphasize that this compartment is, to date, the least studied within the bromeliad phyllosphere. Only one study focused on bromeliad flowers, and another sporadically collected a sample thereof. The most frequent species in this compartment were also common in leaves and tanks: *P. flavescens* and *C. intermedia*. Recent studies highlight that the most frequent species in flowers can vary greatly depending on the host and even between studies (Félix et al., 2021). Furthermore, only *Ph. bogoriensis* was a non-singleton exclusive to the flowers compartment. It was possible to observe that species richness in groups such as the genus *Sporobolomyces*, the class Microbotryomycetes and the subphylum Pucciniomycotina increased in flowers in relation to the other compartments. Abdelfattah et al. (2019) found a significantly greater abundance of *Sporobolomyces* in flowers than in other vine compartments. It is evident that to tolerate the extreme conditions of the phyllosphere, microorganisms must have efficient and versatile strategies that enable the colonization of this habitat. One of the explanations for the prevalence of some of the groups found relies on certain traits that make them more tolerant to phyllosphere stress. For example, when regarding the ability to produce pigments that are related to the tolerance to UV radiation, these pigments can be diverse, but they are often carotenoids that are widely found in members of Sporidiobolales (e.g. *Rhodotorula* and *Sporobolomyces*), or melanin, produced by black yeasts such as *Aureobasidium* and *Hortaea* (Fonseca and Inácio, 2006; Kemler et al., 2017). Furthermore, some species of *Sporobolomyces* and *Bullera* produce ballistospores, which are ejectable spores that have a dispersal function (Fonseca and Inácio, 2006; Kemler et al., 2017). Much of the literature usually mentions flowers as an environment dominated by ascomycetes yeasts and that the basidiomycetes yeasts found in this compartment are usually contaminating groups (Canto et al., 2017; Pozo et al., 2011). However, our results corroborate the

statement that when disregarding exudates such as nectar, at least in bromeliad flowers, most of the community consists of basidiomycetes. Nevertheless, the prevalence of basidiomycetes has been previously reported in hexose-dominant nectar flowers and flowers pollinated by birds (Mittelbach et al., 2015).

The anthoplane is a term that we will use to refer to the plant compartment that was analyzed in the study carried out by Félix et al. (2021), which is defined as the surface of the floral tissue after the removal (at least in a large part) of the floral exudates and consequently of the microbial community that accompanies them. So far, only Félix et al. (2021) has discussed this compartment in bromeliads as a unit (or subunit). Although Landell et al. (2006) have briefly isolated anthoplane yeasts, there was no mention of a distinction between the anthoplane and the anthosphere. Studies of yeast in flowers are commonly performed in anthosphere communities without distinguishing between the tissue surface and exudates, e.g. (Canto et al., 2017; Mittelbach et al., 2015; Pozo et al., 2011). There is not enough information in the literature to prove this, though the general idea is that the anthoplane is a subcompartment of the anthosphere. Thus, all microorganisms in the anthoplane are part of the anthosphere, but the opposite is not true. According to this logic, the prevalence of basidiomycetous yeasts would be characteristic of the anthoplane, as it has a similar configuration to the phylloplane (leaf), and of ascomycetes in the anthosphere, mainly due to the presence of nectar and other exudates that benefit specialists such as members of *Metschnikowiaceae* (Morris et al., 2020). This highlights that flowers are a complex substrate, with greater resemblance to a mosaic with several compartments instead of a single homogeneous compartment.

4.4. Fruits

Fruits are a complex and heterogeneous environment from a microbiological point of view (Ganter et al., 2017). For example, in pineapple fruits yeast density varies from 5.01 ± 0.05 to 4.0 ± 0.6 log CFU/g⁻¹. Density is higher in the fruit's outer ring and it decreases as the innermost regions are measured (Di Cagno et al., 2010). Similar to leaves, fruits are often dominated by oligotrophic and non-fermentative species of basidiomycete yeasts, mainly in immature and/or intact fruits (Fonseca and Inácio, 2006; Maksimova et al., 2009; Péter et al., 2017). However, among the bromeliad compartments, fruits were the most dissimilar to the leaves and

the only that showed the dominance of ascomycetes (78% of the species in this compartment), mainly from the Saccharomycetaceae family. A plausible explanation is that yeasts from bromeliad fruits were always perforated from ripe pineapple fruits (*Ananas comosus* (L.) Merr.), on various substrate types (fermented fruit, pulp, or peels). In addition, enrichment methods were used, which usually overrepresent ascomycetes yeasts. Basidiomycetes yeasts are usually more generalist and capable of assimilating several carbon sources (polytrophic), having an advantage in oligotrophic environments. Therefore, they are frequent and dominant in the phyllosphere, mainly directly associated with the cuticle (Félix et al., 2021; Fonseca and Inácio, 2006; Hagler et al., 1993; Yurkov, 2017).

Furthermore, in fruits, the natural process of senescence that leads to disruption, and later decay, mostly promotes changes in basidiomycetes, oligotrophic and non-fermentative communities, as well as in mostly ascomycetes, copiotrophic predominant communities and with fermenting yeasts (Ganter et al., 2017; Péter et al., 2017). This issue is strengthened as we observe that the most frequent yeasts in bromeliad fruits were ascomycetes fermenters, being frequently found in fruits (Ganter et al., 2017) such as *M. guilliermondii*, *H. uvarum* and *Pi. kudriavzevii*. Species related and exclusive to this bromeliad compartment, such as *C. sorboxylosa*, *Pi. kudriavzevii*, *Pi. fermentans* and *Saccharomyces cerevisiae* are also all fermenters. In turn, plant ascomycetes yeasts, mainly members of Saccharomycetaceae, tend to be specialists (assimilate a more restricted number of carbon sources), fast-growing and dominate copiotrophic environments (rich in simple sugars) such as nectar and ripe or decaying fruits (Félix et al., 2021; Fonseca and Inácio, 2006; Ganter et al., 2017; Péter et al., 2017). For example, approximately 85% of yeasts considered to be nectivores are fast-growing specialist ascomycetes (Pozo et al., 2011). While the surface-associated community of immature and/or intact fruit tends to undergo community displacements, the damaged or fermented fruit community tends to be more recurrent and stable (Fonseca and Inácio, 2006; Péter et al., 2017). For instance, in pineapple fruits, *M. guilliermondii* was the species that dominated the first days of fermentation, while towards the end of the process, the species *H. uvarum* became dominant. This pattern remained in independent collections in two continents (Chanprasartsuk et al., 2010). Moreover, a large number of yeasts capable of producing killer toxins have been found in fruits (mainly

decayed) (Ganter et al., 2017; Starmer and Lachance, 2011), which highlights the competitive character of the habitat, typical of copiotrophic environments (Félix et al., 2021).

4.5. Leaves

The surface of the leaves (phylloplane) is one of the largest habitats in the world and it has an estimated total surface area of 1 billion km²; equivalent to twice the surface area of the Earth (Bringel and Couée, 2015). Bromeliad leaves can vary in several aspects, from length, shape, color, as well as in terms of the serration of the edge. For example, while members of the genus *Bromelia* have a densely serrated edge, members of the subfamily *Tillandsioideae* have smooth edges (Benzing, 2000). Some species of bromeliads depend on the absorption of leaves to access water and nutrients, which reinforces the importance of the associated microbiota for the processes of decomposition and nutrient cycling (Leroy et al., 2016). In addition to the most common environmental disturbers that can affect the leaf yeast community such as rain, wind, and solar radiation (Fonseca and Inácio, 2006), other external factors can impact the community, e.g. epiphytic yeasts can be completely inhibited by air pollution on bromeliad leaves epiphytes of the genus *Tillandsia* (Brighigna et al., 2000). Most of the organic compounds made available to the epiphytic microbiota are made available spontaneously or non-spontaneously by the plant itself, while inorganic compounds are usually from an external source (Fonseca and Inácio, 2006; Kemler et al., 2017). As already mentioned, the phyllosphere (especially the phylloplane) is an extreme and oligotrophic environment, and this may be one of the factors that generate its great diversity. Copiotrophic environments (rich in simple sugars) are usually marked by a rapidly growing specialist community that, under competitive pressure, can diminish local diversity, and oligotrophic environments are marked by populations with lower abundances and higher diversity (Cray et al., 2013; Lachance, 2013; Félix et al. 2021). Furthermore, it is known that the yeast species shift is larger in these compartments in relation to the others (Fonseca and Inácio, 2006).

Compared to the other compartments analyzed (flowers, fruits, and water tank), leaves were the most heterogeneous (variance) and this compartment is the most perennial structures among the others, which can corroborate the heterogeneity

recorded. It is possible that the turnover of species, which can occur for seasonal reasons, host senescence, deterministic or stochastic factors, cause the community to vary over time more intensely than in structures that are restricted to a seasonal or reproductive period. Bromeliad leaves were one of the most diverse plant compartments and, as expected, the basidiomycetes were dominant. The genera *Papiliotrema* and *Carlosrosaea* stood out in this compartment in terms of species richness and the frequency of occurrence. The most frequent species in bromeliad leaves were all basidiomycetes: *P. laurentii*, *Ps. hubeiensis*, *Ca. hohenbergiae* and *P. leoncinii*. Finally, one of the patterns described regarding the richness of yeasts in bromeliad leaves indicates the existence of a significant and positive correlation between the diversity of bromeliads and associated yeasts. This pattern is probably a result of the greater diversity of hosts, which generates a more heterogeneous environment with a greater diversity of available niches (Navarro et al., 2020). However, no significant relationship was found between the beta-diversity of bromeliads and yeasts, which may indicate that there is no species-specific relationship between hosts and yeasts (Navarro et al., 2020).

4.6. Water tank

About half of the bromeliad genera and more than half of the subfamilies contain species capable of forming water tanks (phytotelma) (Ladino et al., 2019; Males and Griffiths, 2017). Depending on the species and environmental conditions, the phytotelm can accumulate from 0.015 to 45 L of water in a single individual (Zotz et al., 2020), and about 50,000 L in a hectare (Ladino et al., 2019). However, this compartment is an adverse environment mainly because it is ephemeral and dependent on external conditions, such as rain. Furthermore, evidence suggests less fluctuation in water conditions in plants with larger tanks and that accumulate more water (Zotz et al., 2020), although it is not known how this affects the phytotelm yeasts. The yeast richness in tanks was the highest among the compartments, closely followed by leaves. Moreover, more than half of the yeast species in bromeliad water tanks were basidiomycetes. The prevalence of basidiomycetes is probably related to the ability of this group to use different sources of nutrition (polytrophy) and to colonize oligotrophic environments, such as the case of water tanks. The polytrophic capacity of bromeliad phytotelma yeasts has already been

verified. According to Hagler et al. (1993), more than half of the yeasts found assimilated more than 20 different carbon sources. It is already known that the microbial community in the tank can vary geographically, interspecifically, and even intraspecifically (Hagler et al., 1993; Louca et al., 2017; Morais et al., 2020). There is evidence to indicate that this environment is extremely variable taxonomically, but at least in bacteria, it is functionally stable (Louca et al., 2017). The most frequent species in this compartment were *C. intermedia*, *R. mucilaginosa*, *P. laurentii* and *N. albida*. All these species are cosmopolitan, but often associated with plant substrates (Rosa and Péter, 2006).

4.7. Community structure

Evolutionary innovations, ecophysiological strategies, besides several water harvesting mechanisms, explain the wide spectrum of occurrence of bromeliads (Benzing, 2000; Crayn et al., 2015; Males and Griffiths, 2017). We can mention the absorbent trichomes of the leaves, the emergence of photosynthetic metabolism type CAM (Crassulacean Acid Metabolism), the capacity to retain water in the tank, and succulence as factors that contributed to the evolutionary success of bromeliads (Benzing, 2000; Crayn et al., 2015; Males and Griffiths, 2017). These traits are important for drought tolerance and, possibly, for the colonization of xerophic environments and high altitudes. Males and Griffths (2017) combined the phylogeny of bromeliads with five functional types previously described, based on the photosynthetic pattern and the water absorption mechanism associated with the bromeliad habit, namely: C₃ terrestrials, C₃ tank epiphytes, CAM atmospheric epiphytes, CAM terrestrials, and CAM tank epiphytes. Among these functional types, morphological, ecological and physiological variations were observed. In addition, there are differences in traits related to drought tolerance, such as water mass per unit area and osmotic potential at full turgor, for example (Males and Griffiths, 2017). Furthermore, these functional groups are indirect indicators of greater or lesser dependence on the root for water absorption. Simply speaking, CAM atmospheric epiphytes are the most adapted for absorption through leaf trichomes, while C₃ terrestrial plants are the most dependent on root absorption (Benzing, 2000; Leroy et al., 2016; Males and Griffiths, 2017). We observed that bromeliad subfamilies, functional types and compartments were shown to have some effect on the structure

of the associated yeast community. Nonetheless, the result was inconclusive since there was a variation in the results depending on the data analyzed. However, even considering the possibility of the significant results being maintained, the explanatory power of the variables was low. The R^2 of the bromeliad subfamily ranged from 3 to 11%, from 7 to 15.6% for the compartments, and from 6.8 to 15.3% for the functional types. The low values of R^2 in these categories may be associated with some classes that had high variance in the compartments, leaves were significantly more heterogeneous (greater variance) than flowers, fruits and water tanks. As previously mentioned, this high heterogeneity in leaves may be related to the shift of species that tends to be large in the phylloplane (Fonseca and Inácio, 2006).

4.8. New Yeast Species

More than twenty new yeast species were described from bromeliads between 2005 and 2020. Several studies on bromeliads have recorded many undescribed species associated with these plants. In bromeliad tanks, Gomes et al. (2015) found 10 new yeast species (28% of the total recorded by the authors); from leaves, Navarro et al. (2020) recorded 32 new species (38% of the total registered); and from flowers, Félix et al. (2021) registered six possible undescribed species (22% of the total registered). Considering all species recorded in bromeliads ($n = 181$), more than a tenth of these species had as initial isolation substrate bromeliads. Little is known about the diversity of existing fungi, only around 5%. Especially in South America, vascular plants are the largest reservoir on the planet for the discovery of new fungi, (Hawksworth and Lücking, 2017). A positive correlation was found between the number of bromeliad hosts and yeast species (Navarro et al., 2020), and about 44 species of bromeliads have been studied so far regarding the yeast community, which is equivalent to only 1% of the richness of bromeliads described. It is evident that we have barely scratched the surface in the knowledge of bromeliad-associated yeasts and that the number of described and undescribed species is even greater than we realize.

4.9. Biases

Some biases can influence this study, mainly because here we have gathered an array of information generated by methodologically heterogeneous

studies. Some possible sources of bias include the criteria used to obtain and select the articles, as well as the choice of databases, besides the different culture media and methodologies used in isolation. Furthermore, different studies showed different sample sizes and the number of hosts analyzed. Another source of bias is the yeast identification method. Older studies tend to use phenotypic identification methods that are known to underestimate the richness, ignoring cryptic species. In addition, some studies, in this case always focusing on fruits (all about pineapples), collected samples available in artificial environments such as markets or plantations. While this methodological difference is a disadvantage because it reduces the explanatory power of the analyses carried out, it is also an opportunity to discuss how these problems can be solved in future studies. For instance, classical (phenotypic) identification may underestimate richness and generate artificial results, but with increasingly more accessible molecular methods and next-generation sequencing (NGS) this type of problem tends to become less frequent.

The culture medium used to seed the samples can be chosen according to a variety of goals and rationales. Depending on the research, it may be a result of legacy issues. For example, the medium that has historically been used in a laboratory, or it may be just because it is the medium that is currently available. Whatever the reason, it is well known that choosing a methodology is, to some degree, choosing tolerable biases, as every methodology has its limitations and problems. For example, in the isolation of yeasts from bromeliads, several culture media have already been used: Fungal Plate Count Agar (PCA), Malt Extract Agar (MEA), Potato Dextrose Agar (PDA), Sabouraud Dextrose Agar (SDA), Wort Agar, Yeast Malt Agar (YM Agar) and Yeast Extract Peptone Dextrose Agar (YEPD Agar). In addition to this variety of media used, there may also be modifications such as acidification and the addition of antibiotics such as chloramphenicol, streptomycin and oxytetracycline. Most studies on bromeliads use YM Agar medium, often with the addition of some antibiotics. The other means mentioned are less often used. In addition to these points, the present study did not consider geoclimatic variables, which can be important predictors of the microbial community, as the special distribution of the studies evaluated was strongly asymmetric, with several gaps and most studies carried out in Brazil (about 65%), mostly in Atlantic Forest areas.

Another point to consider is that the data used in this study were collected until October 2021 and since then other studies have been published expanding the diversity of known yeasts in bromeliads. For example, Nutaratat et al. (2022) described the new genus *Savitreella* and the species *Savitreella phatthalungensis* and *Goffeauzyma siamensis* associated with pineapple leaves. In addition, the species *Sa. phatthalungensis* is the first taphrinomycete yeast recorded in bromeliads. Similarly, Bezerra et al. (2022) described *Valentiella maceioensis*, the new genus and species of black yeast-like associated with bromeliad leaves. Another work that recently described yeasts associated with bromeliad water tank and leaves was Navarro et al. (2022) who described *Tremella ananatis* and *Tremella lamprococci*.

5. CONCLUSIONS

Overall, it was possible to verify the great richness of yeasts associated with bromeliads. Nevertheless, it seems that we are still far from recovering the fullness of the diversity of yeasts in this substrate. However, with the frequent modernization of independent cultivation techniques, this goal may not be so far reaching. Moreover, the predominant yeast community from the bromeliad phyllosphere was similar to what is expected from the phyllosphere in general, though several expected patterns such as the prevalence of basidiomycetes in fruits were not confirmed. However, this is likely associated with the type of material and method applied in isolation. The yeast community diverged among compartments, thus establishing bromeliads as a complex and heterogeneous environment. Leaves and tanks were the most diverse compartments, which may be associated with the fact that leaves are heterogeneous environments (a point that was reaffirmed in this analysis) and tanks are ephemeral environments subject to large fluctuations. Furthermore, both environments are oligotrophic, which can be a factor that increases the richness of associated species.

The structure of the yeast community was related to the variables analyzed (functional bromeliad types, bromeliad subfamilies and compartment types), but all were low explanatory power. As already known, the dynamic of the

phylosphere microbiota is modulated by several factors, both deterministic and stochastic. This complex network of factors must be evaluated with more precise experimental designs so that the patterns can be deeply elucidated. The functional types of bromeliads and compartments seem to be useful and promising variables, which can be used as explanatory units in future studies. Nevertheless, the data used herein had several biases such as the variation in sample size, host and method of yeasts identification. Thus, in future studies, these variables should be controlled in an experimental design that allows verifying the real influence of the variables verified herein. Furthermore, further studies are needed to confirm the influence of scale on the processes and patterns of yeasts in bromeliads.

Bromeliads proved to be a potential substrate for prospecting new yeast species and a precious though still little explored source of biotechnological raw material. From the data collected, it was evident that there is still much to be explored in bromeliads. For instance, no studies were found on nectar. Among the data already available, those on flowers are still very incipient. In these last three decades (since the 90s) of studies on yeasts in bromeliads, most of the works focused on the descriptive survey of communities. The importance of this knowledge framework is undeniable, as these works provided the scientific community with all the knowledge obtained to date. It is necessary to take the next steps and increase the number of studies that seek to understand the processes that structure the epiphytic community and not just verify the patterns.

Acknowledgements

This work was supported by the Brazilian National Council for Scientific and Technological Development (CNPq, number of grants 475378/2013-0, 408718/2013-7, 441657/2016-8, 311553/2018-4, 312889/2021-6), Coordination for the Improvement of Higher Education (CAPES) for the funding granted. We thank PhD. Gustavo Paulino and Msc. Hector Maurício Casanova Navarro for their contribution to the revision of the text.

REFERENCES

- Abdelfattah, A., Sanzani, S.M., Wisniewski, M., Berg, G., Cacciola, S.O., Schena, L., (2019). Revealing Cues for Fungal Interplay in the Plant–Air Interface in Vineyards. *Front. Plant Sci.* 10, 1–10. <https://doi.org/10.3389/fpls.2019.00922>
- Allard, S.M., Ottesen, A.R., Micallef, S.A., (2020). Rain induces temporary shifts in epiphytic bacterial communities of cucumber and tomato fruit. *Sci. Rep.* 10, 1765. <https://doi.org/10.1038/s41598-020-58671-7>
- Amorim, J.C., Piccoli, R.H., Duarte, W.F., (2018). Probiotic potential of yeasts isolated from pineapple and their use in the elaboration of potentially functional fermented beverages. *Food Res. Int.* 107, 518–527. <https://doi.org/10.1016/j.foodres.2018.02.054>
- Araujo, F. V., Medeiros, R.J., Mendonça-Hagler, L.C., Hagler, A.N., (1998). A preliminary note on yeast communities of bromeliad-tank waters of Rio de Janeiro, Brazil. *Rev. Microbiol.* 29, 118–121.
- Araujo, F. V., Rosa, C.A., Freitas, L.F.D., Lachance, M.A., Vaughan-Martini, A., Mendonca-Hagler, L.C., Hagler, A.N., (2012). *Kazachstania bromeliacearum* sp. nov., a yeast species from water tanks of bromeliads. *Int. J. Syst. Evol. Microbiol.* 62, 1002–1006. <https://doi.org/10.1099/ijns.0.031633-0>
- Benzing, D.H., (2000). *Bromeliaceae: profile of an adaptive radiation*. Cambridge University Press, New York.
- Bezerra, J. D., Navarro, H., Almeida, J. H., Félix, C. R., Landell, M. F., (2022). *Valentiella maceioensis* gen. et sp. nov. (Herpotrichiellaceae, Chaetothyriales), a new black yeast-like fungus isolated from bromeliads in Brazil. *Mycological Progress*, 21(2), 1-9. <https://doi.org/10.1007/s11557-022-01783-3>
- Brighigna, L., Gori, A., Gonnelli, S., Favilli, F., (2000). The influence of air pollution on the phyllosphere microflora composition of *Tillandsia* leaves (Bromeliaceae). *Rev. Biol. Trop.* 48, 511–517. <https://doi.org/10.15517/rbt.v48i2-3.18821>
- Bringel, F., Couée, I., (2015). Pivotal roles of phyllosphere microorganisms at the interface between plant functioning and atmospheric trace gas dynamics. *Front. Microbiol.* 6, 1–14. <https://doi.org/10.3389/fmicb.2015.00486>
- Buck, J.W., (2002). In vitro antagonism of *Botrytis cinerea* by phylloplane yeasts. *Can. J. Bot.* 80, 885–891.
- Canto, A., Herrera, C.M., Rodriguez, R., (2017). Nectar-living yeasts of a tropical host plant community: diversity and effects on community-wide floral nectar traits. *PeerJ* 5, e3517. <https://doi.org/10.7717/peerj.3517>
- Chanprasartsuk, O. Prakitchaiwattana, C., Sanguandekul, R., (2013). Comparison of methods for identification of yeasts isolated during spontaneous fermentation of freshly crushed pineapple juices. *J. Agric. Sci. Technol.* 15, 1479–1490.

- Chanprasartsuk, O., Pheanudomkitlert, K., Toonwai, D., (2012). Pineapple wine fermentation with yeasts isolated from fruit as single and mixed starter cultures. *Asian J. Food Agro-Industry* 5, 104–111.
- Chanprasartsuk, O.O., Prakitchaiwattana, C., Sanguandeekul, R., Fleet, G.H., (2010). Autochthonous yeasts associated with mature pineapple fruits, freshly crushed juice and their ferments; and the chemical changes during natural fermentation. *Bioresour. Technol.* 101, 7500–7509.
<https://doi.org/10.1016/j.biortech.2010.04.047>
- Corte, L., di Cagno, R., Groenewald, M., Roscini, L., Colabella, C., Gobbetti, M., Cardinali, G., (2015). Phenotypic and molecular diversity of *Meyerozyma guilliermondii* isolates isolated from food and other environmental niches, hints for an incipient speciation. *Food Microbiol.* 48, 206–215.
<https://doi.org/10.1016/j.fm.2014.12.014>
- Cray, J.A., Bell, A.N., Bhaganna, P., Mswaka, A.Y., Timson, D. J., Hallsworth, J.E., (2013). The biology of habitat dominance; can microbes behave as weeds? *Microbial biotechnology*, 6(5), 453-492. <https://doi.org/10.1111/1751-7915.12027>
- Crayn, D.M., Winter, K., Schulte, K., Smith, J.A.C., (2015). Photosynthetic pathways in Bromeliaceae: phylogenetic and ecological significance of CAM and C 3 based on carbon isotope ratios for 1893 species. *Bot. J. Linn. Soc.* 178, 169–221.
<https://doi.org/10.1111/boj.12275>
- Crous, P.W. et al. (2019). Fungal planet description sheets: 868-950. *Persoonia Mol. Phylogeny Evol. Fungi* 42, 291–473. <https://doi.org/10.3767/persoonia.2019.42.11>
- Crous, P.W. et al. (2018). Fungal planet description sheets: 716–784. *Persoonia Mol. Phylogeny Evol. Fungi* 40, 240–393. <https://doi.org/10.3767/persoonia.2018.40.10>
- Dellacassa, E., Trenchs, O., Fariña, L., Debernardis, F., Perez, G., Boido, E., Carrau, F., (2017). Pineapple (*Ananas comosus* L. Merr.) wine production in Angola: Characterisation of volatile aroma compounds and yeast native flora. *Int. J. Food Microbiol.* 241, 161–167. <https://doi.org/10.1016/j.ijfoodmicro.2016.10.014>
- Di Cagno, R., Cardinali, G., Minervini, G., Antonielli, L., Rizzello, C.G., Ricciuti, P., Gobbetti, M., (2010). Taxonomic structure of the yeasts and lactic acid bacteria microbiota of pineapple (*Ananas comosus* L. Merr.) and use of autochthonous starters for minimally processing. *Food Microbiol.* 27, 381–389.
<https://doi.org/10.1016/j.fm.2009.11.012>
- Dixon, P., (2003). VEGAN, a package of R functions for community ecology. *J. Veg. Sci.* 14, 927–930.
- Félix, C.R., Andrade, D.A., Almeida, J.H., Navarro, H.M.C., Fell, J.W., Landell, M.F., (2020). *Vishniacozyma alagoana* sp. Nov. a tremellomycetes yeast associated with plants from dry and rainfall tropical forests. *Int. J. Syst. Evol. Microbiol.* 70, 3449–3454. <https://doi.org/10.1099/ijsem.0.004193>

- Félix, C.R., Navarro, H.M.C., Almeida, J.H., Landell, M.F., (2021). Behind the nectar : the yeast community in bromeliads inflorescences after the exudate removal. Mycol. Prog. 20, 1191–1202. [https://doi.org/https://doi.org/10.1007/s11557-021-01728-2](https://doi.org/10.1007/s11557-021-01728-2)
- Felix, C.R., Navarro, H.M.C., Paulino, G.V.B., Broetto, L., Landell, M.F., (2017). Carlosrosaea hohenbergiae sp. Nov. and Carlosrosaea aechmeae sp. nov., two tremellaceous yeasts isolated from bromeliads in north-eastern Brazil. Int. J. Syst. Evol. Microbiol. 67, 1752–1757. <https://doi.org/10.1099/ijsem.0.001856>
- Fonseca, Á., Inácio, J., (2006). Phylloplane Yeasts, in: Biodiversity and Ecophysiology of Yeasts. pp. 263–301.
- Formoso, A., Heidrich, D., Felix, C.R., Tenório, A.C.A.C., Leite, B.R.B.R., Pagani, D.M.D.M., Ortiz-Monsalve, S., Ramírez-Castrillón, M., Landell, M.F.M.F., Scroferneker, M.L.M.L., Valente, P., (2015). Enzymatic Activity and Susceptibility to Antifungal Agents of Brazilian Environmental Isolates of *Hortaea werneckii*. Mycopathologia 180, 345–352. <https://doi.org/10.1007/s11046-015-9920-3>
- Frank, J.H., Lounibos, L.P., (1987). Phytotelmata: swamps or islands? Florida Entomol. 70, 14–20.
- Ganter, P.F., Morais, P.B., Rosa, C.A., (2017). Yeasts in Cacti and Tropical Fruit, in: Yeasts in Natural Ecosystems: Diversity. pp. 225–264. <https://doi.org/10.1007/978-3-319-62683-3>
- Glushakova, A.M., Chernov, I.Y., (2010). Seasonal dynamics of the structure of epiphytic yeast communities. Microbiology 79, 830–839. <https://doi.org/10.1134/S0026261710060160>
- Glushakova, A.M., Chernov, I.Y., (2007). Seasonal dynamic of the numbers of epiphytic yeasts. Microbiology 76, 590–595. <https://doi.org/10.1134/S0026261707050128>
- Glushakova, A.M., Kachalkin, A. V, Chernov, I.Y., (2014). Yeasts in the flowers of entomophilic plants of the Moscow region. Microbiol. (Russian Fed. 83, 125–134. <https://doi.org/10.1134/S002626171402009X>
- Goffredi, S.K., Jang, G.E., Haroon, M.F., (2015). Transcriptomics in the tropics: Total RNA-based profiling of Costa Rican bromeliad-associated communities. Comput. Struct. Biotechnol. J. 13, 18–23. <https://doi.org/10.1016/j.csbj.2014.12.001>
- Gomes, F.C.O., Safar, S.V.B., Marques, A.R., Medeiros, A.O., Santos, A.R.O., Carvalho, C., Lachance, M.-A., Sampaio, J.P., Rosa, C.A., (2015). The diversity and extracellular enzymatic activities of yeasts isolated from water tanks of *Vriesea minarum*, an endangered bromeliad species in Brazil, and the description of *Occultifur brasiliensis* f.a., sp. nov. Antonie Van Leeuwenhoek 107, 597–611. <https://doi.org/10.1007/s10482-014-0356-4>
- Gomes, F.C.O., Safar, S.V.B., Santos, A.R.O., Lachance, M.A., Rosa, C.A., (2016). *Kockovaella libkindii* sp. Nov., a yeast species isolated from water tanks of bromeliad. Int. J. Syst. Evol. Microbiol. 66, 5066–5069. <https://doi.org/10.1099/ijsem.0.001471>

- Hagler, A.N., Rosa, C.A., Morais, P.B., Mendonca-Hagler, L.C., (1993). Yeasts and coliform bacteria of water accumulated in bromeliads of mangrove and sand dune ecosystems of southeast Brazil. *Can. J. Microbiol.* 39, 973–977.
<https://doi.org/10.1139/m93-146>
- Hammer, Ø., Harper, D.A.T. a. T., Ryan, P.D., (2001). PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontol. Electron.* 4(1), 1–9. <https://doi.org/10.1016/j.bcp.2008.05.025>
- Hawksworth, D.L., Lücking, R., (2017). Fungal Diversity Revisited: 2.2 to 3.8 Million Species, in: *The Fungal Kingdom*. ASM Press, Washington, DC, USA, pp. 79–95. <https://doi.org/10.1128/9781555819583.ch4>
- Hsieh, T.C., Ma, K.H., Chao, A., (2016). iNEXT: an R package for rarefaction and extrapolation of species diversity. *Methods Ecol. Evol.* 7, 1451–1456.
<https://doi.org/10.1111/2041-210X.12613>
- Inácio, J., Landell, M.F., Valente, P., Wang, P.H., Wang, Y.T., Yang, S.H., Manson, J.S., Lachance, M.A., Rosa, C.A., Fonseca, Á., (2008). *Farysizyma* gen. nov., an anamorphic genus in the Ustilaginales to accommodate three novel epiphytic basidiomycetous yeast species from America, Europe and Asia. *FEMS Yeast Res.* 8, 499–508. <https://doi.org/10.1111/j.1567-1364.2008.00377.x>
- Islam, F., Salam, M.A., Rahman, M.A., Paul, S.I., Das, T.R., Rahman, M.M., Shaha, D.C., Gupta, D.R., Alam, M.S., Islam, T., (2021). Plant endophytic yeasts *Pichia fermentans* and *Meyerozyma caribbica* improve growth, biochemical composition, haematological parameters and morphology of internal organs of premature *Barbonymus gonionotus*. *Aquac. Reports* 19, 1–13.
<https://doi.org/10.1016/j.aqrep.2020.100575>
- Kemler, M., Witfeld, F., Begerow, D., Yurkov, A., (2017). Yeasts in Natural Ecosystems: Diversity. Springer International Publishing, Cham.
<https://doi.org/10.1007/978-3-319-62683-3>
- Khunnamwong, P., Ribeiro, J.R.A., Garcia, K.M., Hagler, A.N., Takashima, M., Ohkuma, M., Endoh, R., Sugita, T., Jindamorakot, S., Limtong, S., (2017). *Occultifur plantarum* f.a., sp. nov., a novel cystobasidiomycetous yeast species. *Int. J. Syst. Evol. Microbiol.* 67, 2628–2633. <https://doi.org/10.1099/ijsem.0.001988>
- Kinkel, L.L., (1997). Microbial population dynamics on leaves. *Annu. Rev. Phytopathol.* 35, 327–347. <https://doi.org/10.1146/annurev.phyto.35.1.327>
- Korres, A.M.N., Buss, D.S., Ventura, J.A., Fernandes, P.M.B., (2011). *Candida krusei* and *Kloeckera apiculata* inhibit the causal agent of pineapple fusariosis, *Fusarium guttiforme*. *Fungal Biol.* 115, 1251–1258. <https://doi.org/10.1016/j.funbio.2011.09.001>
- Korres, A.M.N., Ventura, J.A., Fernandes, P.M.B., (2010). First Report of *Bacterium* and Yeasts Associated with Pineapple Fruit Collapse in Espírito Santo State, Brazil. *Plant Dis.* 94, 1509–1509. <https://doi.org/10.1094/PDIS-04-10-0276>

- Kurtzman, C.P., (2011). Phylogeny of the ascomycetous yeasts and the renaming of *Pichia anomala* to *Wickerhamomyces anomalus*. *Antonie van Leeuwenhoek, Int. J. Gen. Mol. Microbiol.* 99, 13–23. <https://doi.org/10.1007/s10482-010-9505-6>
- Lachance, M.A. (2013). The biodiversity, ecology, and biogeography of ascomycetous yeasts. *The Ecological Genomics of Fungi*, 355-370. <https://doi.org/10.1002/9781118735893.ch16>
- Ladino, G., Ospina-Bautista, F., Estévez Varón, J., Jerabkova, L., Kratina, P., (2019). Ecosystem services provided by bromeliad plants: A systematic review. *Ecol. Evol.* 9, 7360–7372. <https://doi.org/10.1002/ece3.5296>
- Landell, M.F., Billodre, R., Ramos, J.P., Leoncini, O., Vainstein, M.H., Valente, P., (2010). *Candida aechmeae* sp. nov. and *Candida vriesiae* sp. nov., novel yeast species isolated from the phylloplane of bromeliads in Southern Brazil. *Int. J. Syst. Evol. Microbiol.* 60, 244–248. <https://doi.org/10.1099/ijns.0.011577-0>
- Landell, M.F., Brandão, L.R., Barbosa, A.C., Ramos, J.P., Safar, S.V.B., Gomes, F.C.O., Sousa, F.M.P., Morais, P.B., Broetto, L., Leoncini, O., Ribeiro, J.R., Fungsin, B., Takashima, M., Nakase, T., Lee, C.F., Vainstein, M.H., Fell, J.W., Scorzetti, G., Vishniac, H.S., Rosa, C.A., Valente, P., (2014). *Hannaella pagnoccae* sp. nov., a tremellaceous yeast species isolated from plants and soil. *Int. J. Syst. Evol. Microbiol.* 64, 1970–1977. <https://doi.org/10.1099/ijns.0.059345-0>
- Landell, M.F., Brandão, L.R., Safar, S.V.B., Gomes, F.C.O., Fèlix, C.R., Santos, A.R.O., Pagani, D.M., Ramos, J.P., Broetto, L., Mott, T., Vainstein, M.H., Valente, P., Rosa, C.A., (2015). *Bullera vriesiae* sp. Nov., a tremellaceous yeast species isolated from bromeliads. *Int. J. Syst. Evol. Microbiol.* 65, 2466–2471. <https://doi.org/10.1099/ijns.0.000285>
- Landell, M.F., Inácio, J., Fonseca, Á., Vainstein, M.H., Valente, P., (2009). *Cryptococcus bromelialarum* sp. nov., an orange-coloured basidiomycetous yeast isolated from bromeliads in Brazil. *Int. J. Syst. Evol. Microbiol.* 59, 910–913. <https://doi.org/10.1099/ijns.0.005652-0>
- Landell, M.F., Mautone, J.N., Valente, P., (2006). Biodiversity of Yeasts Associated To Bromeliads in Itapuã Park , Viamão / RS. *Biociencias* 14, 144–149.
- Leneveu-jenvrin, C., Quentin, B., Assemat, S., Hoarau, M., Meile, J.C., Remize, F., (2020). Changes of quality of minimally-processed pineapple (*Ananas comosus*, var. ‘queen victoria’) during cold storage: Fungi in the leading role. *Microorganisms* 8. <https://doi.org/10.3390/microorganisms8020185>
- Leroy, C., Carrias, J.-F., Cérégino, R., Corbara, B., (2016). The contribution of microorganisms and metazoans to mineral nutrition in bromeliads. *J. Plant Ecol.* 9, 241–255. <https://doi.org/10.1093/jpe/rtv052>
- Limtong, S., Kaewwichian, R., (2015). The diversity of culturable yeasts in the phylloplane of rice in Thailand. *Ann. Microbiol.* 65, 667–675. <https://doi.org/10.1007/s13213-014-0905-0>

- Limtong, S., Koowadjanakul, N., (2012). Yeasts from phylloplane and their capability to produce indole-3-acetic acid. *World J. Microbiol. Biotechnol.* 28, 3323–3335. <https://doi.org/10.1007/s11274-012-1144-9>
- Liu, D., Howell, K., (2021). Community succession of the grapevine fungal microbiome in the annual growth cycle. *Environ. Microbiol.* 23, 1842–1857. <https://doi.org/10.1111/1462-2920.15172>
- Liu, X.Z., Wang, Q.M., Göker, M., Groenewald, M., Kachalkin, A. V., Lumbsch, H.T., Millanes, A.M., Wedin, M., Yurkov, A.M., Boekhout, T., Bai, F.Y., (2015). Towards an integrated phylogenetic classification of the Tremellomycetes. *Stud. Mycol.* 81, 85–147. <https://doi.org/10.1016/j.simyco.2015.12.001>
- Louca, S., Jacques, S.M.S., Pires, A.P.F., Leal, J.S., Srivastava, D.S., Parfrey, L.W., Farjalla, V.F., Doebeli, M., (2017). High taxonomic variability despite stable functional structure across microbial communities. *Nat. Ecol. Evol.* 1, 0015. <https://doi.org/10.1038/s41559-016-0015>
- Maksimova, I.A., Yurkov, A.M., Chernov, I.Y., (2009). Spatial structure of epiphytic yeast communities on fruits of *Sorbus aucuparia* L. *Biol. Bull.* 36, 613–618. <https://doi.org/10.1134/S1062359009060107>
- Males, J., Griffiths, H., (2017). Functional types in the Bromeliaceae: relationships with drought-resistance traits and bioclimatic distributions. *Funct. Ecol.* 31, 1868–1880. <https://doi.org/10.1111/1365-2435.12900>
- Marques, A.R., Resende, A.A., Gomes, F.C.O., Santos, A.R.O., Rosa, C.A., Duarte, A.A., de Lemos-Filho, J.P., dos Santos, V.L., (2021). Plant growth-promoting traits of yeasts isolated from the tank bromeliad *Vriesea minarum* L.B. Smith and the effectiveness of *Carlosrosaea vriesiae* for promoting bromeliad growth. *Brazilian J. Microbiol.* 52, 1417–1429. <https://doi.org/10.1007/s42770-021-00496-1>
- Mittelbach, M., Yurkov, A.M., Nocentini, D., Nepi, M., Weigend, M., Begerow, D., (2015). Nectar sugars and bird visitation define a floral niche for basidiomycetous yeast on the Canary Islands. *BMC Ecol.* 15, 2. <https://doi.org/10.1186/s12898-015-0036-x>
- Morais, P.B., de Sousa, F.M.P., Rosa, C.A., (2020). Yeast in plant phytotelmata: Is there a “core” community in different localities of rupestrian savannas of Brazil? *Brazilian J. Microbiol.* 51, 1209–1218. <https://doi.org/10.1007/s42770-020-00286-1>
- Morris, M.M., Frixione, N.J., Burkert, A.C., Dinsdale, E.A., Vannette, R.L., (2020). Microbial abundance, composition, and function in nectar are shaped by flower visitor identity. *FEMS Microbiol. Ecol.* 96, 1–14. <https://doi.org/10.1093/femsec/fiaa003>
- Nagy, L.G., Ohm, R. a, Kovács, G.M., Floudas, D., Riley, R., Gácser, A., Sipiczki, M., Davis, J.M., Doty, S.L., de Hoog, G.S., Lang, B.F., Spatafora, J.W., Martin, F.M., Grigoriev, I. V, Hibbett, D.S., (2014). Latent homology and convergent regulatory evolution underlies the repeated emergence of yeasts. *Nat. Commun.* 5, 4471. <https://doi.org/10.1038/ncomms5471>

Nasanit, R., Jaibangyang, S., Tantirungkij, M., Limtong, S., (2016). Yeast diversity and novel yeast D1/D2 sequences from corn phylloplane obtained by a culture-independent approach. Antonie Van Leeuwenhoek. <https://doi.org/10.1007/s10482-016-0762-x>

Nasanit, R., Tangwong-o-thai, A., Tantirungkij, M., Limtong, S., (2015). Assessment of epiphytic yeast diversity in rice (*Oryza sativa*) phyllosphere in Thailand by a culture-independent approach. Antonie Van Leeuwenhoek 119, 1145–1157. <https://doi.org/10.1016/j.funbio.2015.08.021>

Nasir, A., Rahman, S.S., Hossain, M.M., Choudhury, N., (2017). Isolation of *Saccharomyces cerevisiae* from pineapple and orange and study of metal's effectiveness on ethanol production . Eur. J. Microbiol. Immunol. 7, 76–91. <https://doi.org/10.1556/1886.2016.00035>

Navarro, H.M.C., Félix, C.R., Paulino, G.V.B., Almeida, J.H., Valente, P., Landell, M.F., (2020). Richness and biotechnological potential of the yeast community associated with the bromeliad phylloplane in the Brazilian Neotropical Forest. Mycol. Prog. 19, 1387–1401. <https://doi.org/10.1007/s11557-020-01631-2>

Navarro, H. M., Félix, C. R., Tavares, V. D., de Sousa, F. M., Santos, A. R., Morais, P. B., Rosa, C.A., Valente, P., Landell, M. F., (2022). *Tremella ananatis* sp. nov. and *Tremella lamprococci* sp. nov., two yeast species associated with bromeliads. International Journal of Systematic and Evolutionary Microbiology, 72(2), 005261. <https://doi.org/10.1099/ijsem.0.005261>

Nutaratat, P., Boontham, W., Khunnamwong, P., (2022). A Novel Yeast Genus and Two Novel Species Isolated from Pineapple Leaves in Thailand: *Savitreella phatthalungensis* gen. nov., sp. nov. and *Goffeauzyma siamensis* sp. nov. Journal of Fungi, 8(2), 118. <https://doi.org/10.3390/jof8020118>

Pagani, D.M., Brandão, L.R., Santos, A.R.O., Felix, C.R., Ramos, J.P., Broetto, L., Scorzetti, G., Fell, J.W., Rosa, C.A., Valente, P., Landell, M.F., (2016). *Papiliotrema leoncinii* sp. nov. and *Papiliotrema miconiae* sp. nov., two tremellaceous yeast species from Brazil. Int. J. Syst. Evol. Microbiol. 66, 1799–1806. <https://doi.org/10.1099/ijsem.0.000945>

Péter, G., Takashima, M., Čadež, N., (2017). Yeast Habitats: Different but Global, in: Yeasts in Natural Ecosystems: Ecology. pp. 39–71. <https://doi.org/10.1007/978-3-319-61575-2>

Piątek, M., Lutz, M., Sousa, F.M.P., Santos, A.R.O., Félix, C.R., Landell, M.F., Gomes, F.C.O., Rosa, C.A., (2017). *Pattersoniomyces tillandsiae* gen. et comb. nov.: linking sexual and asexual morphs of the only known smut fungus associated with Bromeliaceae. Org. Divers. Evol. 17, 531–543. <https://doi.org/10.1007/s13127-017-0340-8>

Pozo, M.I., Herrera, C.M., Bazaga, P., (2011). Species Richness of Yeast Communities in Floral Nectar of Southern Spanish Plants. Microb. Ecol. 61, 82–91. <https://doi.org/10.1007/s00248-010-9682-x>

- R Team, (2016). R: A language and environment for statistical computing.
- Redford, A.J., Bowers, R.M., Knight, R., Linhart, Y., Fierer, N., (2010). The ecology of the phyllosphere: Geographic and phylogenetic variability in the distribution of bacteria on tree leaves. *Environ. Microbiol.* <https://doi.org/10.1111/j.1462-2920.2010.02258.x>
- Reyes, M.E.Q., Rohrbach, K.G., Paull, R.E., (2004). Microbial antagonists control postharvest black rot of pineapple fruit. *Postharvest Biol. Technol.* 33, 193–203. <https://doi.org/10.1016/j.postharvbio.2004.02.003>
- Rosa, C.A., Péter, G., (2006). 'The yeast handbook - Biodiversity and ecophysiology of yeasts,' 'The yeast handbook - Biodiversity and ecophysiology of yeasts.' <https://doi.org/10.1007/3-540-30985-3>
- Ruivo, C.C.C., Lachance, M.A., Rosa, C.A., Bacci, M., Pagnocca, F.C., (2005). *Candida bromeliacearum* sp. nov. and *Candida ubatubensis* sp. nov., two yeast species isolated from the water tanks of *Canistropsis seidelii* (Bromeliaceae). *Int. J. Syst. Evol. Microbiol.* 55, 2213–2217. <https://doi.org/10.1099/ijss.0.63698-0>
- Safar, S.V.B., Gomes, F.C.O., Marques, A.R., Lachance, M.A., Rosa, C.A., (2013). *Kazachstania rupicola* sp. nov., a yeast species isolated from water tanks of a bromeliad in Brazil. *Int. J. Syst. Evol. Microbiol.* 63, 1165–1168. <https://doi.org/10.1099/ijss.0.048462-0>
- Siqueira-Filho, J. A.; Leme, E.M.C., 2006. Fragmentos de Mata Atlântica do Nordeste - Biodiversidade, Conservação e suas Bromélias.
- Sousa, F.M.P., Morais, P.B., Lachance, M.A., Rosa, C.A., (2014). *Hagleromyces* gen. nov., a yeast genus in the Saccharomycetaceae, and description of *Hagleromyces aurorensis* sp. nov., isolated from water tanks of bromeliads. *Int. J. Syst. Evol. Microbiol.* 64, 2915–2919. <https://doi.org/10.1099/ijss.0.063883-0>
- Starmer, W.T., Lachance, M.-A., (2011). Yeast Ecology, in: *The Yeasts*. Elsevier, pp. 65–83. <https://doi.org/10.1016/B978-0-444-52149-1.00006-9>
- Stone, B.W.G., Jackson, C.R., (2021). Seasonal Patterns Contribute More Towards Phyllosphere Bacterial Community Structure than Short-Term Perturbations. *Microb. Ecol.* 81, 146–156. <https://doi.org/10.1007/s00248-020-01564-z>
- Tangsombatvichit, P., Pisapak, K., Suksaard, P., (2020). The natural lipolytic yeast *Candida* sp. Rmutsb-27 isolated from pineapple for treatment of cooking oil contaminated wastewater. *Environment Asia* 13, 70–79. <https://doi.org/10.14456/ea.2020.43>
- Tschapka, M., Von Helversen, O., (2007). Phenology, nectar production and visitation behaviour of bats on the flowers of the bromeliad *Werauhia gladioliflora* in a Costa Rican lowland rain forest. *J. Trop. Ecol.* 23, 385–395. <https://doi.org/10.1017/S0266467407004129>

- Udota, H.I.J., Urúa, E.E., (2010). Comparative analysis of fungal growth in commercially and laboratory prepared fruit juices - Using orange and pineapple as a case study. *J. Ind. Pollut. Control* 26, 125–130.
- Vacher, C., Hampe, A., Porté, A.J., Sauer, U., Compant, S., Morris, C.E., (2016). The Phyllosphere: Microbial Jungle at the Plant–Climate Interface. *Annu. Rev. Ecol. Evol. Syst.* 47, 1–24. <https://doi.org/10.1146/annurev-ecolsys-121415-032238>
- Valente, P., Boekhout, T., Landell, M.F., Crestani, J., Pagnocca, F.C., Sette, L.D., Passarini, M.R.Z., Rosa, C.A., Brandão, L.R., Pimenta, R.S., Ribeiro, J.R., Garcia, K.M., Lee, C.F., Suh, S.O., Péter, G., Dlauchy, D., Fell, J.W., Scorzetti, G., Theelen, B., Vainstein, M.H., (2012). *Bandoniozyma* gen. nov., a Genus of Fermentative and Non-Fermentative Tremellaceous Yeast Species. *PLoS One* 7. <https://doi.org/10.1371/journal.pone.0046060>
- Vorholt, J.A., (2012). Microbial life in the phyllosphere. *Nat. Rev. Microbiol.* 10, 828–840. <https://doi.org/10.1038/nrmicro2910>
- Vu, D., Groenewald, M., Szöke, S., Cardinali, G., Eberhardt, U., Stielow, B., de Vries, M., Verkleij, G.J.M., Crous, P.W., Boekhout, T., Robert, V., (2016). DNA barcoding analysis of more than 9 000 yeast isolates contributes to quantitative thresholds for yeast species and genera delimitation. *Stud. Mycol.* 85, 91–105. <https://doi.org/10.1016/j.simyco.2016.11.007>
- Wang, Q., Yurkov, A.M., Göker, M., Lumbsch, H.T., Leavitt, S.D., Groenewald, M., Theelen, B., Liu, X., Boekhout, T., Bai, F., (2016). Phylogenetic classification of yeasts and related taxa within Pucciniomycotina. *Stud. Mycol.* 81, 149–189. <https://doi.org/10.1016/j.simyco.2015.12.002>
- Wang, Q.M., Begerow, D., Groenewald, M., Liu, X.Z., Theelen, B., Bai, F.Y., Boekhout, T., (2015). Multigene phylogeny and taxonomic revision of yeasts and related fungi in the Ustilaginomycotina. *Stud. Mycol.* 81, 55–83. <https://doi.org/10.1016/j.simyco.2015.10.004>
- Westgate, M.J., (2019). revtools: An R package to support article screening for evidence synthesis. *Res. Synth. Methods* 10, 606–614. <https://doi.org/10.1002/jrsm.1374>
- Whipps, J.M., Hand, P., Pink, D., Bending, G.D., (2008). Phyllosphere microbiology with special reference to diversity and plant genotype. *J. Appl. Microbiol.* 105, 1744–1755. <https://doi.org/10.1111/j.1365-2672.2008.03906.x>
- Wilke, C.O., (2016). cowplot: streamlined plot theme and plot annotations for “ggplot2.” R Packag. version 0.6.
- Wolowski, M., Freitas, L., (2015). An overview on pollination of the Neotropical Poales. *Rodriguésia* 66, 329–336. <https://doi.org/10.1590/2175-7860201566204>
- Yurkov, A., 2017. Yeast Community Composition and Structure 73–100. <https://doi.org/10.1007/978-3-319-61575-2>

Yurkov, A., Alves, A., Bai, F.Y., Boundy-Mills, K., Buzzini, P., Čadež, N., Cardinali, G., Casaregola, S., Chaturvedi, V., Collin, V., Fell, J.W., Girard, V., Groenewald, M., Hagen, F., Hittinger, C.T., Kachalkin, A.V., Kostrzewa, M., Kouvelis, V., Libkind, D., Liu, X., Maier, T., Meyer, W., Péter, G., Piątek, M., Robert, V., Rosa, C.A., Sampaio, J.P., Sipiczki, M., Stadler, M., Sugita, T., Sugiyama, J., Takagi, H., Takashima, M., Turchetti, B., Wang, Q.M., Boekhout, T., (2021). Nomenclatural issues concerning cultured yeasts and other fungi: why it is important to avoid unneeded name changes. *IMA fungus*, 12(1), 1-20. <https://doi.org/10.1186/s43008-021-00067-x>

Zhou, Q., Zhang, X., He, R., Wang, S., Jiao, C., Huang, R., He, X., Zeng, J., Zhao, D., (2019). The Composition and Assembly of Bacterial Communities across the Rhizosphere and Phyllosphere Compartments of *Phragmites australis*. *Diversity* 11, 98. <https://doi.org/10.3390/d11060098>

Zotz, G., Leja, M., Aguilar-Cruz, Y., Einzmann, H.J.R., (2020). How much water is in the tank? An allometric analysis with 205 bromeliad species. *Flora Morphol. Distrib. Funct. Ecol.* Plants 264, 151557. <https://doi.org/10.1016/j.flora.2020.151557>

1 Periódico proposto: *FEMS Microbiology Ecology*

2 **5. CAPÍTULO 2 - Diversity, rainfall pulses and memory effect: A look**
3 **at yeast-plant system from Brazilian Tropical Semiarid Dryland**

4

5 Ciro Ramon Félix (ORCID 0000-0001-5087-8872)^{1,2}, Bruno Emanuel da Silva
6 Nascimento (ORCID 0000-0001-6690-0595)², Victor Daniel Firmino dos Santos
7 Tavares (ORCID 0000-0003-1492-0097)², Melissa Fontes Landell^{1*} (ORCID 0000-
8 0001-6848-0803)

9

10

11

12 ¹Programa de Pós-graduação em Diversidade Biológica e Conservação nos
13 Trópicos, Universidade Federal de Alagoas, Maceió – AL, Brazil.

14 ²Universidade Federal de Alagoas, Instituto de Ciências Biológicas e da Saúde,
15 Maceió – AL, Brazil.

16

17

18

19 *Correspondence: Melissa Fontes Landell, Setor de Genética - ICBS, Universidade
20 Federal de Alagoas, Av. Lourival Melo Mota, s/n, Tabuleiro dos Martins, CEP: 57072-
21 900, Maceió - AL, Brasil, Tel. (+55) 82-32141995 (e-mail:
22 melissa.landell@gmail.com)

23

24 Abstract

25
26 In drylands, rainfall occurs in pulses (rare events of resource over-availability) and
27 the drought that precedes it affects the intensity of the biological response during the
28 pulse (memory effect). The phyllosphere is the plant aboveground region, it's the
29 gateway to plant-environment interaction and a microbial megadiverse environment.
30 In both ecosystems, water deficit is recurrent and has wide implications for biological
31 communities. Here, we aimed to verify the rainfall, seasonality and environmental
32 memory effect on the diversity and community structure of phyllosphere yeasts in a
33 Brazilian dryland. While taxonomic, phylogenetic and functional alpha-diversity did
34 not differ significantly between seasonal periods, rain, drought and memory had little
35 influence on alpha-diversity but were significantly related to community structure. The
36 taxonomic composition, although functionally stable, diverged significantly over
37 seasons and presented high turnover values. Most of the functional traits did not
38 differ between periods, however there was a tendency towards greater frequency
39 during the rainy season. Our results bring insights into phyllosphere yeasts dynamics
40 in a tropical dryland and how it relates to rainfall pulses, seasonality, and memory
41 effect. In addition, by understanding the effect of dry-wet cycles on the microbiota of
42 the semiarid phyllosphere, we can predict effects on host health due to microbial
43 dysbiosis.

44

45 **Keywords:** Caatinga, seasonality, bromeliad, leaf, phyllosphere and turnover.

46 Introduction

47 Drylands are fundamentally regions with an aridity index (AI) < 0.65, that is,
48 with a high water-deficit (Noy-Meir 1973; Huang *et al.* 2016). These areas, which
49 cover more than 41% of the planet and are considered fragile ecosystems, generally
50 have unique biodiversity and provide important ecosystem services in the global
51 carbon, water and nitrogen cycles, and climate regulation (Huang *et al.* 2016;
52 Maestre *et al.* 2016, 2021). According to Noy-Meir (1973), the three most important
53 attributes for understanding the drylands ecosystem functioning are: i) The
54 precipitation is scarce and, therefore, a dominant factor in the control of biological
55 processes, ii) occurs occasionally and modestly, iii) and is variable and
56 unpredictable.

57 In the context of pulse dynamics, a pulse is defined as an ephemeral event of
58 resource overabundance (Holt 2008; Yang *et al.* 2008; Collins *et al.* 2014). The
59 pulses are a perturbation that generates a transient alternate state (Holt 2008). In
60 response to resource pulses, species can alter their dynamics (e.g. a spike in seed
61 recruitment or microbial metabolic rate) (Holt 2008; Yang *et al.* 2008; Collins *et al.*
62 2014). Although, resource pulses are short-lived events, they can be important
63 triggers that initiate powerful processes with persistent ecological effects long after
64 the pulse ends (Austin *et al.* 2004; Yang *et al.* 2008). In drylands, it can be
65 considered that rainfall inflows occur in 'pulses' (Noy-Meir 1973; Collins *et al.* 2014).
66 The rainfall pulse can influence other processes such as the translocation of
67 nutrients, disruption of the soil surface, redox patterns and the organic matter
68 production, deposition and decomposition (Austin *et al.* 2004; Collins *et al.* 2014).

69 The long periods of drought in drylands create reservoirs of mineral and
70 organic substrates because growth of plants and microorganisms is limited during
71 this time, generating little demand for nutrients, and lower rates of decomposition and
72 cycling (Austin *et al.* 2004; Holt 2008). This reservoir system can be diverse (e.g.
73 water, carbon and nitrogen) and influence ecosystem responses to future
74 precipitation, since depending on the size of the accumulated reservoir,
75 environmental response will be different (Austin *et al.* 2004; Collins *et al.* 2014). For
76 example, the duration of a dry period before a rain event determines the size of the
77 accumulated reservoir of organic matter and N inorganic. Consequently, this can

78 shape the size of denitrification activity generated by rainfall (Austin *et al.* 2004;
79 Reynolds *et al.* 2004). At some level, there is an environmental 'memory' of previous
80 precipitation events, which is one of the keys to understanding the dryland
81 environmental sensitivity, especially in intra-seasonal patterns (Reynolds *et al.* 2004;
82 Schwinning *et al.* 2004).

83 The phyllosphere (aerial part of plants) receives the effects of the external
84 environment and mediates its relationship with the plant (Vacher *et al.* 2016; Koskella
85 2020). Furthermore, this habitat is home to one of the most diverse microbiotas on
86 the planet, which includes groups such as bacteria, archaea, viruses, filamentous
87 fungi, yeasts, microalgae, protozoa and nematodes (Vorholt 2012; Goffredi, Jang and
88 Haroon 2015; Vacher *et al.* 2016; Thapa and Prasanna 2018; Koskella 2020). One of
89 the most challenging and common conditions that phyllosphere microorganisms face
90 is water-deficit stress. Even in more humid environments, the cuticle (the layer that
91 covers the plant) is a hydrophobic surface with low wettability. It is difficult for water to
92 remain in the epiphytic regions (especially in the leaves), generating an environment
93 with little water available and high osmotic pressure due to accumulation of
94 metabolites (Beattie and Lindow 1995; Lindow and Brandl 2003; Vorholt 2012;
95 Bringel and Couée 2015; Vacher *et al.* 2016).

96 Microorganisms of the phyllosphere have essential functions for plants, acting
97 in processes of defense, pathogenicity, nutrient acquisition, growth, reproductive
98 success and evolution (Rodriguez *et al.* 2009; Thapa and Prasanna 2018; Leveau
99 2019; Koskella 2020). For example, phyllosphere yeasts can act in nutrient-cycling
100 processes, antagonize phytopathogens and produce plant hormones that stimulate
101 growth (Buck 2002; Limtong and Koowadjanakul 2012). Because of the oligotrophic
102 nature of the phyllosphere, this environment tends to select microorganisms capable
103 of 'remodeling' it to access resources. For example, narrowing the cuticle, producing
104 hormones, enzymes, exopolysaccharides (EPS) and biosurfactants (Beattie and
105 Lindow 1995; Leveau 2019). Some of these strategies can increase and/or facilitate
106 access to water and drought tolerance, such as the production of biosurfactants that
107 increase leaf wettability (Beattie and Lindow 1995; Leveau 2019; Koskella 2020).
108 Events of intense taxonomic variation are recorded in the phyllosphere microbiome,
109 but recent evidence indicates the existence of a functional core (Louca *et al.* 2017;

110 Lajoie, Maglione and Kembel 2020). Therefore, plants may have active mechanisms
111 to select essential functional groups, often related to the metabolism of amino acids,
112 nucleotides, carbohydrates, and energy (Lajoie, Maglione and Kembel 2020).
113 Morevoer, as the prevailing condition in drylands is water deficit, fungi, rather than
114 bacteria, can dominate the processes of N decomposition and transformation, in
115 addition to generating symbiotic associations with primary producers (Austin *et al.*
116 2004; Collins *et al.* 2014).

117 This study makes relevant contributions to the knowledge of microbial diversity
118 in drylands, as well as some functional traits that may represent ecological and
119 biotechnological insights, such as the production of enzymes and biosurfactants.
120 Additionally, we address the following question: How do the dynamics of rainfall
121 pulses and drought (rainfall volume, storm or single-rain events, seasonality and
122 memory) affect the yeast community structure and diversity of the bromeliad
123 phyllosphere from Brazilian semiarid dryland? Based on this question, the following
124 hypotheses were tested: Hⁱ- Heavier and more continuous rainfall, and longer periods
125 of drought preceding the rain event (memory effect) increase yeast diversity
126 (taxonomic, phylogenetic and functional) and abundance. In view of this, the diversity
127 and abundance of yeasts were greater during the rainy season. Hⁱⁱ- The rainfall,
128 drought duration, memory effect and seasonality modulate the structure of
129 taxonomic, phylogenetic and functional of the phyllosphere yeast community. Hⁱⁱⁱ-
130 Furthermore, the seasonal periods present a distinct taxonomic and phylogenetic
131 structure but are functionally similar.

132 Methods

133 Model plant

134 The plant model species was *Bromelia laciniosa* Mart. ex Schult. & Schult.f
135 (Bromeliaceae, Bromelioideae), an endemic species of the semiarid dryland located
136 in Northeast Brazil (Caatinga) (Ferreira, Fabricante and Siqueira-Filho 2015). Its
137 leaves, flowers and fruits are popularly used in the treating child colic, diarrhea, fever,
138 jaundice, dandruff and hepatitis (de Albuquerque *et al.* 2007). It is a typically
139 terrestrial plant insert in the terrestrial CAM (crassulacean acid metabolism)
140 functional group (Males and Griffiths 2017).

141 *Study area description*

142 The collections were carried out in the Tocaia Reserve (RPPN Tocaia),
143 municipality of Santana do Ipanema, Alagoas-Brazil, located in the semi-arid region
144 of northeastern Brazil (Caatinga) ($9^{\circ}23'08.9"S$ and $37^{\circ}15'22.8"W$) along a transect of
145 5×200 meters. The rainfall occurs irregularly, with an annual volume of 500 - 700
146 mm. The RPPN Tocaia has an area of approximately 20 hectares and a maximum
147 altitude of about 400 meters, it is mainly composed of an arboreal physiognomy. The
148 predominant climate is semi-arid type Bsh (Köppen).

149 *Collections of plant material*

150 Twelve collections were carried out, six in the dry season (October, November
151 and December 2020 and 2021) and six in the rainy season (May, June and July 2021
152 and 2022). In each collection, five individuals of adult and apparently healthy
153 bromeliads were randomly sampled along the previously established transect. Three
154 leaves were randomly selected from each plant individual collected, making a total of
155 60 individuals, and 180 leaves collected during the collections. The leaves were
156 sealed in sterile plastic bags, then transported at room temperature ($27 \pm 2^{\circ}\text{C}$) for
157 approximately 4 h to the laboratory. In the laboratory, the leaves were stored at $10-15$
158 $^{\circ}\text{C}$ and processing was carried out within a maximum of 24 h after the material was
159 collected in the field.

160 *Samples processing*

161 The leaves underwent an initial wash with sterile distilled water to remove
162 contaminants such as dust and other exogenous materials. Then, leaf fragments
163 between 5 and 20 cm^2 were randomly cut contemplating the 3 leaves collected, until
164 a total area of 120 cm^2 was obtained, and the mass was measured, which was
165 allocated in a 250 mL Erlenmeyer flask containing 75 mL of sterile distilled water.
166 The water containing the leaf fragments was stirred for 30 min at 180 RPM. After the
167 shaking, the washing product was collected in a sterile flask and the leaf fragments
168 were preserved in an Erlenmeyer flask. After removing the water from the first wash,
169 75 mL of a 0.5% solution of Tween 80 was added and stirred again under the same
170 time and rotation. The product of the second wash was then added to that of the first,

171 forming a pool. A 100 µL fraction of this product was seeded in petri dishes
172 containing modified YM Agar medium (0.3% yeast extract, 0.3% malt extract, 0.5%
173 bacteriological peptone, 1% glucose, 2% agar, 0.04% chloramphenicol, pH 4.0) in
174 the dilution of 10⁰ and 10⁻¹, in duplicate. The plates were incubated at 25-28 °C and
175 observed daily during 10 days for yeast isolation.

176 ***Isolation and storage of yeasts***

177 The different yeast morphotypes, delimited considering colonial characteristics
178 such as color, margin, texture, shape, and elevation, were isolated on YEPD agar
179 medium (2% glucose, 1% bacteriological peptone, 0.5% yeast extract, and 2% agar).
180 All yeast isolates obtained during 10 days of isolation were preserved in GYMP
181 medium (2% glucose, 2% malt extract, 0.5% yeast extract, 0.2% monobasic sodium
182 phosphate, and 2% agar) for further analysis.

183 ***Functional traits***

184 The assimilation profile was estimated during 21 days of growth based on the
185 ability of the isolates to grow using a single carbon source following Kurtzman *et al.*
186 (2011). The carbon sources used were cellobiose, D-arabinose, galactose, glucose,
187 glycerol, inulin, raffinose, rhamnose, and xylose. In addition, the ability of the isolates
188 to ferment glucose was verified to produce extracellular hydrolytic enzymes such as
189 amylase, caseinase, cellulase, esterase, lipase, and pectinase, and to produce
190 biosurfactants.

191 The diameter of the halos and colonies of each isolate were measured to
192 evaluate the production of extracellular enzymes, and the modified pz equation was
193 used: $pz = \left| \left(\frac{Dc}{(Dc+Dh)} \right) - 1 \right|$. Where pz represents the enzymatic activity, Dh and Dc
194 are, respectively, the diameter of the hydrolysis halo and the diameter of the colony.

195 The inoculum standardization for the evaluation of enzyme production was
196 performed using turbidity as a reference to quantify the number of cells. Cells of the
197 isolates were diluted in 2 mL of sterile distilled water until reaching an approximate
198 concentration of 10⁵ cells/mL, equivalent to degree 1 of the Wickerham card. Then

199 the inoculum was applied as dot on the culture medium using the replicator stamp.
200 The Petri dishes were analyzed after 7-10 days of incubation at 25-28 °C.

201 The amylolytic activity was evaluated using a modified starch agar culture
202 medium (0.5% soluble starch, 0.5% bacteriological peptone, 0.5% yeast extract,
203 0.05% magnesium sulfate, 0.001% iron sulfate, 0.001% sodium chloride, 1.5% agar).
204 After the incubation period, a 1% Lugol solution was applied to the plate to visualize
205 the hydrolysis halo. The isolate with amylolytic activity shows a halo not stained by
206 Lugol and remains clear (Buzzini and Martini 2002; Mautone *et al.* 2010; Carrasco *et*
207 *al.* 2012).

208 The cellulolytic activity was evaluated by modified CMC agar (0.5%
209 carboxymethylcellulose (CMC), 0.1% sodium nitrate, 0.1% monobasic potassium
210 phosphate, 0.1% potassium chloride, 0.05% magnesium sulfate, 0.05% yeast
211 extract, 0.1% glucose and 1.7% agar). The result was visualized with a 0.1% solution
212 of Congo red added to the plates, and after 40 min the plates were washed with 1M
213 NaCl solution. Isolates capable of hydrolyzing CMC showed an orange halo (Buzzini
214 and Martini 2002; Carrasco *et al.* 2012).

215 The esterase activity was evaluated using modified a Tween 80 agar medium
216 (2.5% tween 80, 0.1% glucose, 1% bacteriological peptone, 0.5% sodium chloride,
217 0.1% calcium chloride, and 2% agar). After the incubation period, the isolates that
218 hydrolyzed the substrate showed a halo formed by a whitish precipitate halo (Buzzini
219 and Martini 2002; Carrasco *et al.* 2012).

220 The protease activity was evaluated using a modified casein agar medium
221 (1% casein, 0.5% glucose, and 2% agar pH 7.0). After the incubation period, the
222 caseinase-producing isolates showed a whitish halo (Buzzini and Martini 2002;
223 Carrasco *et al.* 2012).

224 The pectinase activity was evaluated using a modified pectin agar medium
225 (0.67% Yeast Nitrogen Base (YNB), 1% citrus pectin, 1% glucose, and 1.8% agar,
226 pH 7.0). After the incubation period, 1% hexadecyltrimethylammonium bromide
227 (CTAB) was added to the plate. The producer isolates showed a clear halo around
228 the colony (Buzzini and Martini 2002; Carrasco *et al.* 2012).

229 The surfactant/emulsifying activity was evaluated using the emulsion index
230 with stability after 24 h (IE_{24}). For testing, isolates were cultured for 48 hours at 22-25
231 °C in YEPD broth. To verify the activity, after growth in 2 mL of YEPD broth, 2 mL of
232 kerosene was added to the culture, stirred for 2 minutes in the tubes and then left to
233 rest for 24 h. The total height of the crop and the height of the emulsion in
234 centimeters were then measured. The IE_{24} was obtained from the equation $IE_{24} =$
235 $(\frac{\text{Height}_{\text{emulsion}}}{\text{Height}_{\text{total}}}) \times 100$, that is, the proportion of the emulsion height in relation to the
236 total height of the crop, in percentage.

237 ***Molecular identification of yeasts***

238 The total genomic DNA was extracted following the protocol for the small-
239 scale preparation of modified yeast DNA, originally proposed by (Sambrook and
240 Russel 2001). Then, the D1/D2 region of the LSU gene rRNA was amplified by the
241 Polymerase Chain Reaction (PCR) technique using the primers NL-1 and NL-4
242 (Kurtzman and Robnett 1998; Fell *et al.* 2000). The PCR parameters followed
243 (Landell *et al.* 2010). The amplicons were sequenced using the Sanger method and
244 identified by comparing them to the GenBank database using the Basic Local
245 Alignment Search Tool (BLAST) on the International Nucleotide Sequence
246 Collaboration Databases website using the BLASTn algorithm (Altschul *et al.* 1997).

247 ***Statistical analysis***

248 The colonies were counted to estimate the yeast abundance considering each
249 yeast morphotype isolated from the samples, expressed in colony-forming units per
250 square centimeter (CFU/cm²). Taxonomic diversity was estimated using Hill's
251 numbers, which allow assigning weights to abundances based on the q factor: q0
252 (richness/greater weight for rare species), q1 (typical values) and q2 (which assigns
253 greater weight to more abundant species). The Average Taxonomic Distinctness
254 (AvTD) index (Clarke and Warwick 1998) was estimated using the Vegan package
255 (Dixon 2003). The AvTD was calculated using phylogenetic nodes of the genera,
256 family, order, class, subphylum, and phylum, undefined groups were considered
257 polytomies. The complete classification of each species was obtained from the
258 Mycobank website (<https://www.mycobank.org>) and, when necessary, from

259 specialized literature. The functional richness (FRic), functional dispersion (FDis),
260 and functional evenness (FEve) indices were estimated using the yeast abundances
261 in the samples and the proportional frequency of analyzed functional traits (trait
262 frequency/species richness) by the FD package (Laliberté *et al.* 2014). Whittaker's
263 beta-diversity was estimated in the Vegan package (Dixon 2003) using the mean
264 abundances of the species in the seasonal periods. Beta-diversity fractionation to
265 identify the contribution of nestedness and turnover to beta-diversity was
266 implemented using Jaccard index and binary data (presence and absence) for
267 taxonomic, phylogenetic, and functional composition in the betapart package
268 (Baselga and Orme 2012).

269 The explanatory environmental variables used to assess the effect of the
270 rainfall and drought regime in the community were: i) Storm volume in millimeters,
271 this variable refers to the most recent rainfall event in consecutive days before each
272 sampling. According to Reynolds *et al.* (2004), in certain cases greater and/or longer
273 rainfall is necessary for any biological effect to be noticeable. ii) Volume of the most
274 recent rainy day in millimeters, this variable refers to the most recent rain event
275 before each collection, and these are mostly single-day events. iii) The memory
276 effect accounted for from the most recent rain event. This variable considers the
277 number of days of drought that preceded the most recent rainfall event. iv) Rainy
278 days in the month, this variable considers the number of rainy days in the thirty days
279 preceding each sampling. v) Volume of accumulated rainfall in the month (mm). This
280 variable indicates the accumulated rainfall in the thirty days preceding each
281 sampling. The rainfall values were collected by the HidroWeb tool, which integrates
282 the National Water Resources Information System (SNIRH) from the Brazilian
283 national water agency (ANA) database. The code of the consulted hydrological
284 station is 00937032.

285 To check the possible relationship among the explanatory environmental
286 variables on the abundance and alpha-diversity indices (taxonomic, phylogenetic,
287 and functional), we implemented a generalized linear model (GLM) using negative
288 binomial distribution in the MASS package (Ripley *et al.* 2013). For these analyses,
289 the plant individuals were used as an analytical unit. Bromeliad AT33 was excluded
290 from analysis as no yeast isolate was obtained from this individual. To verify the

291 multivariate relationship of explanatory variables with the structure taxonomic and
292 phylogenetic (by composition and abundance data), and functional (by composition
293 and proportional frequency data) of yeast communities, a series of Canonical
294 Analysis of Principal coordinates (CAP) was implemented using the Bray-Curtis
295 distance in the Vegan package (Dixon 2003). To avoid errors and bias, all models
296 were built avoiding inserting autocollinear variables in the same model. The possible
297 autocollinearity of the explanatory variables was verified by the variance inflation
298 factors (VIF).

299 The effect of seasonality on community diversity and structure was tested
300 considering each of the twelve collections as analytical units over time. To verify the
301 ability of the seasonal periods to group and differentiate samples using the
302 taxonomic, phylogenetic or functional traits of the community was employed a
303 Multivariate Permutation Analysis (PERMANOVA). The residuals method was
304 employed under the full model with 999 permutations by a Principal Coordinate
305 Analysis (PCoA) applying Bray-Curtis distance. Additionally, a similarity percentage
306 (SIMPER) analysis using Bray-Curtis distance and 999 permutations was
307 implemented with species abundance data in seasonal periods. The SIMPER
308 analysis indicated that the elements (variables) that most contributed to the
309 dissimilarity between seasonal periods using taxonomic, phylogenetic or functional
310 information. In addition, the analysis also indicated which of these groups showed
311 significantly different abundance (in the case of taxonomic and phylogenetic data)
312 and frequency (in the case of functional traits) between periods. All analyzes were
313 performed in the R software (R Team 2021).


314 Results

315 *Diversity from plant-yeast system in Brazilian dryland*

316 From bromeliad leaves were recorded a richness of 79 species, 43 genera, 27
317 families and 20 orders of yeasts and yeast-like fungi. Species from phylum
318 Basidiomycota totaled 70% (n=56) (50% Agaricomycotina, 32% Pucciniomycotina
319 and 18% Ustilaginomycotina) and from phylum Ascomycota 30% (n=24) (50%
320 Saccharomycotina, 46% Pezizomycotina and 4% Taphrinomycotina) (Supplementary
321 Table 1). The richness in periods (dry and rainy) was identical: 50 species. The

322 proportion between Basidiomycota and Ascomycota species remained stable
323 regardless of the period.

324 The average yeast richness and standard deviation was 3.9 ± 2.1 species
325 per plant, ranging from 0 to 9 yeast species in a single individual. Only one of the
326 sixty plants sampled did not register any yeast. More than half of the species (55%)
327 were singletons (species with a single record) and 75% were singletons or
328 doubletons. The most frequently recorded species in samples were *Aureobasidium*
329 *thailandense* (36%) and *Occultifur brasiliensis* (25%). The yeast abundance average
330 per plant individual was 9.53×10^2 CFU/cm² (i.e. 4.62×10^3 CFU/g) and the mean
331 abundance per yeast isolate was 1.88×10^2 CFU/cm² (i.e. 8.99×10^2 CFU/g). The non-
332 singleton or non-doubleton species with the highest mean abundance were *Candida*
333 *blankii* (4.48×10^2 CFU/cm²) and *Saitozyma ninhbinhensis* (4.12×10^2 CFU/cm²).
334 Additionally, 37% of the isolates found had abundance values lower than 1.0×10^2
335 CFU/cm² and 85% less than 3.0×10^2 CFU/cm². Less than 2% of the isolates showed
336 an abundance greater than 1.0×10^3 CFU/cm². Among the functional traits evaluated,
337 the most frequent was the assimilation raffinose, inulin and glucose, followed by the
338 production of the extracellular hydrolases (esterase, lipase and cellulase) (Figure 1,
339 Supplementary Figure 1). On the other hand, as less frequent, the emulsification and
340 fermentations abilities were observed. The least frequent traits were the production of
341 the extracellular hydrolase pectinase and amylase (Figure 1, Supplementary Figure
342 1).




343 **Figure 1-** Frequency proportional to species richness of functional traits in each bromeliad sample.
344 Values are shown for each seasonal period and combined data (Total). The black bar indicates the
345 standard deviation.

346 ***How do drought, rainfall and memory effect yeast diversity and abundance?***

347 Contrary to the initial hypothesis, the most recent rainfall volume, the number
348 of dry days the community was exposed to, the number of rainy days in a month and
349 the monthly rainfall volume were not significant predictors for any of the analyzed
350 response variables. The memory effect (dry time that preceded the last rain) did not
351 show a significant relationship with most diversity or abundance metrics, except for
352 FRic (Functional richness). However, the relationship between memory and FRic was
353 negative, that is, the longer the period of drought that precedes the rain, the lower the
354 functional richness found in the yeast community ($\alpha=-0.03$, $R^2= 0.08$, $p\text{-value}=0.011$).
355 The volume of the most recent consecutive rain (storm) was positively and
356 significantly related to the q0 component (species richness) ($\alpha=0.008$, $R^2= 0.14$, $p\text{-}$
357 value=0.002) and with FRic ($\alpha=0.03$, $R^2= 0.26$, $p\text{-value}<0.0001$). But it showed no
358 relationship with the other diversity metrics or with the abundance of yeasts.

359 ***Taxonomic, phylogenetic and functional yeast community structure***

360 Among the explanatory variables, the storm volume showed a significant
361 relationship with the taxonomic ($p\text{-value}=0.001$, $R^2=0.05$) and phylogenetic ($p\text{-}$
362 value=0.002, $R^2=0.08$), but not with functional community structure (Figure 2). The
363 memory effect was relevant only in the taxonomic structure ($p\text{-value}=0.03$, $R^2=0.04$).
364 Also, the most recent rainfall volume was not related to the structure of the yeast
365 community in any of its facets. Only the rainy days in a month ($p\text{-value}=0.007$, $R^2=$
366 0.37), monthly rainfall volume ($p\text{-value}=0.005$, $R^2= 0.41$) and drought days duration
367 ($p\text{-value}=0.001$, $R^2= 0.53$) showed a significative relationship only with the functional
368 structure of the community (Figure 2 and Supplementary Figure 1).



369

370
371
372
373
374
375

Figure 2- Non-metric multidimensional scaling (NMDS) using data on species abundance (a). The abundance of phylogenetic groups (genera, family, order, class, subphylum, and phylum) (b). Values of frequency of the functional traits (c). All ordination analysis were constructed by Bray-Curtis distance, and the vectors marked with a red asterisk indicate represent the variables that showed a significant relationship with the structure of the community by Canonical Analysis of Principal coordinates (CAP) result.


376 *The seasonal periods and yeasts*

377 The Hill's components (q_0 , q_1 and q_2) used to estimate taxonomic diversity
 378 did not differ significantly between season periods ($p\text{-value}>0.5$). The same
 379 happened in the estimators of phylogenetic diversity (Average taxonomic distinctness
 380 – AvTD, $p\text{-value}=0.1$) and functional diversity (FRic, FDis, and FEve, $p\text{-value}>0.51$)
 381 (Figure 3). The yeast abundance per plant was 9% lower in the rainy season
 382 (9.29×10^2 CFU/cm²) when compared to the dry season (1.0×10^3 CFU/cm²) and this
 383 difference was not significant ($p=0.23$). No species occurred in all collections carried
 384 out in both seasons. The species *A. thailandense* and *C. blankii* were the only ones
 385 present in all the collections of the rainy period and *Papiliotrema laurentii* and
 386 *Tremella ananatis* were the only ones to be present in all the collections of the dry
 387 period. Additionally, only 25% ($n=20$) of the species were shared between seasons,
 388 and 37.5% ($n=30$) were exclusively recorded in each.

389 Considering species composition and abundance data, the yeast community
390 structure varied between seasonal periods ($p\text{-value}=0.002$, $R^2=0.15$), but not when
391 considering phylogenetic ($p\text{-value}=0.09$) or functional ($p\text{-value}=0.57$) data. This
392 taxonomic difference between periods was also observed in the Whittaker beta-
393 diversity values: taxonomic ($w=0.58$), phylogenetic ($w=0.35$) and functional ($w=0.03$).
394 Considering occurrence (binary) data, 99.5% of the beta-diversity observed in yeast
395 species and phylogenetic groups between seasonal periods was generated by the
396 species replacement component (turnover). However, the functional beta-diversity
397 was negligible, represented fully by the nestedness component.

398 The abundance of no species differed significantly between seasonal periods,
399 however the species that most contributed to the dissimilarity between periods were
400 *O. brasiliensis* (8% contribution, more abundant in the dry period) and *C. blankii* (8%
401 contribution, more abundant in the rainy season) (Supplementary Table 2). Among
402 the phylogenetic groups, the group that diverged significantly was the genus *Candida*
403 and related groups such as: the order Saccharomycetales, the class
404 Saccharomycetes, and the subphylum Saccharomycotina ($p\text{-value}\leq 0.013$)
405 (Supplementary Table 3). The genus *Candida* was the most abundant in the rainy
406 season, which led to an increase in the other related phylogenetic groups.

407 The assimilation of rhamnose was more frequent in the rainy season and was
408 the trait that most contributed to the functional dissimilarity between periods, however
409 it did not differ significantly between dry and rainy seasons. The traits that varied
410 significantly between seasonal periods were galactose assimilation ($p\text{-value}=0.038$)
411 and cellulase production ($p\text{-value}=0.024$), both more frequent in the rainy season
412 (Figure 1, Supplementary Table 4). Even though the majority did not demonstrate
413 significant differences, 76% of the evaluated functional traits were more frequent in
414 the rainy season. Only the production of lipase, pectinase and caseinase enzymes
415 and the ability to emulsify (IE_{24}) were most frequent in the dry period. There was no
416 difference in the number of traits expressed per species between seasonal periods,
417 neither when comparing core species (present in both periods), and when comparing
418 exclusive species ($p\text{-value}\geq 0.5$). Most of the functional traits evaluated were trophic,
419 related to the acquisition and use of nutrients. The results indicate that there is no
420 difference in the polytropy of the species between the seasonal periods.



421

422


423

424

Figure 3- Boxplots with the variances of abundance (a) and alpha-diversity metrics of taxonomic (b, c and d), phylogenetic (e), and functional (f, g and h) between seasonal periods. The asterisk indicates the average.

425 **Beta-diversity partitions**

426 Taxonomic beta-diversity was higher compared to phylogenetic and functional
 427 in all scales analyzed: seasonal, temporal, among collections and plant individuals
 428 (Figure 4). Except for the functional facet, turnover accounted for most of the beta-
 429 diversity observed across all scales. In all different facets, beta-diversity was higher
 430 among bromeliad individuals and the most significant contribution of nestedness to
 431 beta-diversity was observed; however, this participation was always minor compared
 432 to turnover in general. In all observed scales, only functional beta-diversity showed a
 433 more significant contribution from nestedness than turnover. Furthermore, the
 434 contribution of nestedness to beta-diversity was lower on the seasonal scale,
 435 gradually increased on the temporal scale, among collections, and reached the
 436 highest contribution among individuals (Figure 4). This trend was the opposite in
 437 functional beta-diversity and seasonally and temporally the contribution of
 438 nestedness is absolute.



439

440 **Figure 4-** Jaccard beta-diversity values (binary data), at different scales: Seasonal, Temporal
 441 (between the four quarterly collection blocks), Among collections and Among plant individuals. In
 442 addition, we also indicate the beta diversity partitions, mostly turnover is the most representative
 443 partition. However, functional beta-diversity is dominated by nestedness.

444 Discussion**445 Yeasts from Caatinga dryland**

446 Certain groups of microorganisms are more associated with one substrate
447 than with others. In the case of yeasts, the phylum Basidiomycota is prevalent in
448 leaves, intact fruits and flowers surfaces, when nectar is disregarded (Fonseca and
449 Inácio 2006). In addition, it is expected that the community will change over the
450 seasonal and ontogenetic periods. In summer (mainly in temperate environments),
451 the genera *Papiliotrema*, *Rhodotorula*, and *Sporobolomyces* are overrepresented,
452 mainly due to factors such as temperature, sunlight, and day length (Glushakova and
453 Chernov 2004, 2010; Fonseca and Inácio 2006; Vacher *et al.* 2016; Kemler *et al.*
454 2017). On the other hand, Fonseca and Inácio (2006) cite evidence that in the
455 wettest and coldest months, *Cryptococcus* spp. (currently, *Papiliotrema* and others),
456 mainly *Papiliotrema laurentii*, is prevalent. Several species from
457 *Cryptococcus/Papiliotrema*, can produce polysaccharide capsules that confer
458 resistance to desiccation (Fonseca and Inácio 2006; Kemler *et al.* 2017). In addition,
459 traits such as the production of pigments can generate a competitive advantage in an
460 environment exposed to various oxidative stresses, such as the surface of leaves.
461 Pigment producers such as *Rhodotorula* and *Sporobolomyces* (carotenoids), and
462 *Aureobasidium* (melanin) are persistent in the phylloplane (Fonseca and Inácio 2006;
463 Kemler *et al.* 2017).

464 In most studies, the seasonal dynamics of phyllosphere microbiota (including
465 yeasts) are developed in temperate environments, where sudden changes in climatic
466 and physiological conditions mark the seasons. In the tropical environment, the
467 maximum productivity is marked by the rainy season and the minimum by the dry
468 season. In addition, temperatures do not vary as much between seasons. In
469 temperate environments, fungal diversity metrics and abundance tend to vary over
470 seasonal periods due to a dependence on the hydrothermal regime and the
471 ontogenetic period of the host (Glushakova and Chernov 2004, 2010; Fonseca and
472 Inácio 2006). The senescence process favors the appearance of damage to the
473 cutin, making the nutritive components of the leaves more available, decreasing the
474 hydrophobicity of the leaf surface, and potentially decreasing the concentration of

475 antimicrobial substances. This explains the peak of diversity and abundance of the
476 epiphytic microbiota during autumn (Inácio *et al.* 2002; Glushakova and Chernov
477 2004, 2010; Fonseca and Inácio 2006). Some hypotheses claim, that one of the
478 reasons for the increase in the leaf community in autumn may be higher humidity and
479 milder temperatures (Inácio *et al.* 2002; Fonseca and Inácio 2006). In tropical
480 environments, community drivers can vary. For bacteria, the leaf content of the water
481 and phosphorus influences the community's abundance and structure, while for fungi,
482 the leaf aluminum content is significant (Vacher *et al.* 2016). However, it is not easy
483 to extrapolate these associations to all communities in the phyllosphere.

484 Our data indicate an increase in the frequency and abundance of the groups
485 such as Basidiomycota in the dry period, especially the species *O. brasiliensis*, *P.*
486 *laurentii* and *T. ananatis*. On the other hand, the groups that stand out in the rainy
487 season are the phylum Ascomycota, the genus *Candida* and species such as *A.*
488 *thailandense* and *C. blankii*. Other studies have observed that in plants, mainly in
489 leaves: the proportion of Ascomycota reaches its maximum in autumn and its
490 minimum between late winter and early spring (Glushakova and Chernov 2010; Abu-
491 Ghosh, Droby and Korine 2014). In the phyllosphere of mosses of the genus
492 *Sphagnum* in swamp (wetter) ascomycetes yeasts were more abundant than in
493 forest (lower humidity), 18.4 and 10.95% of the abundance, respectively (Kachalkin
494 and Yurkov 2012).

495 In a dryland system, an increase in the proportion of basidiomycetes yeasts in
496 relation to ascomycetes was observed in drier periods (Abu-Ghosh, Droby and
497 Korine 2014). In summer, basidiomycetes reached 86% of the species of the
498 phyllosphere. As the ascomycete species increased in wetter periods, the frequency
499 of phyla was the same during this period (Abu-Ghosh, Droby and Korine 2014).
500 However, considering the whole mycobiome of the soil, greater aridity (greater water
501 deficit) promotes dominance of fungi of the phylum Ascomycota (Tedersoo *et al.*
502 2014; Maestre *et al.* 2016). Our data did not show differences in the proportion of
503 ascomycetes in the rainy or dry season, but that some species and groups of
504 ascomycetes are more frequent and/or abundant in the rainy period. It is possible
505 that the proportion increase between phyllosphere ascomycetes and basidiomycetes
506 in more humid seasonal periods observed in several works is characteristic of

507 temperate environments. In the tropical environment, the yeast phylum proportion
508 tends to be similar between seasonal periods, as well as taxonomic diversity. This
509 statement is supported by the study by Gomes *et al.* (2015), who found similar
510 diversity of yeasts in bromeliad tanks. However, the composition of the community
511 was quite different between periods.

512 ***Does rain and drought affect yeasts on leaves?***

513 In drylands, water availability is considered the central factor in controlling
514 biological activity. Furthermore, the leaf surface is a hydrophobic region subject to
515 intense water limitation. Some factors, such as host age and rainfall regime, can
516 change water availability in the phyllosphere (Vacher *et al.* 2016). Plant aging tends
517 to change physicochemical characteristics in the cuticle, which becomes more
518 wettable and permeable, increasing water adhesion and nutrient availability (Vacher
519 *et al.* 2016; Oso *et al.* 2021). Very waxy leaves have poor wettability; even if water is
520 available through humidity or rain, a film of water will hardly form. In addition to
521 promoting water availability for the microbial community on the leaf, the formation of
522 this water film quickly changes the pH and redistributes nutrients in the epiphytic
523 region (Morris 2001). It is possible that the frequency and intensity of rainfall are also
524 related to the microbial response (Kinkel 1997). Rain fall is known to alter
525 colonization and dispersion rates, washing the leaves and removing microorganisms,
526 while transferring other microorganisms from the atmosphere to the plant (Morris
527 2001; Vacher *et al.* 2016; Leveau 2019).

528 In this way, we expected that the rains would contribute strongly to the
529 phyllosphere yeast community diversity. However, differing from the initial
530 hypothesis, rain did not present significant influence. Only storm volume and memory
531 effect had any significant relationship with certain diversity metrics. Storm volume
532 positively influenced functional and species richness, and memory effect was related
533 only to functional richness. Although, in the case of memory effect, the relationship
534 was negative, diverging from our hypothesis.

535 Studies on epiphytic yeasts in the desert suggest that these microorganisms
536 are resistant to various abiotic stresses, mainly of oxidative nature (Abu-Ghosh,
537 Droby and Korine 2014). However, in experimental studies with grasses,

intermediate drought conditions did not significantly affect the phyllosphere bacterial community, but when drought was intensified, diversity tended to decrease (Bechtold *et al.* 2021). It is plausible that longer droughts than the one we have recorded (a maximum of 58 days without any rain) could affect the epiphytic microbiota. In the phyllosphere of many plants, short-term rainfall events tend not to result in changes in microbiota diversity and abundance. For example, in tomato leaves (Allard, Ottesen and Micallef 2020) and macrophyte leaves (Stone and Jackson 2021) rainfall did not influence the richness and abundance of the bacteria communities in the short term. In contrast, in tomato fruits rainfall increased the richness few days later until reaching close to the richness recorded before the rain (Allard, Ottesen and Micallef 2020). There is evidence that rainfall and other long-term factors, such as seasonality, influence the microbial community of the phyllosphere more than short-term factors (Chen *et al.* 2021; Stone and Jackson 2021; Yan *et al.* 2022).

551 *Taxonomic, phylogenetic and functional structures of the community*

Rainfall metrics did not influence, or not as expected, facets of yeast diversity while environmental variables proved to be relevant for the organization of communities. Interestingly, the volume of the most recent storm was significantly related to the taxonomic and phylogenetic structure of the community. While the most recent rainfall, mostly a single-rain event, showed no significant relationship with the data. This may be an indication that in this environment, a semiarid dryland, not only the pulse dynamics, but also the constancy of the rain is relevant to the relationships and structures of the community. Reynolds *et al.* (2004) argues about minimum thresholds necessary for rain events to be reflected in biological effects. In drylands, rainfall events generally do not exceed 2 mm, in more arid drylands it is estimated between 10-50 rainy days per year, distributed between 3-15 rain events, of which 5 or 6 have the magnitude to generate effects biologically significant (Noy-Meir 1973; Collins *et al.* 2014).

According to Kinkel (1997), the microbial community of phyllosphere is regulated by four population processes: i) immigration, ii) emigration, iii) growth (generation) and iv) death. For Vellend (2010), the mechanisms that regulate ecological communities are diverse, but all of them can be synthesized in just four

types of processes: selection, drift, speciation and dispersion. In this context, selection reflects differences in fitness, drift refers to stochastic changes in the relative abundances of species in a community, speciation refers to the emergence of new species, and dispersion deals with the movement of organisms in space (Vellend 2010; Vacher *et al.* 2016). The memory effect was significant for the taxonomic structure, reinforcing that these variables are more relevant for the structure of the community than for diversity. The functional structure was unrelated to variables such as most recent rainfall, memory, or storm volume. On the other hand, it was strongly related to drought days, monthly volume of rainfall and monthly rainy days.

Other studies, mostly in temperate environments, indicate a gradual increase in yeast diversity in leaves from seasonal periods with a peak of diversity in autumn (Glushakova and Chernov 2004, 2007, 2010). One mechanism that justifies this is the milder temperatures, lower humidity, and more nutrients made available by plants (Fonseca and Inácio 2006; Gouka, Raaijmakers and Cordovez 2022). However, this seasonal cycle tends to occur mainly in short-lived plants and in temperate environments. In our study, the model plants were bromeliads, a tropical group of slow-growing and long-lived plants that do not lose leaves seasonally (Benzing 2000; Ladino *et al.* 2019). We expected that water availability would be translated into greater nutrient availability and diversity during the rainy season, therefore increasing alpha-diversity in all its facets: taxonomic, phylogenetic and functional. However, our hypothesis was falsified, the diversity and abundance of yeasts did not vary significantly between dry and rainy seasons. Gomes *et al.* (2015), found the same pattern when analyzing yeast diversity in bromeliad tanks in Brazil, the yeast community showed stable diversity values and varied compositionally.

Stability in taxonomic alpha-diversity may explain the stability in other diversity facets like functional. The relationship between biodiversity and ecosystem functioning is robustly supported by evidence from observational and experimental studies, therefore, more diverse biological communities are expected to present greater diversity of ecosystem functions (Cardinale *et al.* 2012). The links that mediate this relationship are diverse and not fully understood. Some of these mechanisms are the complementary niche partitioning (when several species

601 complement each other spatio-temporally in resource use), the effect of species
602 identity (when specific species have a disproportionate functional role being crucial to
603 ecosystem functioning) and facilitation (when species facilitate the activities of others
604 by improving ecosystem functioning) (Daam *et al.* 2019). The functional diversity of
605 yeasts was similar between seasonal periods, and this may be a causal relationship
606 with the also “stable” taxonomic diversity. However, we cannot determine whether
607 these factors are just correlated.

608 **Beta-diversity**

609 Between seasonal periods, our data point to a large and significant taxonomic
610 beta-diversity of yeasts, a modest and non-significant phylogenetic variation and a
611 strong functional stability. In *Typha latifolia* macrophyte leaves, a high seasonally
612 bacteria beta-diversity was found, but related to nestedness and not turnover (Stone
613 and Jackson 2021). Host species can influence microbial composition and structure
614 in the phyllosphere (Bechtold *et al.* 2021). In another example, the organization of
615 endophytic and epiphytic bacterial communities differs. The epiphytic community
616 appears to be more widely distributed among distinct species, and the endophytic
617 community tends to be particular to each host species (Yao *et al.* 2020).

618 Similar to our results, endophytic fungi from the phyllosphere and rhizosphere
619 of cacti from Caatinga dryland showed a trend towards stability in species richness,
620 but with intense turnover (Ferreira-Silva *et al.* 2021). Even after being subjected to
621 rain events epiphytic bacteria in tomato leaves showed stability in OTUs richness and
622 temporal variation in community composition (Allard, Ottesen and Micallef 2020). The
623 associated bacterial community can exhibit significant taxonomic variation in
624 bromeliads, even at a few meters of spatial scales (Louca *et al.* 2017). This pattern
625 was also observed in yeasts in bromeliad phytotelmata between seasonal periods, with
626 similar diversity values (Gomes *et al.* 2015). Recent studies suggest that, even using
627 culture-independent approaches, communities can vary substantially by sharing few
628 OTUs between different plants (Yan *et al.* 2022). Multiple variables may influence the
629 discrepancies among studies, both the type of approach (culture- dependent or
630 independent), and factors such as the microbial group studied and the scale of the
631 study.

632 Several studies point to the phyllosphere as a dynamic environment with an
633 equally dynamic and diverse community with significant taxonomic variation in both
634 bacteria and fungi (Andrews *et al.* 1987; Kinkel 1997; Vacher *et al.* 2016; Louca *et al.*
635 2017; Stone and Jackson 2021). We observed a high taxonomic and phylogenetic
636 beta-diversity (almost entirely generated by turnover) in all observed scales:
637 seasonal, temporal, among collections and among plant individuals. However, the
638 functional beta-diversity was low and represented by nestedness. The turnover of
639 epiphytic species is well documented and may have several causes: seasonality,
640 senescence, host type, and temporal and spatial variation (Inácio *et al.* 2002;
641 Glushakova and Chernov 2004, 2010; Fonseca and Inácio 2006; Peñuelas *et al.*
642 2012). This is not the first study to record a large taxonomic variation associated with
643 the stability of functional groups (or traits). Louca *et al.* (2017) verified a large spatial
644 turnover of bacterial species in bromeliad tanks while maintaining functional groups.
645 Lajoie *et al.* (2020) reported that the most abundant (and most consistent) functional
646 groups in the phyllosphere microbiota of several tree species are related to
647 metabolism (about 45%), the main ones being the metabolism of amino acids,
648 nucleotides, carbohydrates, and energy. This functional stability may indicate the
649 existence of a phyllosphere functional core (Lajoie, Maglione and Kembel 2020).
650 There is a discussion about taxonomic and functional decoupling, where the
651 taxonomic structure can vary significantly while keeping its functional configuration
652 stable. Nevertheless, the discussion revolves mainly around prokaryotes, with few
653 mentions of eukaryotic organisms (Louca *et al.* 2018).

654 Stochastic and deterministic mechanisms are important for assembling
655 biological communities, including the phyllosphere microbiota (Vellend 2010; Vacher
656 *et al.* 2016; Gouka, Raaijmakers and Cordovez 2022). The high turnover of species
657 and phylogenetic groups indicates high community dynamics at different scales and
658 a high rate of extinction and colonization. Colonization has stochastic facets and
659 occurs mainly through vectors such as rain, wind and animals (mostly insects and
660 birds. On the other hand, selection involves deterministic processes related to
661 hosting species and the fitness of colonizing species or isolates (Andrews *et al.* 1987;
662 Kinkel 1997; Vellend 2010; Mittelbach *et al.* 2015; Vacher *et al.* 2016; Blackwell
663 2017). The results indicate a stochastic and intense colonization dynamic; however,

664 non-random selection processes are based on functional traits. It is reflected in the
665 high taxonomic turnover and high functional stability observed.

666 ***Functional traits from phyllosphere yeasts***

667 As mentioned, there were no significant differences in relation to the
668 components of functional diversity between periods. The more expressed traits in the
669 phyllosphere were the carbon source assimilation, principally plant-derived oligo and
670 monosaccharides like raffinose, glucose and xylose, but also polysaccharides like
671 inulin. In yeasts isolated from flag leaves of wheat carbohydrates (xylose, D-mannitol
672 e N-acetyl-D-glucosamine) were the most frequent nutrients assimilated, followed by
673 polymers and organic acids (Gouka *et al.* 2022). One of the characteristics that
674 allows yeasts to conquer different habitats, including the phyllosphere, is their ability
675 to use a wide spectrum of carbon and nitrogen sources (polytropy). In the study by
676 Hagler *et al.* (1993), more than half of the yeast community recovered from the
677 phytotelmata of bromeliads were able to assimilate more than 20 different carbon
678 sources. The main source of organic nutrients for microorganisms in the phyllosphere
679 (especially leaves) are exudates provided by plants, while inorganic nutrients are
680 generally obtained from the external environment, for example, by wind deposition
681 (Fonseca and Inácio 2006; Kemler *et al.* 2017). In this way, plant physiology plays a
682 vital role in nutritional availability and, consequently, in the structure of the microbial
683 community (Vacher *et al.* 2016). In dry environments, the dynamics of rain pulses
684 generate biological responses in plants. Thus, it is expected that in rainy seasons the
685 availability and diversity of organic compounds for microbiota becomes greater.

686 Yeasts are heterotrophic organisms and extracellular enzymes play a
687 fundamental role in their nutrition. They hydrolyze macromolecules and make
688 nutrients available for the cell to absorb, thus directly or indirectly mediating
689 decomposition (Fonseca and Inácio 2006). We observed that esterase was the most
690 frequently produced among the evaluated hydrolases. Esterases and other
691 hydrolases such as cellulase produced by fungi, are important in the flow of carbon in
692 environments (Treseder *et al.* 2018). On the other hand, the production of cuticle-
693 degrading enzymes and biosurfactants has been proposed as a mechanism that can
694 increase permeability in leaves and facilitate the movement of bacteria in the

695 phylloplane (Lindow and Brandl 2003; Doan and Leveau 2015; Leveau 2019; Oso *et*
696 *al.* 2021). Furthermore, *in vitro*, esterases produced by the epiphytic yeast species
697 *Pseudozyma antarctica* was able to affect the plant's cutin and influence its water
698 dynamics (Ueda *et al.* 2015). Furthermore, esterase produced by *P. antarctica*
699 aggravated the infection caused by *Botrytis cinerea* in tomato plants (Ueda *et al.*
700 2018). It does not indicate that the yeast is pathogenic but may be a facilitator of the
701 infection.

702 In our research, few traits were individually expressed with significative
703 difference when compared dry and rainy seasons. However, a clear pattern is that
704 most of the traits analyzed were more frequently expressed during the rainy season.
705 That corroborates with the idea that rains generate greater nutritional diversity for
706 microorganisms in the phyllosphere. Interestingly, among the few traits that were
707 most frequently expressed in the dry period is the emulsion index (IE₂₄). This index is
708 indicative of one of the activities of a surfactant. Biosurfactants are chemically
709 diverse molecules produced by various microbial groups whose main characteristic is
710 amphipathic; they have a polar and a non-polar region. These molecules have
711 numerous activities such as: decreasing surface tension, increasing wettability on
712 hydrophobic surfaces, emulsifying between immiscible liquids, and solubilizing
713 hydrocarbons (Beattie and Lindow 1995; Thapa and Prasanna 2018; Zeisler-Diehl,
714 Barthlott and Schreiber 2020; Oso *et al.* 2021). Thus, the production of biosurfactants
715 can be an important tool for epiphytic microorganisms to change their habitat and
716 increase the availability of water and nutrients in the phyllosphere (Lindow and
717 Brandl 2003; Leveau 2019). For example, the phytopathogen *Pseudomonas syringae*
718 can increase the leaf's wettability and local water availability by releasing
719 biosurfactants (Koskella 2020). The higher frequency of emulsion is evidence that in
720 the studied dryland (Caatinga), during the drier period, phyllosphere yeasts increase
721 the production of biosurfactants to improve surface wettability and increase access to
722 water. However, more data and studies are needed to evaluate this hypothesis.

723 Conclusions

724 Given this, our results bring unprecedented insights into the dynamics of
725 phyllosphere yeast microbiota from tropical dryland and how it relates to rainfall

726 pulses, drought, seasonality and memory effect. Furthermore, our study added
727 information about the persistent alpha-diversity of the phyllosphere yeast community
728 to seasonal changes, from a taxonomic, phylogenetic and functional perspective.
729 Contrary to what is usually observed in seasonal studies in temperate regions, the
730 yeast community showed stability in diversity and seasonal variation in terms of
731 composition. Indicating a possible tropical seasonal pattern that may differ from the
732 temperate pattern, where there is usually a change in diversity depending on the
733 season. By understanding the effect of rain and drought on the phyllosphere
734 microbiota, especially in drylands, we can think about how the microbial community
735 of the leaves will be affected by the changes in the dry-wet cycles that can be caused
736 by climate change. The phyllosphere microbiota is an important factor for the
737 resilience and maintenance of plant health. Consequently, this community's dysbiosis
738 can influence the host's health. In addition, our results indicate trends in the
739 production of products of biotechnological interest such as enzymes and emulsifiers
740 in different seasons. This information can serve as a guide for bioprospecting in
741 future studies.

742 Acknowledgements

743 We thank Mr. Alberto Nepomuceno Agra Jr. for authorization and assistance during
744 the collections. PhD Patrícia Valente, PhD Jadson Bezerra and PhD Gilberto Justino
745 for the contribution in this work. We also thank MSc. Elizabeth Teixeira and MSc.
746 Hector M. C. Navarro for collaborating in revising the text.

747 Funding

748 This work was supported by Conselho Nacional de Desenvolvimento Científico e
749 Tecnológico (CNPq, number of grants 475378/2013-0, 312889/2021-6 and
750 311553/2018-4), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior
751 (CAPES) and Fundação de amparo a Pesquisa de Alagoas (FAPEAL). This work is
752 part of the project "INCT Yeasts: Biodiversity, preservation and biotechnological
753 innovation", funded by Conselho Nacional de Desenvolvimento Científico e
754 Tecnológico (CNPq), grant #406564/2022-1.

755

756 Conflicts of interest

757 The authors declare that they have no known competing financial interests or
758 personal relationships that could have appeared to influence the work reported in this
759 paper.

760 References

761 Abu-Ghosh S, Droby S, Korine C. Seasonal and plant-dependent variations in
762 diversity, abundance and stress tolerance of epiphytic yeasts in desert habitats.
763 *Environ Microbiol Rep* 2014;6:373–82.

764 de Albuquerque UP, de Medeiros PM, de Almeida ALS et al. Medicinal plants of the
765 caatinga (semi-arid) vegetation of NE Brazil: A quantitative approach. *J Ethnopharmacol* 2007;114:325–54.

767 Allard SM, Ottesen AR, Micallef SA. Rain induces temporary shifts in epiphytic
768 bacterial communities of cucumber and tomato fruit. *Sci Rep* 2020;10:1765.

769 Altschul SF, Madden TL, Schäffer AA et al. Gapped BLAST and PSI-BLAST: A new
770 generation of protein database search programs. *Nucleic Acids Res* 1997;25:3389–
771 402.

772 Andrews JH, Kinkel LL, Berbee FM et al. Fungi, Leaves, and the Theory of Island
773 Biogeography. *Microb Ecol* 1987;14:277–90.

774 Austin AT, Yahdjian L, Stark JM et al. Water pulses and biogeochemical cycles in
775 arid and semiarid ecosystems. *Oecologia* 2004;141:221–35.

776 Baselga A, Orme CDL. betapart: an R package for the study of beta diversity.
777 *Methods Ecol Evol* 2012;3:808–12.

778 Beattie GA, Lindow SE. The Secret Life of Foliar Bacterial Pathogens on Leaves.
779 *Annu Rev Phytopathol* 1995;33:145–72.

780 Bechtold EK, Ryan S, Moughan SE et al. Phyllosphere Community Assembly and
781 Response to Drought Stress on Common Tropical and Temperate Forage Grasses.
782 *Appl Environ Microbiol* 2021;87:1–19.

783 Benzing DH. *Bromeliaceae: Profile of an Adaptive Radiation*. New York: Cambridge
784 University Press, 2000.

785 Blackwell M. Yeasts in Insects and Other Invertebrates. In: Buzzini P, Lachance M-A,
786 Yurkov A (eds.). *Yeasts in Natural Ecosystems: Diversity*. Cham: Springer
787 International Publishing, 2017, 397–433.

788 Bringel F, Couée I. Pivotal roles of phyllosphere microorganisms at the interface
789 between plant functioning and atmospheric trace gas dynamics. *Front Microbiol*
790 2015;6:1–14.

- 791** Buck JW. In vitro antagonism of *Botrytis cinerea* by phylloplane yeasts. *Canadian
792 Journal of Botany* 2002;80:885–891.
- 793** Buzzini P, Martini A. Extracellular enzymatic activity profiles in yeast and yeast-like
794 isolates isolated from tropical environments. *J Appl Microbiol* 2002;93:1020–1025.
- 795** Cardinale BJ, Duffy JE, Gonzalez A et al. Biodiversity loss and its impact on
796 humanity. *Nature* 2012;486:59–67.
- 797** Carrasco M, Rozas JM, Barahona S et al. Diversity and extracellular enzymatic
798 activities of yeasts isolated from King George Island, the sub-Antarctic region. *BMC
799 Microbiol* 2012;12:251.
- 800** Chen QL, Hu HW, Yan ZZ et al. Precipitation increases the abundance of fungal
801 plant pathogens in Eucalyptus phyllosphere. *Environ Microbiol* 2021;23:7688–700.
- 802** Clarke KR, Warwick RM. A taxonomic distinctness index and its statistical properties.
803 *Journal of Applied Ecology* 1998;35:523–31.
- 804** Collins SL, Belnap J, Grimm NB et al. A multiscale, hierarchical model of pulse
805 dynamics in arid-land ecosystems. *Annu Rev Ecol Evol Syst* 2014;45:397–419.
- 806** Daam MA, Teixeira H, Lillebø AI et al. Establishing causal links between aquatic
807 biodiversity and ecosystem functioning: Status and research needs. *Science of the
808 Total Environment* 2019;656:1145–56.
- 809** Dixon P. VEGAN, a package of R functions for community ecology. *Journal of
810 Vegetation Science* 2003;14:927–30.
- 811** Doan HK, Leveau JHJ. Artificial Surfaces in Phyllosphere Microbiology.
812 *Phytopathology* 2015;105:1036–42.
- 813** Fell JW, Boekhout T, Fonseca A et al. Biodiversity and systematic of
814 basidiomycetous yeast as determined by large submit rDNA D1/D2 domain
815 sequence analysis. *Int J Syst Evol Microbiol* 2000;50:1351–71.
- 816** Ferreira JV, Fabricante JR, Siqueira-Filho JA. Checklist preliminar de Bromeliaceae
817 do Parque Nacional do Catimbaú, Pernambuco, Brasil. *Nat On Line* 2015;13:92–7.
- 818** Ferreira-Silva A, Hughes FM, Rosa CA et al. Higher turnover of endophytic fungal
819 assemblages in the tissues of globose cactus *Melocactus ernestii* from Brazilian
820 semi-arid biome. *Symbiosis* 2021;85:79–91.
- 821** Fonseca Á, Inácio J. Phylloplane Yeasts. *Biodiversity and Ecophysiology of Yeasts*.
822 2006, 263–301.
- 823** Frank JH, Lounibos LP. Phytoelmata: swamps or islands? *Florida Entomologist*
824 1987;70:14–20.
- 825** Glushakova AM, Chernov II. Seasonal dynamics in a yeast population on the Oxalis
826 acetosella L. leaves. *Mikrobiology* 2004;73:226–32.

- 827** Glushakova AM, Chernov IY. Seasonal dynamic of the numbers of epiphytic yeasts.
828 *Microbiology (N Y)* 2007;76:590–5.
- 829** Glushakova AM, Chernov IYu. Seasonal dynamics of the structure of epiphytic yeast
830 communities. *Microbiology (N Y)* 2010;79:830–9.
- 831** Goffredi SK, Jang GE, Haroon MF. Transcriptomics in the tropics: Total RNA-based
832 profiling of Costa Rican bromeliad-associated communities. *Comput Struct
833 Biotechnol J* 2015;13:18–23.
- 834** Gomes FCO, Safar SVB, Marques AR et al. The diversity and extracellular enzymatic
835 activities of yeasts isolated from water tanks of *Vriesea minarum*, an endangered
836 bromeliad species in Brazil, and the description of *Occultifur brasiliensis* f.a., sp. nov.
837 *Antonie Van Leeuwenhoek* 2015;107:597–611.
- 838** Gouka L, Raaijmakers JM, Cordovez V. Ecology and functional potential of
839 phyllosphere yeasts. *Trends Plant Sci* 2022;27:1109–23.
- 840** Gouka L, Vogels C, Hansen LH et al. Genetic, Phenotypic and Metabolic Diversity of
841 Yeasts From Wheat Flag Leaves. *Front Plant Sci* 2022;13, DOI:
842 10.3389/fpls.2022.908628.
- 843** Hagler AN, Rosa CA, Morais PB et al. Yeasts and coliform bacteria of water
844 accumulated in bromeliads of mangrove and sand dune ecosystems of southeast
845 Brazil. *Can J Microbiol* 1993;39:973–7.
- 846** Holt RD. Theoretical perspectives on resource pulses. *Ecology* 2008;89:671–81.
- 847** Huang J, Ji M, Xie Y et al. Global semi-arid climate change over last 60 years. *Clim
848 Dyn* 2016;46:1131–50.
- 849** Inácio J, Pereira P, De Carvalho M et al. Estimation and diversity of phylloplane
850 mycobiota on selected plants in a Mediterranean-type ecosystem in Portugal. *Microb
851 Ecol* 2002;44:344–53.
- 852** Kachalkin A V., Yurkov AM. Yeast communities in *Sphagnum phyllosphere* along the
853 temperature-moisture ecocline in the boreal forest-swamp ecosystem and description
854 of *Candida sphagnicola* sp. nov. *Antonie van Leeuwenhoek, International Journal of
855 General and Molecular Microbiology* 2012, DOI: 10.1007/s10482-012-9710-6.
- 856** Kemler M, Witfeld F, Begerow D et al. Phylloplane Yeasts in Temperate Climates. In:
857 Buzzini P, Lachance M-A, Yurkov A (eds.). *Yeasts in Natural Ecosystems: Diversity*.
858 Cham: Springer International Publishing, 2017, 171–97.
- 859** Kinkel LL. Microbial population dynamics on leaves. *Annu Rev Phytopathol*
860 1997;35:327–47.
- 861** Koskella B. The phyllosphere. *Current Biology* 2020;30:R1143–6.

- 862** Kurtzman CP, Fell JW, Boekhout T *et al.* Methods for isolation, phenotypic
863 characterization and maintenance of yeasts. *The Yeasts*. Elsevier B.V., 2011, 87–
864 110.
- 865** Kurtzman CP, Robnett CJ. Identification and phylogeny of ascomycetous yeasts from
866 analysis of nuclear large subunit (26S) ribosomal DNA partial sequences. *Antonie van Leeuwenhoek, International Journal of General and Molecular Microbiology*
867 1998;73:331–71.
- 869** Ladino G, Ospina-Bautista F, Estévez Varón J *et al.* Ecosystem services provided by
870 bromeliad plants: A systematic review. *Ecol Evol* 2019;9:7360–72.
- 871** Lajoie G, Maglione R, Kembel SW. Adaptive matching between phyllosphere
872 bacteria and their tree hosts in a neotropical forest. *Microbiome* 2020;8:1–10.
- 873** Laliberté E, Legendre P, Shipley B *et al.* Package ‘FD.’ *Measuring functional diversity from multiple traits, and other tools for functional ecology* 2014:0–1.
- 875** Landell MF, Billodre R, Ramos JP *et al.* *Candida aechmeae* sp. nov. and *Candida*
- 876** *vrieseae* sp. nov., novel yeast species isolated from the phylloplane of bromeliads in
- 877** Southern Brazil. *Int J Syst Evol Microbiol* 2010;60:244–8.
- 878** Leal IR, Silva JMC, Tabarelli M *et al.* Mudando o curso da conservação da
879 biodiversidade na Caatinga do Nordeste do Brasil. *Megadiversidade* 2005;1:139–46.
- 880** Leroy C, Corbara B, Dézerald O *et al.* What drives detrital decomposition in
881 neotropical tank bromeliads? *Hydrobiologia* 2017;802:85–95.
- 882** Leveau JH. A brief from the leaf: latest research to inform our understanding of the
883 phyllosphere microbiome. *Curr Opin Microbiol* 2019;49:41–9.
- 884** Limtong S, Koowadjanakul N. Yeasts from phylloplane and their capability to produce
885 indole-3-acetic acid. *World J Microbiol Biotechnol* 2012;28:3323–35.
- 886** Lindow SE, Brandl MT. Microbiology of the Phyllosphere. *Appl Environ Microbiol*
887 2003;69:1875–83.
- 888** Louca S, Jacques SMS, Pires APF *et al.* High taxonomic variability despite stable
889 functional structure across microbial communities. *Nat Ecol Evol* 2017;1:0015.
- 890** Louca S, Polz MF, Mazel F *et al.* Function and functional redundancy in microbial
891 systems. *Nat Ecol Evol* 2018;2:936–43.
- 892** Machado IC, Lopes AV. Floral Traits and Pollination Systems in the Caatinga, a
893 Brazilian Tropical Dry Forest. *Ann Bot* 2004;94:365–76.
- 894** Maestre FT, Benito BM, Berdugo M *et al.* Biogeography of global drylands. *New*
895 *Phytologist* 2021;231:540–58.
- 896** Maestre FT, Eldridge DJ, Soliveres S *et al.* Structure and Functioning of Dryland
897 Ecosystems in a Changing World. *Annu Rev Ecol Evol Syst* 2016;47:215–37.

- 898** Males J, Griffiths H. Functional types in the Bromeliaceae: relationships with
899 drought-resistance traits and bioclimatic distributions. Oliveira R (ed.). *Funct Ecol*
900 2017;31:1868–80.
- 901** Mautone JN, Landell MF, Fuentefria AM et al. Phyloplane yeasts as a source of
902 industrially interesting enzymes. *Brazilian Journal of Biosciences* 2010;8:169–73.
- 903** Menezes RSC, Sampaio EVSB, Giongo V et al. Biogeochemical cycling in terrestrial
904 ecosystems of the Caatinga Biome. *Brazilian journal of biology* 2012;72:643–53.
- 905** Mittelbach M, Yurkov AM, Nocentini D et al. Nectar sugars and bird visitation define a
906 floral niche for basidiomycetous yeast on the Canary Islands. *BMC Ecol* 2015;15:2.
- 907** Moro MF, Lughadha EN, Araújo FS de et al. A Phytogeographical Metaanalysis of
908 the Semiarid Caatinga Domain in Brazil. *The Botanical Review* 2016, DOI:
909 10.1007/s12229-016-9164-z.
- 910** Morris CE. Phyllosphere. *Encyclopedia of life sciences* 2001:1–8.
- 911** Noy-Meir I. Desert Ecosystems: Environment and Producers. *Annu Rev Ecol Syst*
912 1973;4:25–51.
- 913** Oso S, Fuchs F, Übermuth C et al. Biosurfactants Produced by Phyllosphere-
914 Colonizing Pseudomonads Impact Diesel Degradation but Not Colonization of
915 Leaves of Gnotobiotic *Arabidopsis thaliana*. Kivisaar M (ed.). *Appl Environ Microbiol*
916 2021;87, DOI: 10.1128/AEM.00091-21.
- 917** Peñuelas J, Rico L, Ogaya R et al. Summer season and long-term drought increase
918 the richness of bacteria and fungi in the foliar phyllosphere of *Quercus ilex* in a mixed
919 Mediterranean forest. *Plant Biol* 2012;14:565–75.
- 920** R Team. Team. R: A Language and Environment for Statistical Computing, 2015.
921 2021.
- 922** Reynolds JF, Kemp PR, Ogle K et al. Modifying the ‘pulse–reserve’ paradigm for
923 deserts of North America: precipitation pulses, soil water, and plant responses.
924 *Oecologia* 2004;141:194–210.
- 925** Ripley B, Venables B, Bates DM et al. Package ‘MASS.’ *Cran r* 2013;538:113–20.
- 926** Rodriguez RJ, White Jr JF, Arnold AE et al. Fungal endophytes: diversity and
927 functional roles. *New phytologist* 2009;182:314–30.
- 928** Sambrook JR, Russel DW. *Molecular Cloning: A Laboratory Manual.*, 2001.
- 929** Santos JC, Leal IR, Almeida- JS et al. Caatinga: the scientific negligence
930 experienced by a dry tropical forest. *Trop Conserv Sci* 2011;4:276–86.
- 931** Schwinnig S, Sala OE, Loik ME et al. Thresholds, memory, and seasonality:
932 understanding pulse dynamics in arid/semi-arid ecosystems. *Oecologia*
933 2004;141:191–3.

- 934** da Silva JMC, Barbosa LCF, Leal IR *et al.* The Caatinga: Understanding the
935 Challenges. In: Silva José Maria Cardoso da and Leal IR and TM (ed.). *Caatinga:*
936 *The Largest Tropical Dry Forest Region in South America*. Cham: Springer
937 International Publishing, 2017, 3–19.
- 938** da Silva JMC, Leal IR, Tabarelli M. *Caatinga.*, 2017.
- 939** Stone BWG, Jackson CR. Seasonal Patterns Contribute More Towards Phyllosphere
940 Bacterial Community Structure than Short-Term Perturbations. *Microb Ecol*
941 2021;**81**:146–56.
- 942** Tedersoo L, Bahram M, Polme S *et al.* Global diversity and geography of soil fungi.
943 *Science* (1979) 2014;**346**:1052–3.
- 944** Thapa S, Prasanna R. Prospecting the characteristics and significance of the
945 phyllosphere microbiome. *Ann Microbiol* 2018;**68**:229–45.
- 946** Treseder KK, Berlemont R, Allison SD *et al.* Drought increases the frequencies of
947 fungal functional genes related to carbon and nitrogen acquisition. *PLoS One*
948 2018;**13**:1–17.
- 949** Ueda H, Mitsuhashi I, Tabata J *et al.* Extracellular esterases of phylloplane yeast
950 *Pseudozyma antarctica* induce defect on cuticle layer structure and water-holding
951 ability of plant leaves. *Appl Microbiol Biotechnol* 2015;**99**:6405–15.
- 952** Vacher C, Hampe A, Porté AJ *et al.* The Phyllosphere: Microbial Jungle at the Plant–
953 Climate Interface. *Annu Rev Ecol Evol Syst* 2016;**47**:1–24.
- 954** Vellend M. Conceptual synthesis in community ecology. *Quarterly Review of Biology*
955 2010;**85**:183–206.
- 956** Vorholt JA. Microbial life in the phyllosphere. *Nat Rev Microbiol* 2012;**10**:828–40.
- 957** Yan Z, Chen Q, Li C *et al.* Contrasting ecological processes shape the Eucalyptus
958 phyllosphere bacterial and fungal community assemblies. *Journal of Sustainable*
959 *Agriculture and Environment* 2022;**1**:73–83.
- 960** Yang LH, Bastow JL, Spence KO *et al.* *What can we learn from resource pulses?*,
961 2008.
- 962** Yao H, Sun X, He C *et al.* Host identity is more important in structuring bacterial
963 epiphytes than endophytes in a tropical mangrove forest. *FEMS Microbiol Ecol*
964 2020;**96**:1–16.
- 965** Zeisler-Diehl VV, Barthlott W, Schreiber L. Plant Cuticular Waxes: Composition,
966 Function, and Interactions with Microorganisms. *Hydrocarbons, Oils and Lipids:*
967 *Diversity, Origin, Chemistry and Fate*. Cham: Springer International Publishing, 2020,
968 123–38.
- 969**

Periódico proposto: *Brazilian journal of Microbiology*

6. CAPÍTULO 3 - *Carlosrosaea* sp. nov., a new tremellomycetes yeast from Brazilian Seasonally Dry Tropical Forest (Caatinga)

Ciro Ramon Félix^{1,2}, Bruno Emanuel da Silva Nascimento¹, Victor Daniel Firmino dos Santos Tavares¹, Hector M. C. Navarro^{1,2}, Melissa Fontes Landell^{1,*}

¹Universidade Federal de Alagoas, Maceió, AL, Brazil.

²Programa de Pós-graduação em Diversidade Biológica e Conservação nos Trópicos, Universidade Federal de Alagoas, Campus A. C. Simões, Maceió, AL, 57072-900, Brazil.

*Correspondence: Melissa Fontes Landell e-mail: melissa.landell@gmail.com

ORCID of the authors:

Ciro Ramon Félix– 0000-0001-5087-8872

Bruno Emanuel da Silva Nascimento– 0000-0001-6690-0595

Hector M. C. Navarro – 0000-0002-5613-2227

Victor Daniel Firmino dos Santos Tayares - 0000-0003-1492-0097

Melissa Fontes Landell - 0000-0001-6848-0803

26 Abstract

27 Bromeliads are tropical plants that are a substrate rich in yeast taxonomic novelties.
28 In a study carried out in a semi-arid region in Northeastern Brazil, yeast isolates were
29 obtained from leaves of the *Bromelia laciniosa*. By analyzing the regions of the
30 D1/D2 domain of LSU rRNA and the Internal Transcribed Spacer (ITS) region of
31 rRNA, six isolates showed an affinity with the genus *Carlosrosaea*. The dissimilarity
32 with the already described species and the phylogenetic reconstruction indicates that
33 these isolates constitute a new *Carlosrosaea* species. Therefore, we propose the
34 new tremellomycetes yeast *Carlosrosaea* sp. nov. (Trimorphomycetaceae,
35 Basidiomycota). According to genetic analyses, *C. sp. nov.* differs from the
36 *C. vrieseae* by 18-19 bp substitutions in D1/D2 and 50 bp substitutions in ITS and
37 from the *C. foliicola* by 22-23 bp substitutions on D1/D2 and 43 bp substitutions on
38 ITS. Furthermore, *C. sp. nov.* differs from *C. vrieseae* and *C. foliicola* by the
39 ability to produce extracellular starch-like compounds. The holotype of *C. sp. nov.*
40 is CBS XXXX, deposited at Mycobank under code MBXXXX.

41

42 **Keywords:** Caatinga, bromeliad, semiarid, Tremellales, Basidiomycota.

43

44 Introduction

45 Drylands cover 41% of the Earth's continental surface. They are
46 characterized by a water deficit that can vary from severe to moderate, with aridity
47 indexes ≤ 0.65 . These ecosystems are threatened by climate change and human
48 occupation. In addition, they are microbiologically undersampled, with unique
49 biodiversity and provide important ecosystem services in the global cycle of carbon,
50 nitrogen, and water [1–3]. Depending on the intensity of aridity, Drylands can be
51 divided into hyper-arid, arid, semi-arid, and sub-humid [1]. The Caatinga is a semi-
52 arid dryland that occupies 1/10 of the Brazilian territory (~1 million km²) and forms the
53 largest Seasonally Dry Forest in South America [4–6]. In this environment, rainfall is
54 low (between 240 and 1500 mm per year) and is concentrated in three consecutive
55 months [7, 8]. In some points in the Caatinga about 60% of the annual volume of
56 rainfall can occur in a single month [9].

57 The Caatinga is home to 3,347 plant species, about 16% of which are
58 endemic [10]. One of the groups of plants that stands out in this environment is the
59 Bromeliaceae family. This group of plants is almost exclusively neotropical, only one
60 species is found outside the Americas, the species *Pitcarnia feliciana*, which occurs
61 on the West Coast of Africa [11], a biological vestige of when these continents were
62 connected. The family comprises 3,500 species, 50 genera and 8 subfamilies [11,
63 12]. About half of the bromeliad genera and more than half of the subfamilies contain
64 species capable of forming water tanks (phytotelm) [12, 13]. Bromeliads can function
65 as a microscale island ecosystem and perhaps also as a refuge, particularly in times
66 of drought [14].

67 The phyllosphere (aerial part of plants) includes leaves (phyloplane), fruits
68 (carposphere), flowers (anthosphere, including pollen and nectar), stems
69 (caulosphere), and water reservoirs (phytotelm). These environments support one of
70 the greatest diversity of microorganisms on the planet, with representatives of
71 diverse groups such as bacteria, archaea, viruses, filamentous fungi, yeasts,
72 microalgae, protozoa and nematodes [15–19]. Some of the largest reservoirs of
73 taxonomic novelties of filamentous fungi and yeasts are vascular plants (especially in
74 South America), insects and soil [3, 20–22]. There is a large gap in relation to the
75 richness of fungi in the world, a recent estimate is that there are between 2.2 and 3.8

76 million species of fungi, but the number of species correctly named and described is
77 around 120,000 (from 3.15 to 5.45% of the estimated total) [20]. Regarding yeasts,
78 the estimate is around 200,000 species [3] and the diversity of yeasts currently
79 described is between 2,200 and 2,300 species, just over 1% of total wealth [3, 23].

80 The genus *Carlosrosaea* (Trimorphomycetaceae, Tremellales,
81 Basidiomycota) is among the groups of yeasts frequently associated with plants. This
82 genus was proposed by Liu et al. [24] to receive the species *Carlosrosaea vriesea*,
83 isolated from bromeliads in Brazil and previously allocated in the genus *Bullera* [25].
84 Currently, there are five species described in *Carlosrosaea*. The species
85 *Carlosrosaea hohenbergiae* and *Carlosrosaea aechmeae* were described associated
86 with bromeliads from Brazil [26]; and *Carlosrosaea foliicola* and *Carlosrosaea*
87 *simaoensis* associated with leaves from China [27].

88 During studies carried out in bromeliads from Northeast Brazil, eight yeast
89 isolates, through molecular information of the D1/D2 domain of LSU rRNA and the
90 Internal Transcribed Spacer (ITS) region of rRNA, indicated affinity with the genus
91 *Carlosrosaea*. The analyzes indicated that these isolates differ from the five species
92 already described for the group. Therefore, we propose the description of the new
93 tremellomycetes yeasts *Carlosrosaea* sp. nov., the fourth species of the
94 group originating from bromeliads from Brazil.

95

96 Material and methods

97 Sampling area

98 Between 2017 and 2022, leaves of the species *Bromelia laciniosa* Mart. ex
99 Schult. & Schult.f (Bromeliaceae, Bromelioideae) were collected at Tocaia Private
100 Heritage Reserve (RPPN Tocaia), in the municipality of Santana do Ipanema,
101 Alagoas, Northeast Brazil (9°23'08.9"S and 37°15 '22.8"W). The RPPN Tocaia has
102 an area of approximately 20 hectares, a maximum altitude of about 400 meters, and
103 is mainly composed of trees.

104

105 Plant collection, yeast isolation and maintenance

106 Leaves of apparently healthy adult individuals were collected, stored in
107 sterile plastic bags, and transported to the laboratory at room temperature for

108 approximately 4 h. In the laboratory, the leaves were stored at 10-15 °C, and
109 processing was carried out within 24 h after collecting the material in the field. For the
110 processing of the samples, the leaves underwent an initial wash with sterile distilled
111 water to remove possible contaminants such as dust and other exogenous materials.
112 Then, leaf fragments between 5 and 20 cm² were randomly cut until a total area of
113 120 cm² was obtained. Sequentially, the leaf fragments were placed in a 250 mL
114 Erlenmeyer flask containing 75 mL of sterile distilled water. Then this flask was
115 shaken for 30 min at 180 RPM and at room temperature (27±2 °C). After stirring, the
116 wash product was collected in a sterile flask and the leaf fragments were kept in the
117 Erlenmeyer flask. After removing and storing the product from the first wash, 75 mL
118 of sterile Tween 80 solution (0.5% concentration) was added to the flask and the
119 material was stirred again at the same time, rotation and temperature. In the end, the
120 products of the two washes were aggregated, thus forming a pool. A 100 µL fraction
121 of this pool was seeded in duplicate on Petri dishes containing modified YM agar
122 medium (0.3% yeast extract, 0.3% malt extract, 0.5% bacteriological peptone, 1%
123 glucose, 2% agar, 0.04% chloramphenicol, pH 4.0) at concentrations of 10⁰ and 10⁻¹.

124 The plates were incubated at 25-28 °C and observed daily for 10 days for
125 yeast isolation on YEPD agar medium (2% glucose, 1% bacteriological peptone,
126 0.5% yeast extract, and 2% agar). Yeast isolates were preserved in tubes containing
127 GYMP agar medium (2% glucose, 2% malt extract, 0.5% yeast extract, 0.2%
128 monobasic sodium phosphate, and 2% agar) and GYMP broth containing final
129 volume of 30% glycerol.

130

131 **Morphological and physiological characterization**

132 All isolates were morphologically and physiologically characterized by
133 Kurtzman et al. [28]. The carbon and nitrogen assimilation pattern were measured in
134 solid medium using the replica plating technique. To verify the possible production of
135 pseudo-hyphae, true hyphae and/or sexual structures, the isolates were sown on
136 potato dextrose agar (PDA), corn meal agar (CMA) and malt extract agar (MEA) at
137 22-25 °C and observed for 21 days.

138

139 **Molecular analysis**

140 For molecular identification of isolates, genomic DNA was extracted using
141 the protocol for small-scale preparation of yeast DNA modified, originally proposed
142 by Sambrook and Russel [29]. From this, the D1/D2 regions of the 26S LSU and the
143 Internal Trancrit Spacer (ITS) of the rRNA were amplified via PCR. Using primers NL-
144 1 (5'- GCA TATC AAT AAG CGG AGG AAA AG -3') and NL-4 (5'- GG TCC GTG TTT
145 CAA GAC GG -3'); and ITS-1 (5'-TCC GTA GGT GAA CCT GCG G-3') and ITS-4 (5'-
146 TCC TCC GCT TAT TGA TAT GC-3') [30–32]. The reaction parameters were
147 established following Landell et al. [33]. For sequencing, the Sanger method was
148 used through the protocol facilities from the *Plataforma Multusuária de*
149 *Sequenciamento de DNA* at the *Laboratório de Bioinformática e Biologia Evolutiva* of
150 the *Universidade Federal de Pernambuco* in an automated sequencing system ABI
151 3130 Genetic Analyzer using the polymer BigDye v3.1 and POP7 (Life
152 Technologies). Sequence consensus was generated using the Staden Package
153 software [34] and MEGA X. Then, the sequences obtained were compared with
154 others deposited in the GenBank database on the National Center for Biotechnology
155 Information (NCBI) website using the Basic Local Alignment Search Tool (BLAST)
156 and the BLASTn algorithm [35]. To estimate phylogenetic relationships, trees were
157 constructed using D1/D2 regions and ITS applying neighbor-joining method. To
158 reconstruct the phylogenetic relationships, gaps were excluded, and the Kimura two-
159 parameter model was applied to correct the genetic distances [36]. Branch
160 robustness was estimated with the bootstrap resampling method using 1000
161 pseudoreplicates [37].

162

163 Results and Discussion

164 Novel species delineation and identification

165 Eight yeast isolates were isolated from bromeliad leaves of *Bromelia*
166 *laciniosa* Mart. ex Schult. & Schult.f (Bromeliaceae, Bromelioideae) from the
167 Caatinga biome, a semi-arid Dryland inserted in Northeast Brazil (Table 1). Analysis
168 of the D1/D2 regions of the 26S LSU and the Internal transcribed Spacer (ITS) of the
169 rRNA indicated that these isolates have affinity with the genus *Carlosrosaea*, and
170 diverge from type material of all currently described species: *C. vrieseae* (18-19 bp
171 substitutions in D1/D2 and 50 in ITS), *C. hohenbergiae* (20-21 bp substitutions in

172 D1/D2 and 54 in ITS), *C. aechmeae* (22-23 bp substitutions in D1/D2 and 44 in ITS),
 173 *C. foliicola* (22-23 bp substitutions on D1/D2 and 43 on ITS), and *C. simaoensis* (21-
 174 23 bp substitutions on D1/D2 and 46 on ITS) (Fig. 1). Therefore, we propose the
 175 species *C. xxxxxxx sp. nov.* to accommodate these isolates. The isolates of *C.*
 176 *xxxxxxxx sp. nov.* differ from each other by 0-1 bp in the D1/D2 domain and have
 177 identical ITS sequences. Furthermore, *C. xxxxxxx sp. nov.* differs from *C. foliicola*, *C.*
 178 *simaoensis* and *C. vrieseae* by the ability to produce extracellular starch-like
 179 compounds and from *C. aechmeae* by the ability to assimilate sucrose (Table 2).

180

181 **Table 1-** Information on *Carlosrosaea xxxxxxx sp. nov.* isolates: Culture collection
 182 code, GenBank sequence accession number and date of collection.

Culture collection deposit code		Genbank access code		Collection date
CBS	UFMG	UFAL	ITS	D1D2
		BRT303	MH909017	September 21, 2017
		BRT537		October 19, 2021
		BRT586		November 23, 2021
	UFMG-CM- CBS XXT	BRT651		May 24, 2022
	UFMG-CMT	BRT659 ^T		May 24, 2022
		BRT663		May 24, 2022
		BRT696		June 23, 2022
		BRT714		July 19, 2022

183

184 **Table 2-** Selection of physiological/biochemical characteristics that differentiate
 185 *Carlosrosaea xxxxxxx sp. nov.* from other species of the genus.

Characteristic	<i>C. aechmeae</i> ^a	<i>C. foliicola</i> ^b	<i>C. hohenbergiae</i> ^a	<i>C. simaoensis</i> ^b	<i>C. vrieseae</i> ^c	<i>C. xxxxxxx</i>
Carbon source						
Inulin	+	-	+	-	V	D,w
Sucrose	-	+	V	+	V	+
Glycerol	-	+	-	+	V	-
Meso-erythritol	-	+	V	-	-	-
Succinic acid	-	D,w	V	w	V	-
Citrate	-	D,w	-	+	V	-
D-gluconate	-		+		+	w
L-arabinitol	+		+			w
DL-lactate	-	+	-		V	w
Tween 20	-		V		+	w
Nitrogen source						
Cadaverine	+	-	-	-	+	w
Lysine	+	-	+	-	+	w
Starch-like compounds						
			+	-	-	+


186 ^a Data obtained from Félix et al. [26].

187 ^b Data obtained from Li et al. [27].

188 ^c Data obtained from Landell et al. [25].

189

190 *Carlosrosaeae* is a monophyletic genus in the Trimorphomycetaceae
 191 family. The genus is well supported by analysis of the D1/D2 and ITS rRNA regions,
 192 and it has affinity with the genus *Vishniacozyma* and the clade *Tremella sensu stricto*
 193 (Fig. 1). The Trimorphomycetaceae family appears monophyletic from the analysis of
 194 the LSU rRNA region, but its support via bootstrap is not robust [24]. The genus
 195 *Vishniacozyma* and *Tremella* occur in diverse substrates such as soil, plants and
 196 marine environments. In addition, these genera can be found in arctic climate, humid
 197 and semi-arid tropical climates [38–41].
 198



199
 200 **Fig. 1-** Phylogenetic tree of *Carlosrosaea* *xxxxxx* sp. nov obtained by neighbour-
 201 joining (Kimura two-parameter distance method) analysis of the concatenated D1/D2
 202 and ITS regions. Bootstrap values $\geq 50\%$ are show. Bar, 0.05 substitutions per
 203 nucleotide position.

204 Bromeliads are an important substrate for prospecting new yeast isolates
205 and species [42]. More than 180 species of yeasts have already been registered in
206 bromeliads and, together with the phytotelmata, the leaves are one of the most
207 diverse substrates of these plants [42]. All described species of the genus
208 *Carlosrosaea* (Tremellales, Basidiomycota) has association with plant substrates,
209 mainly to bromeliads, with records of at least 13 species of these plants, in flowers,
210 phytotelma and leaves [25–27, 42–44]. The species *C. vriesiae* can stimulate plant
211 growth (evaluated in bromeliads) producing indoleacetic acid (IAA), phosphate
212 solubilization, siderophore production and antagonism to phytopathogens [45].
213 Furthermore, *Carlosrosaea* spp. isolates produced extracellular enzymes such as
214 amylase, cellulase, pectinase, and protease [44]. It indicates, in addition to an
215 important biotechnological potential, a possible ecological role of the genus in
216 stimulating plant growth.

217


218 **Description of *Carlosrosaea* xxxxxxx sp. nov.**

219

Carlosrosaea xxxxxxx

220 After 3 days on YEPD broth at 22–25 °C, yeast cells are globose,
221 subglobose with unipolar or bipolar buds (2.4–3.9 × 3.0–5.1 µm) (Fig. 2a). The
222 colonies are cream to yellowish-white, smooth, and creamy (Fig. 2b). In Dalmau
223 plates after 4 weeks on cornmeal agar, were not observed pseudohypha or hypha
224 formation. Glucose fermentation ability is negative and ballistoconidia production is
225 absent. The species can grow in the presence of 0.01% cycloheximide, but not at
226 0.1%. The ability to grow in medium containing 10 and 16% NaCl was observed, but
227 it was not able to grow in the presence of 50% glucose. There was no growth in the
228 presence of 1% acetic acid. The urease activity, the reaction of diazonium blue B and
229 the starch-like compounds production are positive. The following carbon compounds
230 are assimilated: D-glucose, inulin (slow), sucrose, raffinose, melibiose, D-galactose,
231 lactose (weak), trehalose, maltose (slow), melezitose, soluble starch (variable),
232 cellobiose (weak), L-rhamnose (weak), D-xylose, L-arabinose, D-arabinose (slow), D-
233 ribose, D-ribitol (slow), D-mannitol (weak), D-glucitol (weak), myo-inositol (weak), D-
234 gluconate (weak), D-glucosamine (weak), N-acetylglucosamine (weak), D-
235 galacturonic acid (weak), xylitol (variable), L-arabinitol (weak), DL-lactate (weak),
236 Tween 20 (weak) and Tween 80 (variable). The following carbon compounds are not

237 assimilated: salicin, glycerol, meso-erythritol, succinic acid and citrate. The
 238 assimilated nitrogen compounds were nitrite (weak), cadaverine (weak) and lysine
 239 (weak). The following nitrogen compounds are not assimilated: nitrate, creatinine,
 240 creatine, ethylamine.



241

242 **Fig. 2-** Cellular (a) and colonial (b) morphology of *C. xxxxxxx* sp. nov. grown in YEPD
 243 for 48 h at 22-25 °C.

244

245 The holotype of *C. xxxxxxx* sp. nov. CBS XXXX, is preserved in a
 246 metabolically inactive state in the CBS Yeast Collection of the Westerdijk Fungal
 247 Biodiversity Institute, Utrecht, Netherlands and in the Collection of Microorganisms; in
 248 the Cells of Federal University of Minas Gerais (*Coleção de Microrganismos e*
 249 *Células da Universidade Federal de Minas Gerais- UFMG*), Belo Horizonte, Minas
 250 Gerais, Brazil as the strain UFMG-CM-XXX; and in the Mycotheque of the Molecular
 251 Diversity Laboratory of the Federal University of Alagoas (*Micoteca do Laboratório de*
 252 *Diversidade Molecular da Universidade Federal de Alagoas - UFAL*), as the strain
 253 BRT659. Mycobank number is XXXXXX. Additional isolates are indicated in Table 1.

254

255 Acknowledgements

256 We thank the Mr. Alberto Nepomuceno Agra Jr. for authorization and assistance
 257 during the collections. This work is part of the project "INCT Yeasts: Biodiversity,

258 preservation and biotechnological innovation", funded by Conselho Nacional de
259 Desenvolvimento Científico e Tecnológico (CNPq), grant #406564/2022-1.

260

261 **References**

- 262** 1. Huang J, Ji M, Xie Y, et al (2016) Global semi-arid climate change over last 60
263 years. *Clim Dyn* 46:1131–1150. <https://doi.org/10.1007/s00382-015-2636-8>
- 264** 2. Maestre FT, Eldridge DJ, Soliveres S, et al (2016) Structure and Functioning
265 of Dryland Ecosystems in a Changing World. *Annu Rev Ecol Evol Syst* 47:215–237.
266 <https://doi.org/10.1146/annurev-ecolsys-121415-032311>
- 267** 3. Boekhout T, Amend AS, el Baidouri F, et al (2022) Trends in yeast diversity
268 discovery. *Fungal Divers* 114:491–537. <https://doi.org/10.1007/s13225-021-00494-6>
- 269** 4. Moro MF, Lughadha EN, Araújo FS de, Martins FR (2016) A
270 Phytoogeographical Metaanalysis of the Semiarid Caatinga Domain in Brazil. *The*
271 *Botanical Review*. <https://doi.org/10.1007/s12229-016-9164-z>
- 272** 5. Machado IC, Lopes AV (2004) Floral Traits and Pollination Systems in the
273 Caatinga , a Brazilian Tropical Dry Forest. *Ann Bot* 94:365–376.
274 <https://doi.org/10.1093/aob/mch152>
- 275** 6. da Silva JMC, Leal IR, Tabarelli M (2017) Caatinga
- 276** 7. da Silva JMC, Barbosa LCF, Leal IR, Tabarelli M (2017) The Caatinga:
277 Understanding the Challenges. In: Silva José Maria Cardoso da and Leal IR and TM
278 (ed) *Caatinga: The Largest Tropical Dry Forest Region in South America*. Springer
279 International Publishing, Cham, pp 3–19
- 280** 8. Santos JC, Leal IR, Almeida- JS, et al (2011) Caatinga : the scientific
281 negligence experienced by a dry tropical forest. *Trop Conserv Sci* 4:276–286
- 282** 9. Menezes RSC, Sampaio EVSB, Giongo V, Pérez-Marin AM (2012)
283 Biogeochemical cycling in terrestrial ecosystems of the Caatinga Biome. *Braz J Biol*
284 72:643–653
- 285** 10. Fernandes MF, Cardoso D, de Queiroz LP (2020) An updated plant checklist
286 of the Brazilian Caatinga seasonally dry forests and woodlands reveals high species
287 richness and endemism. *J Arid Environ* 174:.
288 <https://doi.org/10.1016/j.jaridenv.2019.104079>
- 289** 11. Benzing DH (2000) *Bromeliaceae: profile of an adaptive radiation*. Cambridge
290 University Press, New York
- 291** 12. Ladino G, Ospina-Bautista F, Estévez Varón J, et al (2019) Ecosystem
292 services provided by bromeliad plants: A systematic review. *Ecol Evol* 9:7360–7372.
293 <https://doi.org/10.1002/ece3.5296>

- 294** 13. Males J, Griffiths H (2017) Functional types in the Bromeliaceae: relationships
295 with drought-resistance traits and bioclimatic distributions. *Funct Ecol* 31:1868–1880.
296 <https://doi.org/10.1111/1365-2435.12900>
- 297** 14. Frank JH, Lounibos LP (1987) Phytotelmata: swamps or islands? *Flr Entom*
298 70:14–20
- 299** 15. Bringel F, Couée I (2015) Pivotal roles of phyllosphere microorganisms at the
300 interface between plant functioning and atmospheric trace gas dynamics. *Front*
301 *Microbiol* 6:1–14. <https://doi.org/10.3389/fmicb.2015.00486>
- 302** 16. Koskella B (2020) The phyllosphere. *Curr Biol* 30:R1143–R1146.
303 <https://doi.org/10.1016/j.cub.2020.07.037>
- 304** 17. Leveau JH (2019) A brief from the leaf: latest research to inform our
305 understanding of the phyllosphere microbiome. *Curr Opin Microbiol* 49:41–49.
306 <https://doi.org/10.1016/j.mib.2019.10.002>
- 307** 18. Vacher C, Hampe A, Porté AJ, et al (2016) The Phyllosphere: Microbial Jungle
308 at the Plant–Climate Interface. *Annu Rev Ecol Evol Syst* 47:1–24.
309 <https://doi.org/10.1146/annurev-ecolsys-121415-032238>
- 310** 19. Vorholt JA (2012) Microbial life in the phyllosphere. *Nat Rev Microbiol* 10:828–
311 840. <https://doi.org/10.1038/nrmicro2910>
- 312** 20. Hawksworth DL, Lücking R (2017) Fungal Diversity Revisited: 2.2 to 3.8
313 Million Species. In: *The Fungal Kingdom*. ASM Press, Washington, DC, USA, pp 79–
314 95
- 315** 21. Blackwell M (2017) Yeasts in Insects and Other Invertebrates. In: Buzzini P,
316 Lachance M-A, Yurkov A (eds) *Yeasts in Natural Ecosystems: Diversity*. Springer
317 International Publishing, Cham, pp 397–433
- 318** 22. Blackwell M (2011) The fungi: 1, 2, 3 ... 5.1 million species? *Am J Bot* 98:426–
319 438. <https://doi.org/10.3732/ajb.1000298>
- 320** 23. Yurkov A, Alves A, Bai FY, et al (2021) Nomenclatural issues concerning
321 cultured yeasts and other fungi: why it is important to avoid unneeded name
322 changes. *IMA Fungus* 12:. <https://doi.org/10.1186/s43008-021-00067-x>
- 323** 24. Liu XZ, Wang QM, Göker M, et al (2015) Towards an integrated phylogenetic
324 classification of the Tremellomycetes. *Stud Mycol* 81:85–147.
325 <https://doi.org/10.1016/j.simyco.2015.12.001>
- 326** 25. Landell MF, Brandão LR, Safar SVB, et al (2015) *Bullera vrieseae* sp. nov., a
327 tremellaceous yeast species isolated from bromeliads. *Int J Syst Evol Microbiol*
328 65:2466–2471. <https://doi.org/10.1099/ijst.0.000285>
- 329** 26. Felix CR, Navarro HMC, Paulino GVB, et al (2017) *Carlosrosaea*
330 *hohenbergiae* sp. nov. and *Carlosrosaea aechmeae* sp. nov., two tremellaceous

- 331** yeasts isolated from bromeliads in north-eastern Brazil. *Int J Syst Evol Microbiol*
332 67:1752–1757. <https://doi.org/10.1099/ijsem.0.001856>
- 333** 27. Li AH, Yuan FX, Groenewald M, et al (2020) Diversity and phylogeny of
334 basidiomycetous yeasts from plant leaves and soil: Proposal of two new orders, three
335 new families, eight new genera and one hundred and seven new species. *Stud Mycol*
336 96:17–140. <https://doi.org/10.1016/j.simyco.2020.01.002>
- 337** 28. Kurtzman CP, Fell JW, Boekhout T, Robert V (2011) Methods for isolation,
338 phenotypic characterization and maintenance of yeasts. In: *The Yeasts*. Elsevier
339 B.V., pp 87–110
- 340** 29. Sambrook JR, Russel DW (2001) *Molecular Cloning: A Laboratory Manual*
- 341** 30. Kurtzman CP, Robnett CJ (1998) Identification and phylogeny of
342 ascomycetous yeasts from analysis of nuclear large subunit (26S) ribosomal DNA
343 partial sequences. *Anton Leeuw Int J G*, 73:331–371.
344 <https://doi.org/10.1023/A:1001761008817>
- 345** 31. Fell JW, Boekhout T, Fonseca A, et al (2000) Biodiversity and systematic of
346 basidiomycetous yeast as determined by large submit rDNA D1/D2 domain
347 sequence analysis. *Int J Syst Evol Microbiol* 50:1351–1371.
348 <https://doi.org/10.1099/00207713-50-3-1351>
- 349** 32. Schoch CL, Seifert K a., Huhndorf S, et al (2012) Nuclear ribosomal internal
350 transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proc*
351 *Natl Acad Sci USA* 109:1–6
- 352** 33. Landell MF, Billodre R, Ramos JP, et al (2010) *Candida aechmeae* sp. nov.
353 and *Candida vrieseae* sp. nov., novel yeast species isolated from the phylloplane of
354 bromeliads in Southern Brazil. *Int J Syst Evol Microbiol* 60:244–248.
355 <https://doi.org/10.1099/ijsem.0.011577-0>
- 356** 34. Staden R, Beal KF, Bonfield JK (2000) The staden package. *Bioinformatics*
357 methods and protocols 115–130
- 358** 35. Altschul SF, Madden TL, Schäffer AA, et al (1997) Gapped BLAST and PSI-
359 BLAST: A new generation of protein database search programs. *Nucleic Acids Res*
360 25:3389–3402. <https://doi.org/10.1093/nar/25.17.3389>
- 361** 36. Kimura M (1980) A Simple Method for Estimating Evolutionary Rates of Base
362 Substitutions Through Comparative Studies of Nucleotide Sequences. *J Mol Evol*
363 16:111–120
- 364** 37. Felsenstein J (1985) Confidence limits on phylogenies: an approach using the
365 bootstrap. *Evolution (NY)* 39:783–791
- 366** 38. Tsuji M, Tanabe Y, Vincent WF, Uchida M (2019) *Vishniacozyma*
367 *ellesmerensis* sp. nov., a psychrophilic yeast isolated from a retreating glacier in the
368 Canadian High Arctic. *Int J Syst Evol Microbiol* 69:696–700

- 369** 39. Félix CR, Andrade DA, Almeida JH, et al (2020) *Vishniacozyma alagoana* sp.
370 nov. a tremellomycetes yeast associated with plants from dry and rainfall tropical
371 forests. *Int J Syst Evol Microbiol* 70:3449–3454.
372 <https://doi.org/10.1099/ijsem.0.004193>
- 373** 40. Khunnamwong P, Surussawadee J, Ribeiro JRA, et al (2019) *Tremella*
374 *saccharicola* f.a., sp. nov., a novel tremellaceous yeast species isolated from tropical
375 regions. *Int J Syst Evol Microbiol* 69:2010–2016.
376 <https://doi.org/10.1099/ijsem.0.003420>
- 377** 41. Navarro HMC, Félix CR, Tavares VDFS, et al (2022) *Tremella ananatis* sp.
378 nov. and *Tremella lamprococci* sp. nov., two yeast species associated with
379 bromeliads. *Int J Syst Evol Microbiol* 72: <https://doi.org/10.1099/ijsem.0.005261>
- 380** 42. Félix CR, da Silva Nascimento BE, Valente P, Landell MF (2022) Different
381 plant compartments, different yeasts: The example of the bromeliad phyllosphere.
382 *Yeast* 39:363–400. <https://doi.org/10.1002/yea.3804>
- 383** 43. Félix CR, Navarro HMC, Almeida JH, Landell MF (2021) Behind the nectar:
384 the yeast community in bromeliads inflorescences after the exudate removal. *Mycol*
385 *Prog* 20:1191–1202. <https://doi.org/https://doi.org/10.1007/s11557-021-01728-2>
- 386** 44. Navarro HMC, Félix CR, Paulino GVB, et al (2020) Richness and
387 biotechnological potential of the yeast community associated with the bromeliad
388 phylloplane in the Brazilian Neotropical Forest. *Mycol Prog* 19:1387–1401.
389 <https://doi.org/10.1007/s11557-020-01631-2>
- 390** 45. Marques AR, Resende AA, Gomes FCO, et al (2021) Plant growth-promoting
391 traits of yeasts isolated from the tank bromeliad *Vriesea minarum* L.B. Smith and the
392 effectiveness of *Carlosrosaea vrieseae* for promoting bromeliad growth. *Braz J*
393 *Microbiol* 52:1417–1429. <https://doi.org/10.1007/s42770-021-00496-1>
- 394**
- 395**

7. DISCUSSÃO GERAL

Plantas são fontes importantes de micro-organismos e, ao mesmo tempo, os micro-organismos que formam a microbiota das plantas são importantes nos processos ecológicos e evolutivos dos vegetais (KOSKELLA, 2020; MEYER; LEVEAU, 2012; MORRIS, 2001; VACHER et al., 2016; VORHOLT, 2012). Uma fração ínfima da microbiota do planeta é conhecida, dependendo do grupo, apenas 1% do total (BOEKHOUT et al., 2022; HAWKSWORTH; LÜCKING, 2017). Conhecer a biodiversidade se faz cada vez mais necessário, e em um cenário de mudanças climáticas, a interação planta-microbiota pode permitir que os hospedeiros estendam seus limites de tolerância, entretanto, para compreender as minúcias e o real potencial desse e de outros mecanismos é necessário conhecer os grupos microbianos e suas funções.

Bromélias são plantas tropicais ricas em diversidade de leveduras (MORAIS; DE SOUSA; ROSA, 2020; NAVARRO et al., 2020). Mais de 180 espécies de leveduras já foram registradas em bromélias. Grupos de leveduras frequentemente encontrados na filosfera de bromélias envolvem gêneros como *Candida*, *Rhodotorula*, *Hannaella*, *Aureobasidium*, *Meyerozyma*, *Carlosrosaea* e *Pseudozyma*. Grupos anteriormente inserido no gênero *Cryptococcus* e que atualmente estão distribuídos em outros gêneros estão entre os mais frequentes da filosfera, são eles: *Kwoniella*, *Naganishia*, *Papiliotrema*, *Saitozyma* e *Vishniacozyma* (LIU et al., 2015). As espécies que mais se destacaram na filosfera de bromélias foram *Papiliotrema laurentii* e *Papiliotrema flavescens*. Muitos desses grupos já possuem registros como membros dominantes da filosfera, principalmente em folhas (FONSECA; INÁCIO, 2006).

Além disso, essa diversidade de leveduras se distribui por vários compartimentos da filosfera como folhas, flores, frutos e tanques. Cada um desses compartimentos possui uma comunidade de leveduras distinta, que responde a estrutura única de cada compartimento. A interação com a cutícula pode ser um modulador importante da estrutura da comunidade e isso pode explicar por que leveduras em folhas, frutos intactos e flores (quando o néctar é desconsiderado) tende a ser estruturalmente semelhantes (FONSECA; INÁCIO, 2006). Plantas frequentemente apresentam comunidades microbianas distintas a depender do

compartimento estudado (ABDELFATTAH et al., 2019; LIU; HOWELL, 2021). Leveduras em bromélias parecem seguir o mesmo padrão e divergir em configuração dependendo do compartimento.

Não há consenso sobre o efeito das chuvas nas comunidades microbianas da filosfera. A microbiota pode se apresentar mais resilientes a eventos de chuva de curto prazo e não manifestar alterações significativas (e.g., STONE; JACKSON, 2021), ou responder de maneira mais imediata a eventos disruptivos de chuva (e.g., ALLARD; OTTESEN; MICALLEF, 2020). A filosfera é um ambiente dinâmico e sujeito a flutuações em muitos fatores ambientais. Micro-organismos da filosfera em ambientes áridos possuem camadas de sobreposição de estresse hídrico, uma gerada pela cutícula da planta que dificulta a lixiviação de metabólitos e limita a molhabilidade, e outra pelo próprio ambiente árido que na maior parte do tempo se encontra em déficit hídrico (KOSKELLA, 2020; MAESTRE et al., 2016; NOY-MEIR, 1973; OSO et al., 2021; VACHER et al., 2016).

Em ambientes árido, as chuvas não essenciais e controlam os processos biológicos (MAESTRE et al., 2016; NOY-MEIR, 1973). É de se esperar que nesses ambientes a microbiota da filosfera responda aos pulsos de chuva, principalmente porque quantidades pequenas de chuva geralmente são suficientes para eliciar a comunidade microbiana em regiões secas (REYNOLDS et al., 2004). Entretanto, a comunidade de leveduras em folhas de bromélias na Caatinga apresentou pouca influência da chuva na diversidade, ao mesmo tempo que a sazonalidade mostrou um efeito importante na estrutura da comunidade.

As leveduras são organismos heterotróficos e as enzimas extracelulares desempenham um papel fundamental na sua nutrição, elas hidrolisam macromoléculas e disponibilizam nutrientes para que a célula utilize. Dentre as enzimas avaliadas, a esterase foi a mais produzida por leveduras da filosfera de bromélias. As esterases e outras hidrolases, como a celulase produzida por fungos, são importantes no fluxo de carbono nos ambientes (TRESEDER et al. 2018). *In vitro*, as esterases produzidas pela espécie de levedura epífita *Pseudozyma antarctica* foram capazes de afetar a cutícula da planta e influenciar sua dinâmica hídrica (UEDA et al. 2015). Além disso, essa mesma esterase foi capaz de agravar a infecção causada por *Botrytis cinerea* em tomateiros (Ueda et al. 2018).

Em bromélias, não se observou diferenças significativas em relação aos componentes da diversidade funcional entre os períodos. Os traços mais expressos na filosfera foram relacionados a assimilação de fontes de carbono, principalmente oligo e monossacarídeos derivados de plantas como rafinose, glicose e xilose, mas também polissacarídeos como inulina. A capacidade de assimilar uma grande gama de compostos é uma das características que permite às leveduras conquistar diferentes habitats. Hagler et al. (1993), indicaram que leveduras de tanque de bromélias possuem uma natureza politrófica, mas da metade da comunidade de leveduras recuperada assimilou mais de 20 diferentes fontes de carbono.

O índice de emulsão (IE_{24}) foi um dos poucos traços testados que foi mais frequente durante o período seco em leveduras na Caatinga. Este índice é indicativo de uma das atividades de um surfactante. Biossurfactantes são moléculas quimicamente diversas produzidas por vários grupos microbianos cuja principal característica é uma estrutura anfipática. Essas moléculas têm inúmeras atividades, como: diminuir a tensão superficial, aumentar a molhabilidade em superfícies hidrofóbicas, emulsificar líquidos imiscíveis e solubilizar hidrocarbonetos (BEATTIE; LINDOW, 1995; THAPA; PRASANNA, 2018; ZEISLER-DIEHL; BARTHLOTT; SCHREIBER, 2020; OSO et al., 2021). A produção de biossurfactantes pode ser uma ferramenta importante para microrganismos epífitos aumentarem o acesso a água e nutrientes (LINDOW; BRANDL, 2003; LEVEAU, 2019). A maior frequência de emulsão durante a seca pode indicar que na Caatinga esse traço é favorecido durante períodos secos e promove um aumento do acesso à água para as leveduras epifíticas. No entanto, mais dados e estudos são necessários para avaliar a validade dessa hipótese.

8. CONSIDERAÇÕES FINAIS

Este estudo contribuiu com o conhecimento em biodiversidade e ecologia de leveduras da filosfera de bromélias. Corroboramos que plantas são substratos ricos em micro-organismos e que bromélias possuem potencial para prospecção de leveduras nos campos de: novidades taxonômicas, matéria-prima biotecnológica e funcionamento ecossistêmico. Além disso, constata-se que ainda há uma grande

lacuna sobre biodiversidade de leveduras em bromélias, e principalmente sobre os processos que esses micro-organismos desempenham.

É evidente que ainda há muito a ser explorado nas bromélias. Atualmente, nenhum estudo sobre leveduras de néctar de bromélias foi desenvolvido, mesmo com a estrutura singular da família Bromeliaceae possuir, como o grande espectro de polinizadores e de composição química do néctar. Entre os anos 1993-2021, a maioria dos estudos sobre leveduras em bromélias se concentrou no levantamento descritivo de comunidades. A importância desse referencial de conhecimento é inegável. Entretanto, é necessário dar os próximos passos e aumentar o número de estudos que buscam entender os processos que estruturam a comunidade de leveduras e os mecanismos funcionais providos por essa comunidade e não apenas verificar os padrões.

Os resultados trazem informações sem precedentes sobre a dinâmica da microbiota de levedura em uma região semiárida importante como a Caatinga. O presente estudo acrescenta informações sobre a tolerância da comunidade de leveduras da filosfera a mudanças sazonais, de uma perspectiva taxonômica, filogenética e funcional. Além disso, ao entender o efeito da chuva e da seca na microbiota da filosfera, podemos pensar em como a comunidade microbiana das folhas será afetada pelas mudanças nos ciclos seco-úmido que podem ser causadas pelas mudanças climáticas. A microbiota da filosfera é um fator importante para a resiliência e manutenção da fitossanidade, consequentemente, a disbiose dessa comunidade pode influenciar na saúde do hospedeiro.

9. REFERÊNCIAS

- ABDELFATTAH, A. et al. Revealing Cues for Fungal Interplay in the Plant–Air Interface in Vineyards. **Frontiers in Plant Science**, v. 10, p. 1–10, 2019.
- ALLEN, K. et al. Will seasonally dry tropical forests be sensitive or resistant to future changes in rainfall regimes? **Environmental Research Letters**, v. 12, n. 2, 2017.
- ALTSCHUL, S. F. et al. Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. **Nucleic Acids Research**, v. 25, n. 17, p. 3389–3402, 1997.
- ANDREOTE, F. D.; GUMIERE, T.; DURRER, A. Exploring interactions of plant microbiomes. **Scientia agricola**, v. 71, n. 6, p. 528–539, 2014.
- ANDREWS, J. H et al. Fungi, Leaves, and the Theory of Island Biogeography. **Microbial ecology**, v. 14, n. 3, p. 277–290, 1987.
- AUSTIN, A. T. et al. Water pulses and biogeochemical cycles in arid and semiarid ecosystems. **Oecologia**, v. 141, n. 2, p. 221–235, 2004.
- BASÍLIO, G. A. et al. Community ecology of epiphytic Bromeliaceae in a remnant of Atlantic Forest in Zona da Mata, Minas Gerais State, Brazil. **Hoehnea**, v. 42, n. 1, p. 21–31, 2015.
- BASTIDAS, R. J.; HEITMAN, J. Trimorphic stepping stones pave the way to fungal virulence. **PNAS**, v. 106, n. 2, p. 351–352, 2009.
- BEATTIE, G. A; LINDOW, S. E. The Secret Life of Foliar Bacterial Pathogens on Leaves. **Annual Review of Phytopathology**, v. 33, n. 1, p. 145–172, set. 1995.
- BECKNELL, J. M.; KISSING KUCEK, L.; POWERS, J. S. Aboveground biomass in mature and secondary seasonally dry tropical forests: A literature review and global synthesis. **Forest Ecology and Management**, v. 276, p. 88–95, 2012.
- BEGON, M.; TOWNSEND, C. R. **Ecology: from individuals to ecosystems**. John Wiley & Sons, 2020.
- BENZING, D. H. **Bromeliaceae: profile of an adaptive radiation**. New York: Cambridge University Press, 2000.
- BEZERRA, J. D. P. et al. *Valentiella maceioensis* gen. et sp. nov. (Herpotrichiellaceae, Chaetothyriales), a new black yeast-like fungus isolated from bromeliads in Brazil. **Mycological Progress**, v. 21, n. 2. 2022.
- BLACKWELL, M. Yeasts in Insects and Other Invertebrates. In: **Yeasts in Natural Ecosystems: Diversity**. Cham: Springer International Publishing. p. 397–433. 2017
- BOEKHOUT, T. et al. Discussion of Teleomorphic and Anamorphic Basidiomycetous Yeasts. In: **The Yeasts**. p. 293–307. 2011.

- BOEKHOUT, T. et al. Trends in yeast diversity discovery. **Fungal Diversity**, v. 114, n. 1, p. 491–537, 2022.
- BOTHA, A. The importance and ecology of yeasts in soil. **Soil Biology and Biochemistry**, v. 43, n. 1, p. 1–8, 2011.
- BRINGEL, F.; COUÉE, I. Pivotal roles of phyllosphere microorganisms at the interface between plant functioning and atmospheric trace gas dynamics. **Frontiers in Microbiology**, v. 6, p. 1–14, 2015.
- BUCK, J. W. In vitro antagonism of *Botrytis cinerea* by phylloplane yeasts. **Canadian Journal of Botany**, v. 80, p. 885–891, 2002.
- BUZZINI, P.; MARTINI, A. Extracellular enzymatic activity profiles in yeast and yeast-like isolates isolated from tropical environments. **Journal of Applied Microbiology**, v. 93, p. 1020–1025, 2002.
- CANTO, A.; HERRERA, C. M.; RODRIGUEZ, R. Nectar-living yeasts of a tropical host plant community: diversity and effects on community-wide floral nectar traits. **PeerJ**, v. 5, p. 1–22. 2017.
- CARRASCO, M. et al. Diversity and extracellular enzymatic activities of yeasts isolated from King George Island, the sub-Antarctic region. **BMC microbiology**, v. 12, p. 251, 2012.
- CHESSON, P. et al. Resource pulses, species interactions, and diversity maintenance in arid and semi-arid environments. **Oecologia**, v. 141, n. 2, p. 236–253, 2004.
- COLEMAN-DERR, D. et al. Plant compartment and biogeography affect microbiome composition in cultivated and native Agave species. **New Phytologist**, v. 209, p. 798–811, 2016.
- COLLINS, S. L. et al. A multiscale, hierarchical model of pulse dynamics in arid-land ecosystems. **Annual Review of Ecology, Evolution, and Systematics**, v. 45, p. 397–419, 2014.
- COOPER, C. R. Yeasts Pathogenic to Humans. In: **The Yeasts**. Fifth Edition ed. London: Elsevier, p. 9–19. 2011.
- DA SILVA, J. M. C. et al. The Caatinga: Understanding the Challenges. In: **Caatinga: The Largest Tropical Dry Forest Region in South America**. Cham: Springer International Publishing, p. 3–19. 2017.
- DA SILVA, J. M. C; LEAL, I. R.; TABARELLI, M. **Caatinga**. 2017.
- DE ALBUQUERQUE, U. P. et al. Medicinal plants of the caatinga (semi-arid) vegetation of NE Brazil: A quantitative approach. **Journal of Ethnopharmacology**, v. 114, n. 3, p. 325–354. 2007.

- DOAN, H. K.; LEVEAU, J. H. J. Artificial Surfaces in Phyllosphere Microbiology. **Phytopathology**, v. 105, n. 8, p. 1036–1042, 2015.
- FÉLIX, C. R. et al. Different plant compartments, different yeasts: The example of the bromeliad phyllosphere. **Yeast**, v. 39, n. 6–7, p. 363–400, 2022.
- FELL, J. W. et al. Biodiversity and systematic of basidiomycetous yeast as determined by large submit rDNA D1/D2 domain sequence analysis. **International Journal of systematic and Evolutionary Microbiology**, v. 50, p. 1351–1371, 2000.
- FERNANDES, M. F.; CARDOSO, D.; DE QUEIROZ, L. P. An updated plant checklist of the Brazilian Caatinga seasonally dry forests and woodlands reveals high species richness and endemism. **Journal of Arid Environments**, v. 174, 2019, p. 104079, 2020.
- FERREIRA, J. VA; FABRICANTE, J. R.; SIQUEIRA-FILHO, J. A. Checklist preliminar de Bromeliaceae do Parque Nacional do Catimbau, Pernambuco, Brasil. **Natureza on line**, v. 13, n. 2, p. 92–97, 2015.
- FONSECA, Á.; INÁCIO, J. Phyloplane Yeasts. **Biodiversity and Ecophysiology of Yeasts**. p. 263–301. 2006.
- FRANK, J. H.; LOUNIBOS, L. P. Phytotelmata: swamps or islands? **Florida Entomologist**, v. 70, n. 1, p. 14–20, 1987.
- GALETTI, L.; BERNARDELLO, L. M. Extrafloral nectaries that attract ants in Bromeliaceae: structure and nectar composition. **Canadian Journal of Botany**, v. 70, n. 6, p. 1101–1106, 1992.
- GANTER, P. F.; MORAIS, P. B.; ROSA, C. A. Yeasts in Cacti and Tropical Fruit. **Yeasts in Natural Ecosystems: Diversity**. p. 225–264. 2017.
- GLUSHAKOVA, A. M.; CHERNOV, I. Y. Seasonal dynamics in a yeast population on the *Oxalis acetosella* L. leaves. **Mikrobiologiya**, v. 73, n. 2, p. 226–32, 2004.
- GLUSHAKOVA, A. M.; CHERNOV, I. Y. Seasonal dynamic of the numbers of epiphytic yeasts. **Microbiology**, v. 76, n. 5, p. 590–595, 2007.
- GOFFREDI, S. K.; JANG, G. E.; HAROON, M. F. Transcriptomics in the tropics: Total RNA-based profiling of Costa Rican bromeliad-associated communities. **Computational and Structural Biotechnology Journal**, v. 13, p. 18–23, 2015.
- GOUKA, L.; RAAIJMAKERS, J. M.; CORDOVEZ, V. Ecology and functional potential of phyllosphere yeasts. **Trends in Plant Science**, v. 27, n. 11, p. 1109–1123, 2022.
- HAGLER, A. N. et al. Yeasts and coliform bacteria of water accumulated in bromeliads of mangrove and sand dune ecosystems of southeast Brazil. **Canadian Journal of Microbiology**, v. 39, n. 10, p. 973–977, 1993.
- HAWKSWORTH, D. L.; LÜCKING, R. Fungal Diversity Revisited: 2.2 to 3.8 Million Species. **The Fungal Kingdom**. p. 79–95. 2017.

- HOLT, R. D. Theoretical perspectives on resource pulses. **Ecology**, v. 89, n. 3, p. 671–681. 2008.
- HUANG, J. et al. Global semi-arid climate change over last 60 years. **Climate Dynamics**, v. 46, n. 3–4, p. 1131–1150, 2016.
- KANE, R. P. Relationship between the southern oscillation / El Nino and rainfall in some tropical and midlatitude regions. **Proceedings of the Indian Academy of Science**, v. 98, n. 3, p. 223–235, 1989.
- KEMLER, M. et al. Phyloplane Yeasts in Temperate Climates. In: **Yeasts in Natural Ecosystems: Diversity**. Cham: Springer International Publishing, p. 171–197. 2017.
- KINKEL, L. Microbial population dynamics on leaves. **Annual Review of Phytopathology**, v. 35, n. 1, p. 327–347. 1997.
- KOSKELLA, B. The phyllosphere. **Current Biology**, v. 30, n. 19, p. R1143–R1146, 2020.
- KOTTEK, M. et al. World map of the Köppen-Geiger climate classification updated. **Meteorologische Zeitschrift**, v. 15, n. 3, p. 259–263. 2006.
- KRÖMER, T. et al. Nectar sugar composition and concentration in relation to pollination syndromes in Bromeliaceae. **Plant Biology**, v. 10, n. 4, p. 502–511. 2008.
- KURTZMAN, C. P. Discussion of Teleomorphic and Anamorphic Ascomycetous Yeasts and Yeast-like Taxa. **The Yeasts**, p. 293–307, 2011a.
- KURTZMAN, C. P. et al. Methods for isolation, phenotypic characterization and maintenance of yeasts. **The Yeasts**. Elsevier, p. 87–110. 2011.
- KURTZMAN, C. P. et al. Phylogeny of the ascomycetous yeasts and the renaming of *Pichia anomala* to *Wickerhamomyces anomalus*. **Antonie van Leeuwenhoek**, v. 99, n. 1, p. 13–23. 2011b.
- KURTZMAN, C. P.; ROBNETT, C. J. Identification and phylogeny of ascomycetous yeasts from analysis of nuclear large subunit (26S) ribosomal DNA partial sequences. **Antonie van Leeuwenhoek, International Journal of General and Molecular Microbiology**, v. 73, n. 4, p. 331–371, 1998.
- LADINO, G. et al. Ecosystem services provided by bromeliad plants: A systematic review. **Ecology and Evolution**, v. 9, n. 12, p. 7360–7372. 2019.
- LANDELL, M. F. et al. *Bullera vrieseae* sp. nov., a tremellaceous yeast species isolated from bromeliads. **International Journal of Systematic and Evolutionary Microbiology**, v. 65, n. 8, p. 2466–2471, 2015.
- LANDELL, M. F. et al. *Candida aechmeae* sp. nov. and *Candida vrieseae* sp. nov., novel yeast species isolated from the phylloplane of bromeliads in Southern Brazil. **International Journal of Systematic and Evolutionary Microbiology**, v. 60, n. 1, p. 244–248, 2010.

- LANDELL, M. F.; MAUTONE, J. N.; VALENTE, P. Biodiversity of Yeasts Associated to Bromeliads in Itapuã Park, Viamão / Rs. **Biociencias**, v. 14, n. 2, p. 144–149, 2006.
- LEAL, I. R. et al. Mudando o curso da conservação da biodiversidade na Caatinga do Nordeste do Brasil. **Megadiversidade**, v. 1, n. 1, p. 139–146, 2005.
- LEROY, C. et al. The contribution of microorganisms and metazoans to mineral nutrition in bromeliads. **Journal of Plant Ecology**, v. 9, n. 3, p. 241–255, 2016.
- LEROY, C. et al. What drives detrital decomposition in neotropical tank bromeliads? **Hydrobiologia**, v. 802, n. 1, p. 85–95. 2017.
- LEVEAU, J. H. J. A brief from the leaf: latest research to inform our understanding of the phyllosphere microbiome. **Current Opinion in Microbiology**, v. 49, p. 41–49, 2019.
- LI, A. H. et al. Diversity and phylogeny of basidiomycetous yeasts from plant leaves and soil: Proposal of two new orders, three new families, eight new genera and one hundred and seven new species. **Studies in Mycology**, v. 96, p. 17–140. 2020.
- LINDOW, S. E.; BRANDL, M. T. Microbiology of the Phyllosphere. **Applied and environmental microbiology**, v. 69, n. 4, p. 1875–1883, 2003.
- LINDOW, S. E.; LEVEAU, J. H. J. Phyllosphere microbiology. **Current Opinion in Biotechnology**, v. 13, n. 3, p. 238–243, 2002.
- LIU, D.; HOWELL, K. Community succession of the grapevine fungal microbiome in the annual growth cycle. **Environmental Microbiology**, v. 23, n. 4, p. 1842–1857, 2021.
- LIU, H.; BRETTELL, L. E.; SINGH, B. Linking the Phyllosphere Microbiome to Plant Health. **Trends in Plant Science**, v. 25, n. 9, p. 841–844, 2020.
- MACHADO, I. C.; LOPES, A. V. Floral Traits and Pollination Systems in the Caatinga, a Brazilian Tropical Dry Forest. **Annals of Botany**, v. 94, n. 3, p. 365–376, 2004.
- MAESTRE, F. T. et al. Biogeography of global drylands. **New Phytologist**, v. 231, n. 2, p. 540–558. 2021.
- MAESTRE, F. T. et al. Structure and Functioning of Dryland Ecosystems in a Changing World. **Annual Review of Ecology, Evolution, and Systematics**, v. 47, p. 215–237. 2016.
- MALES, J.; GRIFFITHS, H. Functional types in the Bromeliaceae: relationships with drought-resistance traits and bioclimatic distributions. **Functional Ecology**, v. 31, n. 10, p. 1868–1880, 2017.
- MANETTI, L. M.; DEIAPORTE, R. H.; LAVERDE, A. Metabólitos secundários da família Bromeliaceae. **Química Nova**, v. 32, n. 7, p. 1885–1897, 2009.

MARENGO, J. A et al. Variabilidade e mudanças climáticas no semiárido brasileiro. **Recursos Hídricos em Regiões Semiáridas**. Instituto Nacional do Semiárido Campina Grande, p. 384–422. 2011.

MARENGO, J. A. et al. Increase risk of drought in the semiarid lands of northeast Brazil due to regional warming above 4 °C. **Climate Change Risks in Brazil**, p. 181–200, 2018.

MARENGO, J. A.; TORRES, R. R.; ALVES, L. M. Drought in Northeast Brazil—past, present, and future. **Theoretical and Applied Climatology**, v. 129, n. 3–4, p. 1189–1200. 2017.

MAUTONE, J. N. et al. Phylloplane yeasts as a source of industrially interesting enzymes. **Brazilian Journal of Biosciences**, v. 8, n. 9, p. 169–173, 2010.

MENEZES, R. S. C. et al. Biogeochemical cycling in terrestrial ecosystems of the Caatinga Biome. **Brazilian journal of biology**, v. 72, n. 3, p. 643–653, 2012.

MEYER, K. M.; LEVEAU, J. H. J. Microbiology of the phyllosphere: a playground for testing ecological concepts. **Oecologia**, v. 168, n. 3, p. 621–629. 2012.

MITTELBACH, M. et al. Nectar sugars and bird visitation define a floral niche for basidiomycetous yeast on the Canary Islands. **BMC Ecology**, v. 15, n. 1, p. 2, 2015.

MMA, Ministério do Meio Ambiente. **Monitoramento do Desmatamento nos Biomas Brasileiros por Satélite. Acordo de Cooperação Técnica MMA/IBAMA: Monitoramento do Bioma Caatinga 2008 a 2009**. Ministério do Meio Ambiente. Brasília, DF, 2011.

MOLLER, L.; LERM, B.; BOTHA, A. Interactions of arboreal yeast endophytes: An unexplored discipline. **Fungal Ecology**, 2016.

MORO, M. F. et al. A Phytogeographical Metaanalysis of the Semiarid Caatinga Domain in Brazil. **The Botanical Review**, 2016.

MORRIS, C. E. Phyllosphere. **Encyclopedia of life sciences**, p. 1–8, 2001.

NAGY, L. G. et al. Latent homology and convergent regulatory evolution underlies the repeated emergence of yeasts. **Nature communications**, v. 5, p. 4471, 2014.

NOY-MEIR, I. Desert Ecosystems: Environment and Producers. **Annual Review of Ecology and Systematics**, v. 4, n. 1, p. 25–51, 1973.

OGLE, K.; REYNOLDS, J. F. Plant responses to precipitation in desert ecosystems: Integrating functional types, pulses, thresholds, and delays. **Oecologia**, v. 141, n. 2, p. 282–294, 2004.

ORTIZ-CASTELLANOS, L.; RUIZ-HERRERA, J.. Phylogenetic relationships of the wall-synthesizing enzymes of Basidiomycota confirm the phylogeny of their subphyla. **Folia Microbiologica**, v. 60, n. 2, p. 143–150, 2015.

- OSO, S. et al. Biosurfactants Produced by Phyllosphere-Colonizing Pseudomonads Impact Diesel Degradation but Not Colonization of Leaves of Gnotobiotic *Arabidopsis thaliana*. **Applied and Environmental Microbiology**, v. 87, n. 9. 2021.
- PULLA, S. et al. Assessing the resilience of global seasonally dry tropical forests. **International Forestry Review**, v. 17, n. 2, p. 91–113. 2015.
- REISBERG, E. E. et al. Distinct phyllosphere bacterial communities on *Arabidopsis* wax mutant leaves. **PLoS ONE**, v. 8, n. 11, p. 1–12. 2013.
- REYNOLDS, J. F. et al. Modifying the ‘pulse–reserve’ paradigm for deserts of North America: precipitation pulses, soil water, and plant responses. **Oecologia**, v. 141, n. 2, p. 194–210. 2004.
- ROBERTS, I. N.; OLIVER, S. G. The yin and yang of yeast: Biodiversity research and systems biology as complementary forces driving innovation in biotechnology. **Biotechnology Letters**, v. 33, n. 3, p. 477–487. 2011.
- RODRIGUEZ, R. J. et al. Fungal endophytes: diversity and functional roles. **New Phytologist**, v. 182, n. 2, p. 314–330. 2009.
- ROSA, C. A; PÉTER, G. **Biodiversity and Ecophysiology of Yeasts**. Berlin/Heidelberg: Springer-Verlag, 2006.
- SAFRIEL, U. et al. Dryland Systems. In: **Ecosystems and Human Well-Being: Current State and Trends**. Island Press, p. 623–662. 2005.
- SAMBROOK, J. R.; RUSSEL, D. W. **Molecular Cloning: A Laboratory Manual**. 2001.
- SANTOS, J. C. et al. Caatinga: the scientific negligence experienced by a dry tropical forest. **Tropical Conservation Science**, v. 4, n. 3, p. 276–286. 2011.
- SAPKOTA, R. et al. Host genotype is an important determinant of the cereal phyllosphere mycobiome. **New Phytologist**, v. 207, n. 4, p. 1134–1144. 2015.
- SCHAEFFER, R. N. et al. Consequences of a nectar yeast for pollinator preference and performance. **Functional Ecology**, v. 31, n. 3, p. 613–621. 2017.
- SCHOCH, C. L. et al. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. **Proceedings of the National Academy of Sciences of the United States of America**, v. 109, n. 16, p. 1–6. 2012.
- SCHOCH, C. L. et al. The ascomycota tree of life: A phylum-wide phylogeny clarifies the origin and evolution of fundamental reproductive and ecological traits. **Systematic Biology**, 2009.
- SCHWINNING, S. et al. Thresholds, memory, and seasonality: understanding pulse dynamics in arid/semi-arid ecosystems. **Oecologia**, v. 141, n. 2, p. 191–193. 2004.

- SELMECKI, Anna M et al. Polyploidy can drive rapid adaptation in yeast. **Nature**, v. 519, n. 7543, p. 349–352, 2015.
- SIQUEIRA-FILHO, J. A.; LEME, E. M. C. **Fragments de Mata Atlântica do Nordeste - Biodiversidade, Conservação e suas Bromélias**. 2006.
- SPRIBILLE, T. et al. Basidiomycete yeasts in the cortex of ascomycete macrolichens. **Science**, p. 1–10, 2016.
- STARMER, W. T.; LACHANCE, M. Yeast Ecology. In: **The Yeasts**. Elsevier, 2011. p. 65–83.
- TEDERSOO, L. et al. Global diversity and geography of soil fungi. **Science**, v. 346, n. 6213, p. 1052–1053, 2014.
- THAPA, S.; PRASANNA, R. Prospecting the characteristics and significance of the phyllosphere microbiome. **Annals of Microbiology**, v. 68, n. 5, p. 229–245, 2018.
- UEDA, H. et al. Disease severity enhancement by an esterase from non-phytopathogenic yeast *Pseudozyma antarctica* and its potential as adjuvant for biocontrol agents. **Scientific Reports**, v. 8, n. 1. 2018.
- VACHER, C. et al. The Phyllosphere: Microbial Jungle at the Plant–Climate Interface. **Annual Review of Ecology, Evolution, and Systematics**, v. 47, n. 1, p. 1–24, 2016.
- VELLEND, M. Conceptual Synthesis in Community Ecology. **The Quarterly Review of Biology**, v. 85, n. 2, p. 183–206. 2010.
- VOORDECKERS, K. et al. Adaptation to High Ethanol Reveals Complex Evolutionary Pathways. **PLoS Genetics**, v. 11, n. 11, p. 1–31, 2015.
- VORHOLT, J. A. Microbial life in the phyllosphere. **Nature Reviews Microbiology**, v. 10, n. 12, p. 828–840. 2012.
- WHIPPS, J. M. et al. Phyllosphere microbiology with special reference to diversity and plant genotype. **Journal of Applied Microbiology**, v. 105, n. 6, p. 1744–1755. 2008.
- WINAGRASKI, E. et al. Diversity of arbuscular mycorrhizal fungi in forest ecosystems of Brazil: A review. **Cerne**, v. 25, n. 1, p. 25–35, 2019.
- WOLOWSKI, M.; FREITAS, L. An overview on pollination of the Neotropical Poales. **Rodriguésia**, v. 66, n. 2, p. 329–336. 2015.
- YANG, L. H. et al. What can we learn from resource pulses? **Ecology**. 2008.
- YURKOV, A. et al. Nomenclatural issues concerning cultured yeasts and other fungi: why it is important to avoid unneeded name changes. **IMA Fungus**, v. 12, n. 1. 2021.

ZEISLER-DIEHL, V. V.; BARTHLOTT, W.; SCHREIBER, L. Plant Cuticular Waxes: Composition, Function, and Interactions with Microorganisms. **Hydrocarbons, Oils and Lipids: Diversity, Origin, Chemistry and Fate**. Springer International Publishing, p. 123–138. 2020.

ZOTZ, G. et al. How much water is in the tank? An allometric analysis with 205 bromeliad species. **Flora**, v. 264. 2020.

ANEXOS

Material suplementar (Capítulo 1, secção 4)

Supplementary Table 1- New yeast species that have been described associated with bromeliads.

Phyla Subphyla	Yeast species	Host	Site	Substrate	Ref.
Ascomycota					
Saccharomycotina					
	<i>Candida aechmeae</i>	<i>Aechmea recurvata</i> and <i>Billbergia nutans</i>	Itapuã State Park, Viamão- Brazil	Phylloplane	(Landell et al., 2010)
	<i>Candida bromeliacearum</i>	<i>Canistropis seidelii</i>	Picinguaba, Serra do Mar State Park- Brazil	Phytotelma	(Ruivo et al., 2005)
	<i>Candida ubatubensis</i>	<i>C. seidelii</i>	Picinguaba, Serra do Mar State Park- Brazil	Phytotelma	(Ruivo et al., 2005)
	<i>Candida vriesiae</i>	<i>Vriesea gigantea</i>	Itapuã State Park, Viamão- Brazil	Phytotelma	(Landell et al., 2010)
	<i>Hagleromyces aurorensis</i>	<i>Bromelia karatas</i>	Aurora do Tocantins, Tocantins State- Brazil	Phytotelma	(Sousa et al., 2014)
	<i>Kazachstania bromeliacearum</i>	<i>Quesnelia quesneliana</i> , <i>Nidularium procerum</i> , <i>Neoregelia cruenta</i> , <i>Aechmea nudicaulis</i> and <i>Vriesea procera</i>	Mangrove of Coroa Grande (Poço das Antas) and Sand dune area of Maricá- Brazil	Phytotelma	(Araujo et al., 2012)
	<i>Kazachstania rupicola</i>	<i>Vriesea minarum</i>	Serra da Piedade region, Caeté city- Brazil	Phytotelma	(Safar et al., 2013)
Basidiomycota					
Agaricomycotina					
	<i>Carcinomyces nordestinensis</i>	<i>Bromelia antiacantha</i>	Tocaia reserve, Santana do Ipanema- Brazil	Phylloplane	(Crous et al., 2019)
	<i>Carlosrosaea aechmeae</i>	<i>Aechmea constantinii</i>	Murici ecological reserve, Murici- Brazil	Phylloplane	(Felix et al., 2017)
	<i>Carlosrosaea hohenbergiae</i>	<i>Hohenbergia ramageana</i> , <i>Tillandsia sp.</i>	Serra da Barriga, União dos Palmares- Brazil; Serra da Saudinha, Maceió- Brazil and Murici ecological reserve, Murici- Brazil	Phylloplane and Anthosphaera	(Felix et al., 2017)
	<i>Carlosrosaea vriesiae</i>	<i>Portea leptantha</i> , <i>Canistrum alagoanum</i> and <i>Aechmea fulgens</i>	Serra da Piedade region, Caeté city- Brazil and Itapuã State Park, Viamão- Brazil	Phylloplane and Phytotelma	(Landell et al., 2015)
	<i>Genolevuria bromelialarum</i>	<i>V. minarum</i> , <i>Vriesea fibburgensis</i> and <i>Tillandsia gardneri</i>	Itapuã State Park, Viamão- Brazil	Phylloplane	(Crestani et al., 2009)
	<i>Hannaella pagnoeae</i>	<i>V. procera</i> , <i>V. fibburgensis</i> and <i>T. gardneri</i>	Itapuã State Park, Viamão- Brazil	Phylloplane	(Landell et al., 2014)
	<i>Kockovaella libkindii</i>	<i>V. gigantea</i> , <i>Tillandsia geminiflora</i> , <i>V. minarum</i> , <i>Encholirium sp.</i> and <i>B. karatas</i>	Itapuã State Park, Viamão- Brazil; South of Brazil, Aurora do Tocantins, Tocantins State- Brazil and Serra da Piedade region, Caeté city- Brazil	Phylloplane and Phytotelma	(Landell et al., 2014)
	<i>Papiliotrema leoncinii</i>	<i>V. minarum</i>	Serra da Piedade region, Caeté city- Brazil	Phytotelma	(Gomes et al., 2016)
		<i>Tillandsia crocata</i> , <i>B. antiacantha</i> , <i>T. gardneri</i> , <i>V. gigantea</i> , <i>A. recurvata</i> , <i>V. fibburgensis</i> , <i>C. alagoanum</i> , <i>A. fulgens</i> , <i>Canistrum aurantiacum</i> , <i>P. leptantha</i> and <i>Bromelia sp.</i>	Itapuã State Park, Viamão- Brazil; Murici ecological reserve, Murici- Brazil; Tocaia reserve, Santana do Ipanema- Brazil and Pedra Talhada Biological Reserve, Quebrangulo- Brazil	Phylloplane	(Paganini et al., 2016)
	<i>Rhynchogastrema complexa</i>	<i>N. cruenta</i> and <i>Ananas comosus</i>	Hsinchu, Taiwan and Marica Resting, Rio de Janeiro-Brazil	Phylloplane	(Valente et al., 2012)
	<i>Vishniacozyma alagoana</i>	<i>Aechmea froesii</i> , <i>B. antiacantha</i> , <i>Hohenbergia stellata</i> and unidentified Bromeliaceae	Serra da Barriga, União dos Palmares- Brazil; Serra da Caiçara, Maravilha- Brazil; Tocaia Reserve, Santana do Ipanema- Brazil	Phylloplane	(Félix et al., 2020)
Pucciniomycotina					
	<i>Occultifur brasiliensis</i>	<i>Vriesea minarum</i>	Serra da Piedade region, Caeté city- Brazil	Phytotelma	(Gomes et al., 2015)
	<i>Occultifur plantarum</i>	<i>N. cruenta</i>	Coastal dune habitat, Maricá- Brazil	Phylloplane	(Khunnamwong et al., 2017)
	<i>Queiroziella brasiliensis</i>	<i>P. leptantha</i> , <i>T. geminiflora</i> and <i>V. gigantea</i>	Serra da Barriga, União dos Palmares- Brazil and Itapuã State Park, Viamão- Brazil	Phylloplane	(Crous et al., 2018)
Ustilaginomycotina					
	<i>Farysia itapuensis</i>	<i>V. fibburgensis</i> , <i>V. procera</i> , <i>T. gardneri</i> , <i>T. geminiflora</i> and <i>Dyckia sp.</i>	Itapuã State Park, Viamão- Brazil	Phylloplane	(Inácio et al., 2008)
	<i>Pattersoniomycetes tillandsiae</i>	<i>Canistrum improcerum</i> , <i>V. minarum</i> , <i>Tillandsia leiboldiana</i> , <i>Tillandsia flabellata</i>	Vera Cruz- Mexico; Vallee de Cordova- Mexico; Honduras; Guatemala; Serra da Saudinha, Maceió- Brazil and Serra do Cipó, Santana do Riacho- Brazil	Phylloplane and Phytotelma	(Piątek et al., 2017)

Supplementary Table 2- Yeast species isolated from bromeliads with some biotechnological potential registered and the paper reference.

Yeast taxon	Production of indoleacetic acid (IAA) ^a	Lipase activity ^b	Production of fermented beverages	Phosphate solubilization	Siderophore production	Antagonism to phytopathogens ^c	Promotion of plant growth ^c	Use as animal feed ^d	Air quality bioindicator	Probiotic potential	Production of extracellular enzymes				
											Amylase	Celulase	Esterase	Pectinase	Protease
															Xylanase
<i>Anomalomyces panicis</i>	(Morais et al., 2020)			(Morais et al., 2020)	(Morais et al., 2020)						(Gomes et al., 2015)	(Gomes et al., 2015)		(Gomes et al., 2015)	(Gomes et al., 2015)
<i>Aureobasidium leucospermi</i>											(Navarro et al., 2020)	(Navarro et al., 2020)	(Navarro et al., 2020)	(Navarro et al., 2020)	
<i>Aureobasidium pullulans</i>	(Morais et al., 2020)							(Brighigna et al., 2000)			(Gomes et al., 2015)	(Gomes et al., 2015)	(Gomes et al., 2015)	(Gomes et al., 2015)	(Gomes et al., 2015)
<i>Aureobasidium thailandense</i>											(Navarro et al., 2020)	(Navarro et al., 2020)	(Navarro et al., 2020)	(Navarro et al., 2020)	
<i>Candida intermedia</i>	(Morais et al., 2020)			(Morais et al., 2020)	(Morais et al., 2020)										(Gomes et al., 2015)
<i>Candida melibiosica</i>	(Morais et al., 2020)			(Morais et al., 2020)	(Morais et al., 2020)						(Gomes et al., 2015)	(Gomes et al., 2015)		(Gomes et al., 2015)	(Gomes et al., 2015)
<i>Candida membranifaciens</i>	(Morais et al., 2020)			(Morais et al., 2020)	(Morais et al., 2020)										(Gomes et al., 2015)
<i>Candida</i> spp.		(Tangsombatvichit et al., 2020)						(Brighigna et al., 2000)							(Gomes et al., 2015)
<i>Candida ubatubensis</i>	(Morais et al., 2020)			(Morais et al., 2020)	(Morais et al., 2020)										
<i>Carlosrosaea</i> sp.											(Navarro et al., 2020)	(Navarro et al., 2020)	(Navarro et al., 2020)	(Navarro et al., 2020)	
<i>Carlosrosaea vriesiae</i>	(Morais et al., 2020)			(Morais et al., 2020)	(Morais et al., 2020)		(Marques et al., 2021)								
<i>Colacogloea</i> sp.	(Morais et al., 2020)			(Morais et al., 2020)	(Morais et al., 2020)										
<i>Cystobasidium laryngis</i>															
<i>Cryptococcus</i> spp.							(Reyes et al., 2004)		(Brighigna et al., 2000)		(Gomes et al., 2015)	(Gomes et al., 2015)	(Gomes et al., 2015)	(Gomes et al., 2015)	
<i>Dioszegia</i> sp.	(Morais et al., 2020)			(Morais et al., 2020)							(Gomes et al., 2015)				(Gomes et al., 2015)
<i>Fellomyces penicillatus</i>	(Morais et al., 2020)			(Morais et al., 2020)	(Morais et al., 2020)						(Gomes et al., 2015)		(Gomes et al., 2015)	(Gomes et al., 2015)	
<i>Fellomyces</i> sp.															
<i>Hannaella pagnocciae</i>	(Morais et al., 2020)			(Morais et al., 2020)							(Gomes et al., 2015)	(Gomes et al., 2015)			(Gomes et al., 2015)
<i>Hannaella sinensis</i>															(Gomes et al., 2015)

2015																
<i>Hanseniaspora guilliermondii</i>		(Dellacassa et al., 2017)														
<i>Hanseniaspora opuntiae</i>		(Dellacassa et al., 2017)														
<i>Hanseniaspora uvarum</i>		(Chanprasartsuk et al., 2012, 2010; Dellacassa et al., 2017)														
<i>Kazachstania rupicola</i>		(Morais et al., 2020)	(Morais et al., 2020)	(Gomes et al., 2015)												
<i>Kloeckera apiculata</i> var. <i>apis</i>				(Korres et al., 2011)												
<i>Kockovaella libkindii</i>		(Morais et al., 2020)														
<i>Kodamaea ohmeri</i>		(Morais et al., 2020)	(Morais et al., 2020)	(Morais et al., 2020)	(Gomes et al., 2015)											
<i>Meira</i> sp.		(Morais et al., 2020)	(Morais et al., 2020)	(Morais et al., 2020)	(Gomes et al., 2015)											
<i>Metschnikowia koreensis</i>					(Gomes et al., 2015)											
<i>Metschnikowia koreensis</i>		(Morais et al., 2020)	(Morais et al., 2020)	(Morais et al., 2020)	(Gomes et al., 2015)											
<i>Meyerozyma caribbica</i>			(Amorim et al., 2018)		(Islam et al., 2021)	(Amorim et al., 2018)	(Gomes et al., 2015)									
<i>Meyerozyma guilliermondii</i>		(Morais et al., 2020)	(Chanprasartsuk et al., 2010; Dellacassa et al., 2017)	(Morais et al., 2020)	(Reyes et al., 2004)		(Gomes et al., 2015; Navarro et al., 2020)	(Navarro et al., 2020)	(Navarro et al., 2020)	(Gomes et al., 2015)	(Gomes et al., 2015)	(Gomes et al., 2015)				
<i>Meyerozyma</i> sp.		(Morais et al., 2020)	(Morais et al., 2020)	(Morais et al., 2020)												
<i>Moesziomyces aphidis</i>							(Gomes et al., 2015)					(Gomes et al., 2015)	(Gomes et al., 2015)			
<i>Myriangium</i> sp.		(Morais et al., 2020)	(Morais et al., 2020)				(Gomes et al., 2015)	(Gomes et al., 2015)				(Gomes et al., 2015)	(Gomes et al., 2015)			
<i>Occultifur brasiliensis</i>		(Morais et al., 2020)	(Morais et al., 2020)	(Morais et al., 2020)			(Gomes et al., 2015)	(Gomes et al., 2015)				(Gomes et al., 2015)	(Gomes et al., 2015)			
<i>Papiliotrema flavescens</i>		(Morais et al., 2020)		(Morais et al., 2020)	(Morais et al., 2020)							(Gomes et al., 2015)	(Gomes et al., 2015)			
<i>Papiliotrema laurentii</i>		(Morais et al., 2020)		(Morais et al., 2020)	(Morais et al., 2020)		(Gomes et al., 2015)	(Gomes et al., 2015)				(Gomes et al., 2015)	(Gomes et al., 2015)			
<i>Papiliotrema nemorosa</i>		(Morais et al., 2020)		(Morais et al., 2020)	(Morais et al., 2020)		(Gomes et al., 2015)	(Gomes et al., 2015)				(Gomes et al., 2015)	(Gomes et al., 2015)			
<i>Papiliotrema rajasthanensis</i>		(Morais et al., 2020)		(Morais et al., 2020)	(Morais et al., 2020)							(Gomes et al., 2015)	(Gomes et al., 2015)			
<i>Papiliotrema</i> sp.		(Morais et al., 2020)		(Morais et al., 2020)								(Gomes et al., 2015)	(Gomes et al., 2015)			

<i>Pattersoniomycetes tillandsiae</i>	(Morais et al., 2020)	(Morais et al., 2020)						
<i>Pichia fermentans</i>					(Islam et al., 2021)			
<i>Pichia kudriavzevii</i>				(Korres et al., 2011)				
<i>Pseudozyma hubeiensis</i>								(Gomes et al., 2015)
<i>Pseudozyma</i> sp.								(Gomes et al., 2015)
<i>Rhodosporidium diobovatum</i>	(Morais et al., 2020)	(Morais et al., 2020)	(Morais et al., 2020)				(Gomes et al., 2015)	(Gomes et al., 2015)
<i>Rhodotorula mucilaginosa</i>								(Gomes et al., 2015)
<i>Rhodotorula</i> spp.	(Morais et al., 2020)		(Morais et al., 2020)	(Reyes et al., 2004)				
<i>Saccharomyces cerevisiae</i>		(Nasir et al., 2017)						
<i>Saccharomyces ludwigii</i>		(Chanprasartsuk et al., 2012)						
<i>Saitozyma flava</i>						(Navarro et al., 2020)	(Navarro et al., 2020)	(Navarro et al., 2020)
<i>Saitozyma podzolica</i>	(Morais et al., 2020)		(Morais et al., 2020)			(Gomes et al., 2015)		(Gomes et al., 2015)
<i>Saturnispora silvae</i>	(Morais et al., 2020)		(Morais et al., 2020)	(Morais et al., 2020)				(Gomes et al., 2015)
<i>Sporobolomyces</i> spp.					(Brighigna et al., 2000)			
<i>Symmetrospora marina</i>						(Navarro et al., 2020)	(Navarro et al., 2020)	(Navarro et al., 2020)
<i>Tremella fuciformis</i>							(Navarro et al., 2020)	(Navarro et al., 2020)
<i>Wickerhamomyces anomalus</i>		(Dellacassa et al., 2017)						

- a- From the absence or presence of L-Tryptophan.
- b- Potential treatment of oil contamination and biodiesel production.
- c- *In vitro*.
- d- Used in fish farming.




Fig. Supple. 1- Scheme of the search, sorting and selection of papers used in the analyses.




Fig. Supple. 2- Extrapolation-rarefaction curve considering data from all bromeliad phylosphere compartments. The curve was elaborated with 1000 pseudoreplica bootstrap and the extrapolation was performed with the Chao1 estimator.

Material suplementar (Capítulo 2, secção 5)

Supplementary Table 1- Species recorded in the study, as well as their frequency of occurrence and average abundance considering the samples (individuals of bromeliads) as the analytical unit.

Phyla	Subphyla	Species	Dry	Rainy			
			Frequency (%)	Average abundance (CFU/g)	Frequency (%)		
Ascomycota							
Pezizomycotina							
		<i>Aureobasidium melanogenum</i>	3.3	125.0	3.4		
		<i>Aureobasidium thailandense</i>	4.0	251.6	55.2		
		<i>Aureobasidium tremulum</i>	3.3	12.5	3.4		
		<i>Chaetosphaeria</i> sp.			3.4		
		<i>Exophiala bergeri</i>	3.3	125.0			
		<i>Exophiala spinifera</i>	1.0	191.7	13.8		
		<i>Hortaea werneckii</i>	3.3	2625.0	106.3		
		<i>Parapyrenis conica</i>			3.4		
		<i>Pseudosydowia</i> sp.			3.4		
		<i>Selenophoma</i> sp.	6.7	125.0			
		<i>Tricellula aurantiaca</i>			3.4		
Saccharomycotina							
		<i>Candida blankii</i>	6.7	100.0	24.1		
		<i>Candida diddensiae</i>			3.4		
		<i>Candida orthopsilosis</i>	3.3	125.0			
		<i>Candida parapsilosis</i>			6.9		
		<i>Candida</i> sp. 1	3.3	125.0	125.0		
		<i>Candida</i> sp. 2	3.3	25.0	6.9		
		<i>Candida</i> sp. 3			6.9		
		<i>Candida</i> sp. 4			3.4		
		<i>Hanseniaspora opuntiae</i>			3.4		
		<i>Meyerozyma guilliermondii</i>	16.7	96.3	2.7		

	<i>Starmerella ilheusensis</i>			3.4	68.8
	<i>Yueomyces</i> sp.	3.3	12.5		
Taphrinomycotina	<i>Taphrina</i> sp.	3.3	125.0		
Basidiomycota					
Agaricomycotina					
	<i>Carcinomyces nordestinensis</i>	3.3	125.0	1.3	375.0
	<i>Carlosrosaea</i> sp. 1	3.3	125.0	3.4	125.0
	<i>Carlosrosaea</i> sp. 2	6.7	187.5	17.2	425.0
	<i>Carlosrosaea</i> sp. 3			3.4	125.0
	<i>Fellomyces penicillatus</i>			3.4	125.0
	<i>Hannaella phetchabunensis</i>			6.9	356.3
	<i>Hannaella siamensis</i>	3.3	125.0	6.9	65.6
	<i>Hannaella sinensis</i>	3.3	62.5		
	<i>Hannaella taiwanensis</i>			13.8	225.0
	<i>Hannaella zae</i>	3.3	375.0		
	<i>Kwoniella dejecticola</i>	3.3	87.5		
	<i>Kwoniella dendrophila</i>			3.4	12.5
	<i>Kwoniella heveanensis</i>	16.7	145.0		
	<i>Kwoniella mangrovensis</i>	3.3	125.0		
	<i>Papiliotrema flavescens</i>	6.7	21.9	6.9	75.0
	<i>Papiliotrema laurentii</i>	36.7	152.8	13.8	182.8
	<i>Papiliotrema miconiae</i>			6.9	81.3
	<i>Papiliotrema rajasthanensis</i>	3.3	125.0	3.4	25.0
	<i>Papiliotrema</i> sp. 1	3.3	250.0		
	<i>Papiliotrema</i> sp. 2			3.4	87.5
	<i>Rhynchogastrema noutii</i>	3.3	125.0		
	<i>Saitozyma flava</i>			6.9	43.8
	<i>Saitozyma ninhbinhensis</i>	1.0	412.5		
	<i>Saitozyma podzolica</i>	1.0	115.6	3.4	406.3
	<i>Tremella ananatis</i>	33.3	360.0	17.2	203.8
	<i>Tremella</i> sp.			3.4	50.0
	<i>Vishniacozyma alagoana</i>	16.7	197.5	6.9	125.0

Pucciniomycotina					
<i>Boekhoutia</i> sp.			3.4	125.0	
<i>Colacogloea</i> sp.	3.3	125.0			
<i>Cyrenella elegans</i>			3.4	125.0	
<i>Cystobasidium keelungensis</i>	3.3	125.0			
<i>Cystobasidium</i> sp.	3.3	125.0			
<i>Erythrobasidium</i> sp.	3.3	125.0			
<i>Halobasidium xiangyangense</i>			3.4	375.0	
<i>Hasegawazyma</i> sp.	6.7	125.0	3.4	37.5	
<i>Occultifur brasiliensis</i>	36.7	434.1	31.3	229.2	
<i>Occultifur externus</i>	3.3	125.0			
<i>Occultifur plantarum</i>	3.3	12.5	3.4	12.5	
<i>Rhodosporidiobolus ruineniae</i>	1.0	87.5			
<i>Rhodotorula mucilaginosa</i>	6.7	131.3			
<i>Rhodotorula paludigena</i>	3.3	62.5			
<i>Rosettozyma</i> sp.	6.7	125.0			
<i>Sakaguchia oryzae</i>			3.4	131.3	
<i>Symmetrospora marina</i>			6.9	56.3	
<i>Symmetrospora suhii</i>	3.3	125.0	13.8	209.4	
Ustilaginomycotina					
<i>Anthracocystis anthraeoideispora</i>			3.4	125.0	
<i>Kordyana</i> sp.			3.4	125.0	
<i>Microstroma</i> sp. 1			6.9	125.0	
<i>Microstroma</i> sp. 2	6.7	68.8			
<i>Moesziomyces antarcticus</i>	3.3	12.5			
<i>Moesziomyces aphidis</i>	3.3	12.5			
<i>Pseudozyma hubeiensis</i>			3.4	375.0	
<i>Pseudozyma pruni</i>	3.3	125.0			
<i>Pseudozyma tsukubaensis</i>	3.3	125.0			
<i>Ustilago maydis</i>			3.4	12.5	

Supplementary Table 2- Results of the similarity percentage analysis (SIMPER) between seasonal periods, the analysis was performed with 999 permutations and considering the taxonomic data (species abundance) and the twelve collections as sample units.

Species	Contribution to average dissimilarity	Standard deviation (sd)	Average to sd ratio	Average abundance Dry	Average abundance Rainy	Cumulative contribution	p-value
<i>Occultifur externus</i>	0.08443	0.08171	1.0333	159.17	68.75	0.099	0.795
<i>Candida blankii</i>	0.0831	0.148	0.5614	6.67	127.71	0.196	0.128
<i>Aureobasidium thailandense</i>	0.06777	0.08241	0.8224	100.62	46.46	0.276	0.238
<i>Tremella</i> sp.	0.05485	0.04805	1.1417	120	33.96	0.34	0.416
<i>Candida</i> sp. 3	0.05319	0.12471	0.4265	0	64.79	0.402	0.288
<i>Carlosrosaea</i> sp. 3	0.04791	0.05608	0.8544	12.5	70.83	0.458	0.334
<i>Saitozyma podzolica</i>	0.03096	0.05525	0.5603	41.25	0	0.495	0.334
<i>Kordyana</i> sp.	0.02969	0.06765	0.4389	87.5	0	0.53	0.892
<i>Microstroma</i> sp. 1	0.02893	0.04056	0.7133	16.04	46.67	0.564	0.791
<i>Papiliotrema miconiae</i>	0.02528	0.0184	1.3737	56.04	24.38	0.593	0.499
<i>Yueomyces</i> sp.	0.02312	0.0235	0.9839	32.92	8.33	0.62	0.202
<i>Carlosrosaea</i> sp. 1	0.02127	0.03606	0.5899	4.17	37.5	0.645	0.782
<i>Hannaella zaeae</i>	0.021	0.02641	0.7952	0	30	0.67	0.068
<i>Kwoniella mangrovensis</i>	0.0193	0.03404	0.5671	24.17	0	0.692	0.209
<i>Fellomyces penicillatus</i>	0.01751	0.0223	0.7854	19.17	14.17	0.713	0.63
<i>Hannaella siamensis</i>	0.01724	0.04104	0.4202	0	23.75	0.733	0.396
<i>Taphrina</i> sp.	0.01634	0.02165	0.7549	4.17	27.92	0.752	0.338
<i>Hannaella phetchabunensis</i>	0.01126	0.02725	0.4132	0	12.5	0.765	0.262
<i>Sakaguchia oryzae</i>	0.0111	0.01851	0.5995	7.4	13.54	0.778	0.6
<i>Hanseniaspora opuntiae</i>	0.00921	0.0213	0.4324	12.5	0	0.789	0.577
<i>Candida diddensiae</i>	0.00751	0.01816	0.4132	0	8.33	0.798	0.262
<i>Colacogloea</i> sp.	0.00751	0.01816	0.4132	0	8.33	0.807	0.262
<i>Rhodotorula mucilaginosa</i>	0.0066	0.01461	0.4518	8.75	0	0.815	0.527
<i>Saitozyma flava</i>	0.00653	0.00963	0.6781	8.33	0	0.822	0.235

<i>Papiliotrema</i> sp. 2	0.00635	0.01471	0.4319	8.33	0	0.83	0.561
<i>Pseudozyma pruni</i>	0.00616	0.01436	0.4288	0	12.5	0.837	0.56
<i>Carlosrosaea</i> sp. 2	0.00582	0.01026	0.5673	4.17	4.17	0.844	0.699
<i>Microstroma</i> sp. 2	0.00563	0.00856	0.6573	0	8.33	0.85	0.217
<i>Hannaella sinensis</i>	0.00535	0.00851	0.6292	4.17	4.38	0.857	0.77
<i>Candida parapsilosis</i>	0.0052	0.01227	0.424	0	8.33	0.863	0.466
<i>Hortaea werneckii</i>	0.00503	0.00754	0.668	8.33	1.25	0.869	0.423
<i>Papiliotrema laurentii</i>	0.00398	0.00518	0.7695	1.46	5	0.878	0.273
<i>Moesziomyces antarcticus</i>	0.00369	0.00856	0.431	4.58	0	0.883	0.586
<i>Papiliotrema rajasthanensis</i>	0.00366	0.00868	0.4221	0	5.42	0.887	0.402
<i>Rhodotorula paludigena</i>	0.00347	0.00577	0.6002	8.75	0	0.891	0.375
<i>Candida</i> sp. 4	0.0034	0.0081	0.4202	0	4.69	0.895	0.396
<i>Candida</i> sp. 1	0.00335	0.00778	0.431	4.17	0	0.899	0.586
<i>Cystobasidium</i> sp.	0.00335	0.00778	0.431	4.17	0	0.903	0.586
<i>Occultifur plantarum</i>	0.00335	0.00778	0.431	4.17	0	0.907	0.586
<i>Rhodosporidiobolus ruineniae</i>	0.00335	0.00778	0.431	4.17	0	0.911	0.586
<i>Tremella ananatis</i>	0.00335	0.00778	0.431	4.17	0	0.915	0.586
<i>Rosettozyma</i> sp.	0.00318	0.00736	0.4319	4.17	0	0.918	0.561
<i>Candida orthopsilosis</i>	0.00307	0.0071	0.4324	4.17	0	0.922	0.577
<i>Boekhoutia</i> sp.	0.00303	0.0072	0.4202	0	4.17	0.925	0.396
<i>Chaetosphaeria</i> sp.	0.00303	0.0072	0.4202	0	4.17	0.929	0.396
<i>Kwoniella dejecticola</i>	0.00303	0.0072	0.4202	0	4.17	0.932	0.396
<i>Ustilago maydis</i>	0.00303	0.0072	0.4202	0	4.17	0.936	0.396
<i>Candida</i> sp. 2	0.00299	0.00602	0.4976	0.83	2.92	0.94	0.451
<i>Papiliotrema</i> sp. 1	0.00298	0.00544	0.5485	4.17	0.83	0.943	0.764
<i>Starmerella ilheusensis</i>	0.00283	0.00644	0.4389	8.33	0	0.946	0.887
<i>Selenophoma</i> sp.	0.00273	0.00644	0.424	0	4.38	0.95	0.466
<i>Pseudosydowia</i> sp.	0.0026	0.00614	0.424	0	4.17	0.953	0.466
<i>Cyrenella elegans</i>	0.00248	0.00569	0.4351	4.17	0	0.956	0.652
<i>Erythrobasidium</i> sp.	0.00248	0.00569	0.4351	4.17	0	0.959	0.652
<i>Exophiala bergeri</i>	0.00248	0.00569	0.4351	4.17	0	0.961	0.652
<i>Parapyrenis conica</i>	0.00243	0.00584	0.416	0	2.92	0.964	0.306
<i>Symmetrospora suhii</i>	0.00243	0.00483	0.503	0	3.75	0.967	0.268

<i>Aureobasidium tremulum</i>	0.00239	0.00459	0.5196	0.42	4.17	0.97	0.65
<i>Anthracocystis anthracoidespora</i>	0.00205	0.00479	0.4288	0	4.17	0.972	0.56
<i>Cystobasidium keelungensis</i>	0.00205	0.00479	0.4288	0	4.17	0.975	0.56
<i>Halobasidium xiangyangense</i>	0.00205	0.00479	0.4288	0	4.17	0.977	0.56
<i>Hasegawazyma</i> sp.	0.00205	0.00479	0.4288	0	4.17	0.98	0.56
<i>Saitozyma ninhbinhensis</i>	0.00182	0.0043	0.424	0	2.92	0.982	0.466
<i>Kwoniella dendrophila</i>	0.00173	0.00399	0.4351	2.92	0	0.984	0.652
<i>Aureobasidium melanogenum</i>	0.00163	0.00313	0.521	4.17	0.42	0.986	0.903
<i>Pseudozyma hubeiensis</i>	0.00156	0.00368	0.424	0	2.5	0.987	0.466
<i>Symmetrospora marina</i>	0.00143	0.00338	0.424	0	2.29	0.989	0.466
<i>Exophiala spinifera</i>	0.00141	0.00322	0.4389	4.17	0	0.991	0.887
<i>Meyerozyma guilliermondii</i>	0.00141	0.00322	0.4389	4.17	0	0.992	0.887
<i>Pseudozyma tsukubaensis</i>	0.00141	0.00322	0.4389	4.17	0	0.994	0.887
<i>Tricellula aurantiaca</i>	0.00139	0.00334	0.416	0	1.67	0.996	0.306
<i>Hannaella taiwanensis</i>	0.00124	0.00285	0.4351	2.08	0	0.997	0.652
<i>Rhynchogastrema noutii</i>	0.00071	0.00161	0.4389	2.08	0	0.998	0.887
<i>Papiliotrema flavescens</i>	0.00054	0.00094	0.5766	0.42	0.42	0.999	0.584
<i>Kwoniella heveanensis</i>	0.00035	0.00083	0.416	0	0.42	0.999	0.306
<i>Occultifur brasiliensis</i>	0.00034	0.00078	0.431	0.42	0	0.999	0.586
<i>Vishniacozyma alagoana</i>	0.00026	0.00061	0.424	0	0.42	1	0.466
<i>Moesziomyces aphidis</i>	0.00014	0.00032	0.4389	0.42	0	1	0.887
<i>Yueomyces</i> sp.	0.00014	0.00032	0.4389	0.42	0	1	0.887

Supplementary Table 3- Results of the similarity percentage analysis (SIMPER) between seasonal periods, the analysis was performed with 999 permutations and considering the phylogenetic data (taxa abundance) and the twelve collections as sample units.

Taxon	Contribution to average dissimilarity	Standard deviation (sd)	Average to sd ratio	Average abundance Dry	Average abundance Rainy	Cumulative contribution	p-value
Basidiomycota	3.85E-02	2.29E-02	1.68E+00	5.85E+02	4.35E+02	0.064	0.673
Ascomycota	3.27E-02	2.39E-02	1.37E+00	2.61E+02	3.54E+02	0.118	0.191
Saccharomycetales	2.72E-02	2.49E-02	1.09E+00	3.23E+01	2.70E+02	0.163	0.013
Saccharomycetes	2.72E-02	2.49E-02	1.09E+00	3.23E+01	2.70E+02	0.208	0.013
Saccharomycotina	2.72E-02	2.49E-02	1.09E+00	3.23E+01	2.70E+02	0.254	0.013
<i>Candida</i>	2.46E-02	2.67E-02	9.20E-01	1.58E+01	2.17E+02	0.294	0.017
Tremellales	1.99E-02	1.45E-02	1.37E+00	3.51E+02	2.78E+02	0.368	0.758
Tremellomycetes	1.99E-02	1.45E-02	1.37E+00	3.51E+02	2.78E+02	0.401	0.758
Agaricomycotina	1.99E-02	1.45E-02	1.37E+00	3.51E+02	2.78E+02	0.434	0.758
Pucciniomycotina	1.94E-02	1.46E-02	1.33E+00	2.21E+02	1.27E+02	0.466	0.866
Pezizomycotina	1.90E-02	1.63E-02	1.17E+00	2.24E+02	8.44E+01	0.498	0.356
Cystobasidiomycetes	1.77E-02	1.42E-02	1.25E+00	1.89E+02	1.23E+02	0.527	0.947
Dothideomycetes	1.61E-02	1.46E-02	1.10E+00	2.01E+02	5.35E+01	0.554	0.299
Cystobasidiaceae	1.50E-02	1.29E-02	1.17E+00	1.72E+02	8.17E+01	0.579	0.904
Cystobasidiales	1.50E-02	1.29E-02	1.17E+00	1.72E+02	8.17E+01	0.603	0.904
<i>Occultifur</i>	1.46E-02	1.37E-02	1.07E+00	1.64E+02	6.92E+01	0.628	0.813
Trimorphomycetaceae	1.26E-02	1.16E-02	1.09E+00	6.95E+01	9.56E+01	0.648	0.802
Dothideales	1.12E-02	1.36E-02	8.18E-01	1.14E+02	5.35E+01	0.667	0.356
<i>Aureobasidium</i>	1.11E-02	1.37E-02	8.10E-01	1.05E+02	5.10E+01	0.685	0.325
Dothioraceae	1.11E-02	1.37E-02	8.09E-01	1.14E+02	5.10E+01	0.704	0.343
<i>Carlosrosaea</i>	9.26E-03	9.92E-03	9.34E-01	1.67E+01	7.92E+01	0.719	0.381
<i>Tremella</i>	8.91E-03	8.02E-03	1.11E+00	1.20E+02	3.56E+01	0.734	0.468
Tremellaceae	8.91E-03	8.02E-03	1.11E+00	1.20E+02	3.56E+01	0.749	0.468

<i>Hannaella</i>	7.28E-03	1.08E-02	6.76E-01	1.88E+01	5.81E+01	0.776	0.58
<i>Saitozyma</i>	7.01E-03	9.18E-03	7.64E-01	5.28E+01	1.65E+01	0.787	0.396
<i>Papiliotrema</i>	6.13E-03	4.30E-03	1.43E+00	7.00E+01	3.85E+01	0.797	0.647
<i>Hortaea</i>	4.95E-03	1.13E-02	4.39E-01	8.75E+01	0.00E+00	0.806	0.88
Teratosphaeriaceae	4.95E-03	1.13E-02	4.39E-01	8.75E+01	0.00E+00	0.814	0.88
Capnodiales	4.95E-03	1.13E-02	4.39E-01	8.75E+01	0.00E+00	0.822	0.88
<i>Meyerozyma</i>	4.82E-03	6.76E-03	7.13E-01	1.60E+01	4.67E+01	0.83	0.775
Debaryomycetaceae	4.82E-03	6.76E-03	7.13E-01	1.60E+01	4.67E+01	0.838	0.775
<i>Vishniacozyma</i>	3.85E-03	3.92E-03	9.84E-01	3.29E+01	8.30E+00	0.845	0.22
Bulleribasidiaceae	3.85E-03	3.92E-03	9.84E-01	3.29E+01	8.30E+00	0.851	0.22
<i>Kwoniella</i>	3.73E-03	5.38E-03	6.93E-01	3.12E+01	4.00E-01	0.857	0.054
<i>Carcinomyces</i>	3.55E-03	6.01E-03	5.90E-01	4.20E+00	3.75E+01	0.863	0.77
Carcinomycetaceae	3.55E-03	6.01E-03	5.90E-01	4.20E+00	3.75E+01	0.869	0.77
Microbotryomycetes	3.30E-03	3.20E-03	1.03E+00	3.21E+01	0.00E+00	0.874	0.056
Ustilaginomycotina	3.26E-03	3.47E-03	9.38E-01	1.38E+01	2.96E+01	0.88	0.671
<i>Symmetrospora</i>	3.11E-03	4.07E-03	7.65E-01	4.20E+00	3.17E+01	0.885	0.289
Symmetrosporaceae	3.11E-03	4.07E-03	7.65E-01	4.20E+00	3.17E+01	0.89	0.289
<i>Exophiala</i>	2.96E-03	3.68E-03	8.03E-01	2.33E+01	1.42E+01	0.9	0.768
Herpotrichiellaceae	2.96E-03	3.68E-03	8.03E-01	2.33E+01	1.42E+01	0.905	0.768
Chaetothyriales	2.96E-03	3.68E-03	8.03E-01	2.33E+01	1.42E+01	0.91	0.768
Eurotiomycetes	2.96E-03	3.68E-03	8.03E-01	2.33E+01	1.42E+01	0.915	0.768
Ustilaginaceae	2.05E-03	2.98E-03	6.87E-01	9.20E+00	1.71E+01	0.918	0.848
Ustilaginales	2.05E-03	2.98E-03	6.87E-01	9.20E+00	1.71E+01	0.922	0.848
Ustilaginomycetes	2.05E-03	2.98E-03	6.87E-01	9.20E+00	1.71E+01	0.925	0.848
<i>Halobasidium</i>	1.88E-03	4.54E-03	4.13E-01	0.00E+00	1.25E+01	0.928	0.258
Sporidiobolaceae	1.80E-03	2.42E-03	7.43E-01	1.96E+01	0.00E+00	0.931	0.272
Sporidiobolales	1.80E-03	2.42E-03	7.43E-01	1.96E+01	0.00E+00	0.934	0.272
Sordariomycetes	1.68E-03	3.02E-03	5.59E-01	0.00E+00	1.25E+01	0.937	0.126
Exobasidiomycetes	1.68E-03	2.35E-03	7.16E-01	4.60E+00	1.25E+01	0.94	0.697
<i>Pseudozyma</i>	1.63E-03	2.30E-03	7.07E-01	8.30E+00	1.25E+01	0.942	0.824

<i>Chaetosphaeria</i>	1.25E-03	3.03E-03	4.13E-01	0.00E+00	8.30E+00	0.944	0.258
<i>Chaetosphaeriaceae</i>	1.25E-03	3.03E-03	4.13E-01	0.00E+00	8.30E+00	0.947	0.258
<i>Chaetosphaeriales</i>	1.25E-03	3.03E-03	4.13E-01	0.00E+00	8.30E+00	0.949	0.258
<i>Erythrobasidiales</i>	1.22E-03	1.34E-03	9.14E-01	1.25E+01	1.20E+00	0.951	0.215
<i>Microstroma</i>	1.20E-03	1.61E-03	7.43E-01	4.60E+00	8.30E+00	0.953	0.696
<i>Microstromataceae</i>	1.20E-03	1.61E-03	7.43E-01	4.60E+00	8.30E+00	0.955	0.696
<i>Microstromatales</i>	1.20E-03	1.61E-03	7.43E-01	4.60E+00	8.30E+00	0.957	0.696
<i>Rhodosporidiobolus</i>	1.10E-03	2.44E-03	4.52E-01	8.80E+00	0.00E+00	0.958	0.561
<i>Rosettozyma</i>	1.09E-03	1.61E-03	6.78E-01	8.30E+00	0.00E+00	0.96	0.249
<i>Rosettozymaceae</i>	1.09E-03	1.61E-03	6.78E-01	8.30E+00	0.00E+00	0.962	0.249
<i>Rosettozymales</i>	1.09E-03	1.61E-03	6.78E-01	8.30E+00	0.00E+00	0.964	0.249
<i>Cystobasidium</i>	9.70E-04	1.45E-03	6.69E-01	8.30E+00	0.00E+00	0.965	0.322
<i>Hasegawazyma</i>	8.40E-04	1.26E-03	6.68E-01	8.30E+00	1.20E+00	0.967	0.427
<i>Rhodotorula</i>	7.00E-04	1.22E-03	5.72E-01	1.08E+01	0.00E+00	0.969	0.41
<i>Saccharomycetaceae</i>	5.90E-04	8.73E-04	6.75E-01	4.00E-01	6.50E+00	0.97	0.249
<i>Taphrina</i>	5.60E-04	1.30E-03	4.31E-01	4.20E+00	0.00E+00	0.971	0.6
<i>Taphrinaceae</i>	5.60E-04	1.30E-03	4.31E-01	4.20E+00	0.00E+00	0.972	0.6
<i>Taphrinales</i>	5.60E-04	1.30E-03	4.31E-01	4.20E+00	0.00E+00	0.973	0.6
<i>Taphrinomycetes</i>	5.60E-04	1.30E-03	4.31E-01	4.20E+00	0.00E+00	0.974	0.6
<i>Taphrinomycotina</i>	5.60E-04	1.30E-03	4.31E-01	4.20E+00	0.00E+00	0.975	0.6
<i>Rhynchogastrema</i>	5.30E-04	1.23E-03	4.32E-01	4.20E+00	0.00E+00	0.976	0.593
<i>Rhynchogastremaceae</i>	5.30E-04	1.23E-03	4.32E-01	4.20E+00	0.00E+00	0.977	0.593
<i>Boekhoutia</i>	5.00E-04	1.20E-03	4.20E-01	0.00E+00	4.20E+00	0.978	0.382
<i>Kordyana</i>	5.00E-04	1.20E-03	4.20E-01	0.00E+00	4.20E+00	0.978	0.382
<i>Tricellula</i>	5.00E-04	1.20E-03	4.20E-01	0.00E+00	4.20E+00	0.979	0.382
<i>Brachybasidiaceae</i>	5.00E-04	1.20E-03	4.20E-01	0.00E+00	4.20E+00	0.98	0.382
<i>Calloriaceae</i>	5.00E-04	1.20E-03	4.20E-01	0.00E+00	4.20E+00	0.981	0.382
<i>Chionosphaeraceae</i>	5.00E-04	1.20E-03	4.20E-01	0.00E+00	4.20E+00	0.982	0.382
<i>Agaricostilbales</i>	5.00E-04	1.20E-03	4.20E-01	0.00E+00	4.20E+00	0.983	0.382
<i>Exobasidiales</i>	5.00E-04	1.20E-03	4.20E-01	0.00E+00	4.20E+00	0.983	0.382

Helotiales	5.00E-04	1.20E-03	4.20E-01	0.00E+00	4.20E+00	0.984	0.382
Agaricostilbomycetes	5.00E-04	1.20E-03	4.20E-01	0.00E+00	4.20E+00	0.985	0.382
Leotiomycetes	5.00E-04	1.20E-03	4.20E-01	0.00E+00	4.20E+00	0.986	0.382
<i>Selenophoma</i>	4.70E-04	1.07E-03	4.39E-01	8.30E+00	0.00E+00	0.987	0.88
<i>Sakaguchia</i>	4.60E-04	1.07E-03	4.24E-01	0.00E+00	4.40E+00	0.987	0.437
Sakaguchiaceae	4.60E-04	1.07E-03	4.24E-01	0.00E+00	4.40E+00	0.988	0.437
Sakaguchiales	4.60E-04	1.07E-03	4.24E-01	0.00E+00	4.40E+00	0.989	0.437
<i>Parapyrenis</i>	4.30E-04	1.02E-03	4.24E-01	0.00E+00	4.20E+00	0.99	0.437
Requienellaceae	4.30E-04	1.02E-03	4.24E-01	0.00E+00	4.20E+00	0.99	0.437
Xylariales	4.30E-04	1.02E-03	4.24E-01	0.00E+00	4.20E+00	0.991	0.437
<i>Colacogloea</i>	4.10E-04	9.49E-04	4.35E-01	4.20E+00	0.00E+00	0.992	0.672
<i>Erythrobasidium</i>	4.10E-04	9.49E-04	4.35E-01	4.20E+00	0.00E+00	0.993	0.672
Erythrobasiidaeae	4.10E-04	9.49E-04	4.35E-01	4.20E+00	0.00E+00	0.993	0.672
Heterogastridiaceae	4.10E-04	9.49E-04	4.35E-01	4.20E+00	0.00E+00	0.994	0.672
Heterogastridiales	4.10E-04	9.49E-04	4.35E-01	4.20E+00	0.00E+00	0.995	0.672
<i>Anthracocystis</i>	3.40E-04	7.98E-04	4.29E-01	0.00E+00	4.20E+00	0.995	0.538
<i>Cyrenella</i>	3.40E-04	7.98E-04	4.29E-01	0.00E+00	4.20E+00	0.996	0.538
<i>Fellomyces</i>	3.40E-04	7.98E-04	4.29E-01	0.00E+00	4.20E+00	0.996	0.538
<i>Hanseniaspora</i>	3.40E-04	7.98E-04	4.29E-01	0.00E+00	4.20E+00	0.997	0.538
Cuniculitremaceae	3.40E-04	7.98E-04	4.29E-01	0.00E+00	4.20E+00	0.997	0.538
<i>Pseudosydowia</i>	2.60E-04	6.14E-04	4.24E-01	0.00E+00	2.50E+00	0.999	0.437
Saccotheciaceae	2.60E-04	6.14E-04	4.24E-01	0.00E+00	2.50E+00	0.999	0.437
<i>Starmerella</i>	2.40E-04	5.63E-04	4.24E-01	0.00E+00	2.30E+00	1	0.437
<i>Moesziomyces</i>	8.00E-05	1.30E-04	6.10E-01	8.00E-01	0.00E+00	1	0.403
<i>Ustilago</i>	4.00E-05	1.02E-04	4.24E-01	0.00E+00	4.00E-01	1	0.436
<i>Yueomyces</i>	2.00E-05	5.40E-05	4.39E-01	4.00E-01	0.00E+00	1	0.877

Supplementary Table 4- Results of the similarity percentage analysis (SIMPER) between seasonal periods, the analysis was performed with 999 permutations and considering the functional data (traits frequency) and the twelve collections as sample units.

Traits	Contribution to average dissimilarity	Standard deviation (sd)	Average to sd ratio	Average abundance Dry	Average abundance Rainy	Ordered cumulative contribution	p-value
Ramnose	0.018078	0.014416	1.254	0.605	0.8217	0.086	0.169
Galactose	0.016903	0.0138	1.2248	0.6817	0.9133	0.167	0.038
Xylose	0.016656	0.015116	1.1019	0.7	0.8983	0.247	0.173
Cellobiose	0.016389	0.011915	1.3755	0.6533	0.8167	0.325	0.272
Glucose	0.016285	0.012515	1.3012	0.695	0.9283	0.403	0.058
Inulina	0.015743	0.013409	1.174	0.725	0.9083	0.478	0.162
Glycerol	0.015555	0.011719	1.3273	0.6883	0.8833	0.553	0.104
D_arabinose	0.014497	0.012264	1.182	0.72	0.845	0.622	0.291
Raffinose	0.014035	0.012752	1.1006	0.7417	0.915	0.689	0.154
Esterase	0.011318	0.007612	1.4868	0.25	0.3583	0.743	0.351
Celullase	0.010855	0.005879	1.8464	0.0517	0.21	0.795	0.024
Lipase	0.009775	0.00649	1.5061	0.2333	0.2317	0.842	0.997
Fermentation	0.008649	0.006469	1.3369	0.2317	0.29	0.883	0.558
Caseinase	0.007192	0.00496	1.4499	0.2533	0.2233	0.918	0.255
IE ₂₄	0.006061	0.004289	1.4131	0.2983	0.235	0.947	0.312
Amylase	0.00603	0.004363	1.3821	0.07	0.1267	0.975	0.413
Pectinase	0.005139	0.007482	0.6868	0.095	0	1	0.408

Supplementary Table 5- Synthesis of information on environmental variables and diversity metrics in the twelve collections carried out in the study.

Collection ID	Collection date	Period	Storm volume	Rain volume	Rain memory	Drought period	Rainy days in the moth	Rain volume in the moth	Average yeast abundance	q0	q1	q2	AvTD	FDis	FRic	FEve
			(mm)	(mm)	(days)	(days)		(mm)	(CFU/g)							
D01	October 7, 2020	Dry	12.2	2.1	1	27	2	5.6	510	3.8	3.21	3.05	76.00	3.18	1.52	0.44
D02	November 10, 2020	Dry	62.7	13	0	4	3	62.7	623.75	4	3.13	2.73	53.95	2.62	4.72	0.60
D03	December 2, 2020	Dry	62.7	6.8	4	21	4	69.5	2412.5	5.4	3.08	2.51	84.89	1.44	9.49	0.55
D04	October 19, 2021	Dry	9.5	13.4	25	23	1	13.4	396.25	3	2.38	2.13	59.57	1.55	1.03	0.48
D05	November 23, 2021	Dry	9.5	13.4	25	58	0	0	1433.75	4.6	3.46	2.98	84.69	2.35	2.16	0.66
D06	December 7, 2021	Dry	9.5	45	63	8	1	45	678.125	2.8	2.02	1.72	76.55	1.45	0.39	0.38
R01	May 18, 2021	Rainy	79.7	0.6	0	1	16	136.2	1318.75	6	4.83	4.21	82.78	2.45	6.01	0.72
R02	June 15, 2021	Rainy	46.2	12.3	1	8	12	72.8	545	3.6	2.47	2.12	66.94	1.46	2.63	0.30
R03	July 20, 2021	Rainy	11	2.3	1	1	15	124.4	800	1.8	1.88	1.62	40.19	1.17	0.46	0.17
R04	May 24, 2022	Rainy	36.2	30.9	9	2	7	99.9	1116.875	5.4	3.95	3.28	87.89	1.88	1.62	0.58
R05	June 23, 2022	Rainy	18.7	7.8	0	1	23	304.6	996.875	4	3.25	2.88	85.61	2.37	1.30	0.65
R06	July 19, 2022	Rainy	18.2	4.3	1	2	22	228.4	611.25	2.6	2.28	2.13	67.36	1.77	0.70	0.32

Supplementary Table 6- List of isolates observed in the study.

ID	Species	Class	Subphyla	Phyla	Collection date	Seasonal period	Abundance (CFU/cm ²)	GenBank number	Collections number
BRT461	<i>Anthracocystis anthraeoideispora</i>	Ustilaginomycetes	Ustilaginomycotina	Basidiomycota	18/05/2021	Chuvoso	125		
BRT632	<i>Aureobasidium melanogenum</i>	Dothideomycetes	Pezizomycotina	Ascomycota	24/05/2022	Chuvoso	12.5		
BRT403	<i>Aureobasidium melanogenum</i>	Dothideomycetes	Pezizomycotina	Ascomycota	02/12/2020	Seco	125		
BRT616	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	07/12/2021	Seco	12.5		
BRT624	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	24/05/2022	Chuvoso	12.5		
BRT700	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	19/07/2022	Chuvoso	12.5		
BRT542	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	50		
BRT609	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	07/12/2021	Seco	62.5		
BRT524	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	19/10/2021	Seco	75		
BRT538	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	137.5		
BRT633	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	24/05/2022	Chuvoso	212.5		
BRT618	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	07/12/2021	Seco	12.5		
BRT392	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	10/11/2020	Seco	12.5		
BRT440	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	18/05/2021	Chuvoso	12.5		
BRT472	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	15/06/2021	Chuvoso	12.5		
BRT474	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	15/06/2021	Chuvoso	12.5		
BRT482	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	15/06/2021	Chuvoso	12.5		
BRT489	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	15/06/2021	Chuvoso	12.5		
BRT491	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	15/06/2021	Chuvoso	12.5		
BRT494	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	15/06/2021	Chuvoso	12.5		
BRT516	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	19/10/2021	Seco	12.5		
BRT525	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	19/10/2021	Seco	12.5		
BRT594	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	07/12/2021	Seco	12.5		
BRT600	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	07/12/2021	Seco	12.5		
BRT676	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/06/2022	Chuvoso	12.5		
BRT623	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	24/05/2022	Chuvoso	18.75		
BRT454	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	18/05/2021	Chuvoso	25		
BRT518	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	19/10/2021	Seco	25		
BRT610	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	07/12/2021	Seco	25		
BRT475	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	15/06/2021	Chuvoso	31.25		
BRT541	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	37.5		
BRT557	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	37.5		
BRT590	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	07/12/2021	Seco	37.5		
BRT445	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	18/05/2021	Chuvoso	50		

BRT503	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	20/07/2021	Chuvoso	62.5
BRT552	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	62.5
BRT602	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	07/12/2021	Seco	68.75
BRT545	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	75
BRT480	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	15/06/2021	Chuvoso	87.5
BRT462	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	18/05/2021	Chuvoso	125
BRT510	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	20/07/2021	Chuvoso	125
BRT514	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	20/07/2021	Chuvoso	125
BRT547	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	125
BRT548	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	125
BRT550	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	125
BRT560	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	125
BRT568	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	125
BRT595	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	07/12/2021	Seco	125
BRT707	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	19/07/2022	Chuvoso	125
BRT705	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	19/07/2022	Chuvoso	131.25
BRT669	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/06/2022	Chuvoso	150
BRT562	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	250
BRT569	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	250
BRT576	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	250
BRT597	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	07/12/2021	Seco	250
BRT579	<i>Aureobasidium thailandense</i>	Dothideomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	500
BRT520	<i>Aureobasidium tremulum</i>	Dothideomycetes	Pezizomycotina	Ascomycota	19/10/2021	Seco	12.5
BRT456	<i>Aureobasidium tremulum</i>	Dothideomycetes	Pezizomycotina	Ascomycota	18/05/2021	Chuvoso	125
BRT698	<i>Boekhoutia</i> sp.	Agaricostilbomycetes	Pucciniomycotina	Basidiomycota	23/06/2022	Chuvoso	125
BRT399	<i>Candida blankii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	02/12/2020	Seco	75
BRT505	<i>Candida blankii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	20/07/2021	Chuvoso	12.5
BRT506	<i>Candida blankii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	20/07/2021	Chuvoso	12.5
BRT507	<i>Candida blankii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	20/07/2021	Chuvoso	12.5
BRT493	<i>Candida blankii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	15/06/2021	Chuvoso	62.5
BRT673	<i>Candida blankii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	23/06/2022	Chuvoso	81.25
BRT467	<i>Candida blankii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	18/05/2021	Chuvoso	125
BRT513	<i>Candida blankii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	20/07/2021	Chuvoso	125
BRT589	<i>Candida blankii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	07/12/2021	Seco	125
BRT701	<i>Candida blankii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	19/07/2022	Chuvoso	125
BRT499	<i>Candida blankii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	20/07/2021	Chuvoso	2750
BRT706	<i>Candida diddensiae</i>	Saccharomycetes	Saccharomycotina	Ascomycota	19/07/2022	Chuvoso	250
BRT596	<i>Candida orthopsilosis</i>	Saccharomycetes	Saccharomycotina	Ascomycota	07/12/2021	Seco	125

BRT664	<i>Candida parapsilosis</i>	Saccharomycetes	Saccharomycotina	Ascomycota	24/05/2022	Chuviso	125
BRT662	<i>Candida parapsilosis</i>	Saccharomycetes	Saccharomycotina	Ascomycota	24/05/2022	Chuviso	125
BRT360	<i>Candida</i> sp. 1	Saccharomycetes	Saccharomycotina	Ascomycota	07/10/2020	Seco	125
BRT708	<i>Candida</i> sp. 2	Saccharomycetes	Saccharomycotina	Ascomycota	19/07/2022	Chuviso	12.5
BRT374	<i>Candida</i> sp. 2	Saccharomycetes	Saccharomycotina	Ascomycota	10/11/2020	Seco	25
BRT702	<i>Candida</i> sp. 2	Saccharomycetes	Saccharomycotina	Ascomycota	19/07/2022	Chuviso	75
BRT435	<i>Candida</i> sp. 3	Saccharomycetes	Saccharomycotina	Ascomycota	18/05/2021	Chuviso	68.75
BRT473	<i>Candida</i> sp. 3	Saccharomycetes	Saccharomycotina	Ascomycota	15/06/2021	Chuviso	1875
BRT674	<i>Candida</i> sp. 4	Saccharomycetes	Saccharomycotina	Ascomycota	23/06/2022	Chuviso	140.625
BRT508	<i>Carcinomyces nordestinensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	20/07/2021	Chuviso	125
BRT535	<i>Carcinomyces nordestinensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	19/10/2021	Seco	125
BRT448	<i>Carcinomyces nordestinensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	18/05/2021	Chuviso	500
BRT457	<i>Carcinomyces nordestinensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	18/05/2021	Chuviso	500
BRT536	<i>Carlosrosaea</i> sp. 1	Tremellomycetes	Agaricomycotina	Basidiomycota	19/10/2021	Seco	125
BRT697	<i>Carlosrosaea</i> sp. 1	Tremellomycetes	Agaricomycotina	Basidiomycota	23/06/2022	Chuviso	125
BRT586	<i>Carlosrosaea</i> sp. 2	Tremellomycetes	Agaricomycotina	Basidiomycota	23/11/2021	Seco	125
BRT537	<i>Carlosrosaea</i> sp. 2	Tremellomycetes	Agaricomycotina	Basidiomycota	19/10/2021	Seco	250
BRT663	<i>Carlosrosaea</i> sp. 2	Tremellomycetes	Agaricomycotina	Basidiomycota	24/05/2022	Chuviso	250
BRT714	<i>Carlosrosaea</i> sp. 2	Tremellomycetes	Agaricomycotina	Basidiomycota	19/07/2022	Chuviso	250
BRT659	<i>Carlosrosaea</i> sp. 2	Tremellomycetes	Agaricomycotina	Basidiomycota	24/05/2022	Chuviso	375
BRT651	<i>Carlosrosaea</i> sp. 2	Tremellomycetes	Agaricomycotina	Basidiomycota	24/05/2022	Chuviso	625
BRT696	<i>Carlosrosaea</i> sp. 2	Tremellomycetes	Agaricomycotina	Basidiomycota	23/06/2022	Chuviso	625
BRT692	<i>Carlosrosaea</i> sp. 3	Tremellomycetes	Agaricomycotina	Basidiomycota	23/06/2022	Chuviso	125
BRT710	<i>Chaetosphaeria</i> sp.	Sordariomycetes	Pezizomycotina	Ascomycota	19/07/2022	Chuviso	250
BRT587	<i>Colacogloea</i> sp.	Microbotryomycetes	Pucciniomycotina	Basidiomycota	23/11/2021	Seco	125
BRT452	<i>Cyrenella elegans</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	18/05/2021	Chuviso	125
BRT357	<i>Cystobasidium keelungensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	07/10/2020	Seco	125
BRT581	<i>Cystobasidium</i> sp.	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	23/11/2021	Seco	125
BRT565	<i>Erythrobasidium</i> sp.	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	23/11/2021	Seco	125
BRT418	<i>Exophiala bergeri</i>	Eurotiomycetes	Pezizomycotina	Ascomycota	02/12/2020	Seco	125
BRT646	<i>Exophiala spinifera</i>	Eurotiomycetes	Pezizomycotina	Ascomycota	24/05/2022	Chuviso	93.75
BRT522	<i>Exophiala spinifera</i>	Eurotiomycetes	Pezizomycotina	Ascomycota	19/10/2021	Seco	12.5
BRT582	<i>Exophiala spinifera</i>	Eurotiomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	62.5
BRT496	<i>Exophiala spinifera</i>	Eurotiomycetes	Pezizomycotina	Ascomycota	15/06/2021	Chuviso	81.25
BRT466	<i>Exophiala spinifera</i>	Eurotiomycetes	Pezizomycotina	Ascomycota	18/05/2021	Chuviso	125
BRT654	<i>Exophiala spinifera</i>	Eurotiomycetes	Pezizomycotina	Ascomycota	24/05/2022	Chuviso	125
BRT558	<i>Exophiala spinifera</i>	Eurotiomycetes	Pezizomycotina	Ascomycota	23/11/2021	Seco	500
BRT468	<i>Fellomyces</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	18/05/2021	Chuviso	125


<i>penicillatus/borneensis/polyborus</i>							
BRT716	<i>Halobasidium xiangyangense</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	19/07/2022	Chuvoso	375
BRT672	<i>Hannaella phetchabunensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	23/06/2022	Chuvoso	581.25
BRT670	<i>Hannaella siamensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	23/06/2022	Chuvoso	118.75
BRT483	<i>Hannaella siamensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	15/06/2021	Chuvoso	12.5
BRT353	<i>Hannaella siamensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	07/10/2020	Seco	125
BRT553	<i>Hannaella sinensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	23/11/2021	Seco	62.5
BRT703	<i>Hannaella taiwanensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	19/07/2022	Chuvoso	75
BRT712	<i>Hannaella taiwanensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	19/07/2022	Chuvoso	125
BRT458	<i>Hannaella taiwanensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	18/05/2021	Chuvoso	250
BRT675	<i>Hannaella taiwanensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	23/06/2022	Chuvoso	450
BRT612	<i>Hannaella zaeae</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	07/12/2021	Seco	375
BRT438	<i>Hanseniaspora opuntiae</i>	Saccharomycetes	Saccharomycotina	Ascomycota	18/05/2021	Chuvoso	125
BRT444	<i>Hasegawazyma</i> sp.	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	18/05/2021	Chuvoso	37.5
BRT368	<i>Hasegawazyma</i> sp.	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	07/10/2020	Seco	125
BRT426	<i>Hasegawazyma</i> sp.	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	02/12/2020	Seco	125
BRT433	<i>Hortaea werneckii</i>	Dothideomycetes	Pezizomycotina	Ascomycota	02/12/2020	Seco	2625
BRT694	<i>Kordyana</i> sp.	Exobasidiomycetes	Ustilaginomycotina	Basidiomycota	23/06/2022	Chuvoso	125
BRT555	<i>Kwoniella dejecticola</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	23/11/2021	Seco	87.5
BRT495	<i>Kwoniella dendrophila</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	15/06/2021	Chuvoso	12.5
BRT406	<i>Kwoniella heveanensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	02/12/2020	Seco	12.5
BRT375	<i>Kwoniella heveanensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	10/11/2020	Seco	75
BRT515	<i>Kwoniella heveanensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	19/10/2021	Seco	125
BRT376	<i>Kwoniella heveanensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	10/11/2020	Seco	137.5
BRT377	<i>Kwoniella heveanensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	10/11/2020	Seco	375
	<i>Meyerozyma</i>						
BRT517	<i>caribbica/guilliermondii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	19/10/2021	Seco	62.5
	<i>Meyerozyma</i>						
BRT439	<i>caribbica/guilliermondii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	18/05/2021	Chuvoso	334.375
	<i>Meyerozyma</i>						
BRT437	<i>caribbica/guilliermondii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	18/05/2021	Chuvoso	706.25
	<i>Meyerozyma</i>						
BRT551	<i>caribbica/guilliermondii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	23/11/2021	Seco	12.5
	<i>Meyerozyma</i>						
BRT627	<i>caribbica/guilliermondii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	24/05/2022	Chuvoso	12.5
	<i>Meyerozyma</i>						
BRT408	<i>caribbica/guilliermondii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	02/12/2020	Seco	125
	<i>Meyerozyma</i>						
BRT509	<i>caribbica/guilliermondii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	20/07/2021	Chuvoso	125

BRT549	<i>Meyerozyma caribbica/guilliermondii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	23/11/2021	Seco	125
BRT436	<i>Meyerozyma caribbica/guilliermondii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	18/05/2021	Chuvoso	153.125
BRT526	<i>Meyerozyma caribbica/guilliermondii</i>	Saccharomycetes	Saccharomycotina	Ascomycota	19/10/2021	Seco	156.25
BRT685	<i>Microstroma</i> sp. 1	Exobasidiomycetes	Ustilaginomycotina	Basidiomycota	23/06/2022	Chuvoso	125
BRT354	<i>Microstroma</i> sp. 2	Exobasidiomycetes	Ustilaginomycotina	Basidiomycota	07/10/2020	Seco	12.5
BRT350	<i>Microstroma</i> sp. 2	Exobasidiomycetes	Ustilaginomycotina	Basidiomycota	07/10/2020	Seco	125
BRT409	<i>Moesziomyces antarcticus</i>	Ustilaginomycetes	Ustilaginomycotina	Basidiomycota	02/12/2020	Seco	12.5
BRT355	<i>Moesziomyces aphidis</i>	Ustilaginomycetes	Ustilaginomycotina	Basidiomycota	07/10/2020	Seco	12.5
BRT397	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	02/12/2020	Seco	412.5
BRT389	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	10/11/2020	Seco	12.5
BRT534	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	19/10/2021	Seco	12.5
BRT477	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	15/06/2021	Chuvoso	25
BRT415	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	02/12/2020	Seco	75
BRT362	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	07/10/2020	Seco	125
BRT366	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	07/10/2020	Seco	125
BRT419	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	02/12/2020	Seco	125
BRT420	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	02/12/2020	Seco	125
BRT427	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	02/12/2020	Seco	125
BRT447	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	18/05/2021	Chuvoso	125
BRT471	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	18/05/2021	Chuvoso	125
BRT652	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	24/05/2022	Chuvoso	125
BRT683	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	23/06/2022	Chuvoso	125
BRT684	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	23/06/2022	Chuvoso	125
BRT449	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	18/05/2021	Chuvoso	200
BRT645	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	24/05/2022	Chuvoso	212.5
BRT381	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	10/11/2020	Seco	250
BRT384	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	10/11/2020	Seco	250
BRT356	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	07/10/2020	Seco	375
BRT463	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	18/05/2021	Chuvoso	375
BRT446	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	18/05/2021	Chuvoso	625
BRT405	<i>Occultifur brasiliensis</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	02/12/2020	Seco	2762.5
BRT349	<i>Occultifur externus</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	07/10/2020	Seco	125
BRT383	<i>Occultifur plantarum</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	10/11/2020	Seco	12.5
BRT492	<i>Occultifur plantarum</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	15/06/2021	Chuvoso	12.5
BRT411	<i>Papiliotrema flavescens</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	02/12/2020	Seco	12.5
BRT486	<i>Papiliotrema flavescens</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	15/06/2021	Chuvoso	12.5


BRT372	<i>Papiliotrema flavesrens</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	10/11/2020	Seco	31.25
BRT484	<i>Papiliotrema flavesrens</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	15/06/2021	Chuvoso	50
BRT500	<i>Papiliotrema flavesrens</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	20/07/2021	Chuvoso	87.5
BRT351	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	07/10/2020	Seco	12.5
BRT476	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	15/06/2021	Chuvoso	12.5
BRT556	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	23/11/2021	Seco	12.5
BRT591	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	07/12/2021	Seco	12.5
BRT504	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	20/07/2021	Chuvoso	37.5
BRT528	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	19/10/2021	Seco	50
BRT599	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	07/12/2021	Seco	75
BRT498	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	20/07/2021	Chuvoso	100
BRT358	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	07/10/2020	Seco	125
BRT359	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	07/10/2020	Seco	125
BRT388	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	10/11/2020	Seco	125
BRT511	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	20/07/2021	Chuvoso	125
BRT543	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	23/11/2021	Seco	143.75
BRT481	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	15/06/2021	Chuvoso	187.5
BRT410	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	02/12/2020	Seco	250
BRT637	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	24/05/2022	Chuvoso	268.75
BRT428	<i>Papiliotrema laurentii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	02/12/2020	Seco	625
BRT502	<i>Papiliotrema miconiae</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	20/07/2021	Chuvoso	37.5
BRT512	<i>Papiliotrema miconiae</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	20/07/2021	Chuvoso	125
BRT488	<i>Papiliotrema rajasthanensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	15/06/2021	Chuvoso	25
BRT572	<i>Papiliotrema rajasthanensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	23/11/2021	Seco	125
BRT385	<i>Papiliotrema</i> sp. 1	Tremellomycetes	Agaricomycotina	Basidiomycota	10/11/2020	Seco	250
BRT487	<i>Papiliotrema</i> sp. 2	Tremellomycetes	Agaricomycotina	Basidiomycota	15/06/2021	Chuvoso	87.5
BRT641	<i>Parapyrenis conica</i>	Sordariomycetes	Pezizomycotina	Ascomycota	24/05/2022	Chuvoso	125
BRT621	<i>Pseudosydowia</i> sp.	Dothideomycetes	Pezizomycotina	Ascomycota	24/05/2022	Chuvoso	75
BRT460	<i>Pseudozyma hubeiensis</i>	Ustilaginomycetes	Ustilaginomycotina	Basidiomycota	18/05/2021	Chuvoso	375
BRT421	<i>Pseudozyma pruni</i>	Ustilaginomycetes	Ustilaginomycotina	Basidiomycota	02/12/2020	Seco	125
BRT364	<i>Pseudozyma tsukubaensis</i>	Ustilaginomycetes	Ustilaginomycotina	Basidiomycota	07/10/2020	Seco	125
BRT379	<i>Rhodosporidiobolus ruineniae</i>	Microbotryomycetes	Pucciniomycotina	Basidiomycota	10/11/2020	Seco	125
BRT554	<i>Rhodosporidiobolus ruineniae</i>	Microbotryomycetes	Pucciniomycotina	Basidiomycota	23/11/2021	Seco	12.5
BRT378	<i>Rhodosporidiobolus ruineniae</i>	Microbotryomycetes	Pucciniomycotina	Basidiomycota	10/11/2020	Seco	125
BRT402	<i>Rhodotorula mucilaginosa</i>	Microbotryomycetes	Pucciniomycotina	Basidiomycota	02/12/2020	Seco	125
BRT592	<i>Rhodotorula mucilaginosa</i>	Microbotryomycetes	Pucciniomycotina	Basidiomycota	07/12/2021	Seco	37.5
BRT398	<i>Rhodotorula mucilaginosa</i>	Microbotryomycetes	Pucciniomycotina	Basidiomycota	02/12/2020	Seco	100
BRT413	<i>Rhodotorula paludigena</i>	Microbotryomycetes	Pucciniomycotina	Basidiomycota	02/12/2020	Seco	62.5

BRT373	<i>Rhynchogastrema noutii</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	10/11/2020	Seco	125
BRT367	<i>Rosettozyma</i> sp.	Microbotryomycetes	Pucciniomycotina	Basidiomycota	07/10/2020	Seco	125
BRT395	<i>Rosettozyma</i> sp.	Microbotryomycetes	Pucciniomycotina	Basidiomycota	10/11/2020	Seco	125
BRT643	<i>Saitozyma flava</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	24/05/2022	Chuvoso	12.5
BRT622	<i>Saitozyma flava</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	24/05/2022	Chuvoso	75
BRT348	<i>Saitozyma ninhbinhensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	07/10/2020	Seco	125
BRT363	<i>Saitozyma ninhbinhensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	07/10/2020	Seco	125
BRT588	<i>Saitozyma ninhbinhensis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	07/12/2021	Seco	987.5
BRT601	<i>Saitozyma podzolica</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	07/12/2021	Seco	96.875
BRT431	<i>Saitozyma podzolica</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	02/12/2020	Seco	125
BRT640	<i>Saitozyma podzolica</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	24/05/2022	Chuvoso	406.25
BRT530	<i>Saitozyma podzolica</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	19/10/2021	Seco	125
BRT634	<i>Sakaguchia oryzae</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	24/05/2022	Chuvoso	131.25
BRT404	<i>Selenophoma</i> sp.	Dothideomycetes	Pezizomycotina	Ascomycota	02/12/2020	Seco	125
BRT423	<i>Selenophoma</i> sp.	Dothideomycetes	Pezizomycotina	Ascomycota	02/12/2020	Seco	125
BRT626	<i>Starmerella ilheusensis</i>	Saccharomycetes	Saccharomycotina	Ascomycota	24/05/2022	Chuvoso	68.75
BRT657	<i>Symmetrospora marina</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	24/05/2022	Chuvoso	100
BRT497	<i>Symmetrospora marina</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	15/06/2021	Chuvoso	12.5
BRT451	<i>Symmetrospora suhii</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	18/05/2021	Chuvoso	125
BRT584	<i>Symmetrospora suhii</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	23/11/2021	Seco	125
BRT648	<i>Symmetrospora suhii</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	24/05/2022	Chuvoso	212.5
BRT470	<i>Symmetrospora suhii</i>	Cystobasidiomycetes	Pucciniomycotina	Basidiomycota	18/05/2021	Chuvoso	375
BRT369	<i>Taphrina</i> sp.	Taphrinomycetes	Taphrinomycotina	Ascomycota	07/10/2020	Seco	125
BRT453	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	18/05/2021	Chuvoso	62.5
BRT380	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	10/11/2020	Seco	125
BRT566	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	23/11/2021	Seco	125
BRT450	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	18/05/2021	Chuvoso	68.75
BRT361	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	07/10/2020	Seco	125
BRT533	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	19/10/2021	Seco	125
BRT583	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	23/11/2021	Seco	125
BRT678	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	23/06/2022	Chuvoso	137.5
BRT598	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	07/12/2021	Seco	225
BRT386	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	10/11/2020	Seco	250
BRT387	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	10/11/2020	Seco	250
BRT432	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	02/12/2020	Seco	250
BRT464	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	18/05/2021	Chuvoso	250
BRT713	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	19/07/2022	Chuvoso	500
BRT417	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	02/12/2020	Seco	875

BRT422	<i>Tremella ananatis</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	02/12/2020	Seco	1125
BRT485	<i>Tremella</i> sp.	Tremellomycetes	Agaricomycotina	Basidiomycota	15/06/2021	Chuvoso	50
BRT644	<i>Ustilago maydis</i>	Ustilaginomycetes	Ustilaginomycotina	Basidiomycota	24/05/2022	Chuvoso	12.5
BRT606	<i>Vishniacozyma alagoana</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	07/12/2021	Seco	375
BRT611	<i>Vishniacozyma alagoana</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	07/12/2021	Seco	50
BRT455	<i>Vishniacozyma alagoana</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	18/05/2021	Chuvoso	125
BRT469	<i>Vishniacozyma alagoana</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	18/05/2021	Chuvoso	125
BRT575	<i>Vishniacozyma alagoana</i>	Tremellomycetes	Agaricomycotina	Basidiomycota	23/11/2021	Seco	375
BRT407	<i>Yueomyces</i> sp.	Saccharomycetes	Saccharomycotina	Ascomycota	02/12/2020	Seco	12.5



Supplementary Figure 1- Frequencies in bromeliad samples in percentage and categories of functional traits expressed by yeasts in different periods, dry and rainy.



Supplementary Figure 2- Relationship of the first component (CAP1) from the Canonical Analysis of Principal coordinates (CAP) and the environmental variables indicating the significant relationships observed.