

**UNIVERSIDADE FEDERAL DE ALAGOAS
INSTITUTO DE CIÊNCIAS BIOLÓGICAS E DA SAÚDE
Programa de Pós-Graduação em Diversidade Biológica e Conservação nos
Trópicos**

JAYNE SILVA SANTOS

**INFLUÊNCIA DO TIPO DE SOLO EM RESPOSTAS ECOFISIOLÓGICAS
VINCULADAS À ESCASSEZ HÍDRICA E À REIDRATAÇÃO EM *Schinus
terebinthifolia* Raddi, UMA ESPÉCIE ARBÓREA TROPICAL**

**MACEIÓ - ALAGOAS
Maio/2019**

JAYNE SILVA SANTOS

**INFLUÊNCIA DO TIPO DE SOLO EM RESPOSTAS ECOFISIOLÓGICAS
VINCULADAS À ESCASSEZ HÍDRICA E À REIDRATAÇÃO EM *Schinus
terebinthifolia* Raddi, UMA ESPÉCIE ARBÓREA TROPICAL**

Dissertação apresentada ao Programa de Pós-Graduação em Diversidade Biológica e Conservação nos Trópicos, Instituto de Ciências Biológicas e da Saúde da Universidade Federal de Alagoas, como requisito para obtenção do título de Mestre em CIÊNCIAS BIOLÓGICAS, área de concentração em CONSERVAÇÃO DA BIODIVERSIDADE TROPICAL.

**Orientador: Prof Dr. Gilberto Costa
Justino**

**MACEIÓ - ALAGOAS
Maio/2019**

**Catalogação na fonte
Universidade Federal de Alagoas
Biblioteca Central**

Bibliotecário: Marcelino de Carvalho

S237i Santos, Jayne Silva.

Influência do tipo de solo em respostas ecofisiológicas vinculadas à escassez hídrica e à reidratação em *Schinus terebinifolia* Raddi, uma espécie arbórea tropical / Jayne Silva Santos. – 2019.

89 f. : il.

Orientador: Gilberto Costa Justino.

Dissertação (Mestrado em Ciências Biológicas) – Universidade Federal de Alagoas. Instituto de Ciências Biológicas e da Saúde. Programa de Pós-Graduação em Diversidade Biológica e Conservação. Maceió, 2019.

Inclui bibliografias.

1. Desidratação. 2. Adaptação fisiológica. 3. *Anacardiaceae*. 4. Fenômenos fisiológicos vegetais. I. Título.

CDU: 582.765:581.1


Folha de aprovação

Jayne Silva Santos


**INFLUÊNCIA DO TIPO DE SOLO EM RESPOSTAS
ECOFISIOLÓGICAS VINCULADAS À ESCASSEZ HÍDRICA E À
REIDRATAÇÃO EM *Schinus terebinifolius* RADDI, UMA
ESPÉCIE ARBÓREA TROPICAL**


Dissertação apresentada ao Programa de Pós-Graduação em Diversidade Biológica e Conservação nos Trópicos, Instituto de Ciências Biológicas e da Saúde da Universidade Federal de Alagoas, como requisito para obtenção do título de Mestre em CIÊNCIAS BIOLÓGICAS na área da Biodiversidade.

Dissertação aprovada em 28 de maio de 2019.


Prof. Dr. Gilberto Costa Justino/UFAL
Orientador


Profa. Dra. Keila Rego Mendes/INPA
(membro titular)


Prof. Dr. Marcelo Teixeira de Oliveira/UFPE
(membro titular)


Prof. Dr. Guilherme Ramos Demétrio Ferreira/UFAL
(membro titular)

MACEIÓ - AL

Maio/2019

DEDICATÓRIA

*À minha família, eles são minha estrutura, minha base, minha força e coragem.
Eles me ensinaram sobre amor, respeito, fé e paciência, coisas que nenhum curso no mundo é capaz de ensinar.*

AGRADECIMENTOS

À Universidade Federal de Alagoas (UFAL), através do Programa de Pós-Graduação em Diversidade Biológica e Conservação dos Trópicos (PPG-DIBICT), pela oportunidade na realização deste curso.

À Fundação de Amparo à Pesquisa do Estado de Alagoas (FAPEAL) pelo fornecimento da bolsa de mestrado que recebi durante o curso, bolsa que me manteve na capital Alagoana, longe da casa dos meus pais.

Ao Prof. Dr. Gilberto Costa Justino, meu orientador, pelo acolhimento, conhecimentos repassados, pela paciência e compreensão, que juntamente com o Prof. Dr. Laurício Endres, responsável pelo Laboratório de Fisiologia Vegetal - CECA, incentivam a pesquisa, o avanço do conhecimento e da ciência no âmbito da botânica.

A todos os integrantes do Laboratório de Ecofisiologia Vegetal do Instituto de Ciências Biológicas e da Saúde (ICBS - UFAL) e do Laboratório de fisiologia Vegetal Centro de Ciências agrárias (CECA – UFAL), que, sem eles esse trabalho se tornaria impossível de ser realizado.

Aos companheiros de curso, da turma de mestrado e doutorado do ano de 2017, por todo companheirismo e amizade de alguns.

À toda minha família, em especial à minha mãe Edna, meu pai José, minha irmã Jayllene, minha vó Eunice (em memória), meu namorado Italo e aos amigos e que me incentivam nos estudos e lutas diárias.

À todos estes, meu eterno agradecimento pelo apoio nesta difícil fase da vida acadêmica, foram eles os responsáveis pela minha formação ética, moral, pessoal, espiritual e profissional.

AGRADEÇO!

EPÍGRAFE

“Pra me reconhecer de volta, pra me reprender e me apreender de novo”

Ana Caetano / Tiago Iorc

RESUMO

Schinus terebinthifolia Raddi (Anacardiaceae) é uma espécie arbórea tropical recomendada para a restauração florestal e que pode ser encontrada em várias localidades do litoral, principalmente em formações florestais da restinga, floresta semidecídua e floresta tropical aberta. Por ser encontrada em ambientes tão diversos, e apresentar grande potencial invasivo, acredita-se que esta espécie apresente grande capacidade de adaptação ao estresse hídrico e diferentes tipos de solo, podendo colonizar ambientes degradados e o semiárido brasileiro. Diante disso, o objetivo deste trabalho foi verificar o comportamento ecofisiológico de *Schinus terebinthifolia* Raddi sob condições de seca seguida de reidratação em três tipos de solo, oriundos de três diferentes tipologias da Mata Atlântica do nordeste do Brasil. O estudo foi dividido em dois experimentos. No experimento 1, foi utilizado em apenas um tipo de solo, oriundo da zona da mata alagoana. Já no experimento 2, foram usados 2 tipos de solo: (i) restinga e (ii) e semiárido para submeter as plantas ao déficit hídrico e a reidratação. Os experimentos foram conduzidos em casa de vegetação com distribuição completamente aleatória em dois tratamentos, sendo: (a) controle e (b) irrigação suspensa. As plantas do tratamento “irrigação suspensa” tiveram a hidratação interrompida até que a média dos valores de fotossíntese chegasse a valores próximos ou iguais a zero, e, após isso, as plantas deste tratamento foram reidratadas. Todos os tipos de solos passaram por análises em densímetro, peneiramento, grau de floculação e análise de fertilidade. No dia de maior estresse e após a reidratação foram realizadas análises de trocas gasosas, fluorescência da clorofila e quantificação de compostos osmorreguladores. Dados microclimáticos da casa de vegetação foram acompanhados durante todo o experimento. A primeira reação ao estresse foi a diminuição da condutância estomática que reduziu gradativamente conforme o déficit hídrico se instalava independentemente do tipo solo, indo de 0,4 a 0,01 ($\text{mmol H}_2\text{O m}^{-2}\text{s}^{-1}$) em média. A aroeira também conservou a integridade dos pigmentos cloroplastídicos importantíssimos na captura de energia luminosa, e regulou seu potencial osmótico principalmente com o aumento de aminoácidos e prolina. Com isso, plantas jovens de *S. terebinthifolia* foram resistentes à seca severa com rápida capacidade de recuperação nos três tipos de solo, tendo respostas mais severas em solo com feições argilosas, provavelmente, devido a maior

compactação do solo seco e de características de disponibilidade hídrica neste tipo de solo.

Palavras-chave: estresse hídrico, plasticidade fisiológica, aroeira.

ABSTRACT

Schinus terebinthifolia Raddi (Anacardiaceae) is a neotropical tree species recommended for forest restoration that can be found in several coastal locations, mainly in 'restinga', semideciduous forest and open tropical forest. The ability to grow in diverse environments may be related to its capacity to adapt to water stress and different soil types, allowing it to colonize degraded environments and even the Brazilian semi-arid (caatinga) region. The objective of this work was therefore to verify the ecophysiological behavior of *Schinus terebinthifolia* Raddi under dry conditions followed by rehydration in three types of soil, from two different types of Atlantic Forest in northeastern Brazil, more one from the Caatinga. The study was divided into two experiments. In experiment 1, only soil from an area of Atlantic forest forest was used. In experiment 2, two types of soil were used: (i) restinga soil, and; (ii) soil from the semi-arid biome. The experiments were conducted in a greenhouse with completely random distribution in two treatments: (i) control, and; (ii) water stress ('suspended irrigation'). Irrigation was suspended until average photosynthesis values reached levels near or equal to zero. Plants of this treatment were then rehydrated. All soil types were analyzed for density, screening, flocculation and fertility. Gas exchange, chlorophyll fluorescence and osmoregulatory compounds were quantified on the day of greatest stress and after rehydration. Microclimatic data from the greenhouse were collected throughout the experiment. The first reaction to stress was a decrease in stomatal conductance, which reduced gradually as water deficit settled (independent of soil type) from 0.4 to 0.01 ($\text{mmol H}_2\text{O m}^{-2}\text{s}^{-1}$). Chloroplastid pigments retained their integrity throughout the treatment, and osmotic potential was regulated mainly through an increase in amino acids and proline. Young *S. terebinthifolia* were thus resistant to severe drought and had the capacity for rapid recovery in all three soil types. Physiological responses were more severe in the clay soil, probably due to root damage associated with soil compaction during drying.

Key words: water stress, physiological plasticity, aroeira.

LISTA DE FIGURAS

Capítulo 1

Figure. 1 Average hourly values of air temperature (°C), relative air humidity (%) and vapor pressure deficit (Kpa), recorded every thirty minutes and during twelve days of the experiment.....	34
Figure. 2 (A) Soil water potential (B) Soil moisture during the experiment.....	37
Figure. 3 Gaseous exchanges of <i>Schinus terebinthifolia</i> Raddi accompanied during twelve days of experiment: (A) Photosynthesis, (B) Stomatal conductance, (C) Transpiration, (D) Carboxylation efficiency, (E) Instantaneous water use efficiency and (F) Intrinsic water use efficiency.....	38
Figure. 4 Values of leaf water potential in <i>Schinus terebinthifolia</i> Raddi plants were verified at the first, seventh and twelfth day of the experiment at two times of day: (A) beforehand and (B) Midday.....	39
Figure. 5 (A) Quantification of soluble carbohydrates, (B) Free amino acid content and (C) Proline in leaves of <i>Schinus terebinthifolia</i> Raddi plants verified on the seventh and twelfth day of experiment.....	40
Figure. 6 Ureide content in leaves of <i>Schinus terebinthifolia</i> Raddi plants on the seventh and twelfth day of experiment: (A) Allantoic acid, (B) Allantoin and (C) Total ureides and (D) Nitrate.....	41
Figure. 7 Photochemical efficiency of <i>Schinus terebinthifolia</i> Raddi observed at the first, seventh and twelfth day of experiment: (A) Maximum quantum yield (F_v/F_m) evaluated at 4:00 am; (B) Maximum quantum production (F_v/F_m) evaluated at 12:00 o'clock; (C) Effective quantum efficiency of photosystem II.....	42

Capítulo 2

Figure 1. Average hourly values of air temperature (°C), relative humidity (%), radiation ($\mu\text{mol/m}^2/\text{sec}$) and vapor pressure deficit (Kpa), recorded every fifteen minutes and during nineteen experimental periods.....	61
Figure 2. Temporal monitoring of soil moisture in both soil types (soil 1 and soil 2) and in both water treatments (control and drought). The arrow indicates the day of greatest stress (maximum stress).....	67
Figure 3. Gaseous exchanges of young plants of <i>Schinus terebinthifolia</i> Raddi for 19 days. A Control 1 and 2 (A); A Dry only 1 and 2 (B); gs control soil 1 and 2 (C); gs dry soil 1 and 2 (D); E control solo 1 and 2 (E); E dry only 1 and 2 (F); Ci control soil 1 and 2 (G); Ci Dry 1 and 2 (H); A/Ci controls only 1 and 2 (I); A/Ci Dry 1 and 2 (J); EUA control soil 1 and 2 (K); EUA dry soil 1 and 2 (L); EIUA control only 1 and 2 (M); EIUA dry soil 1 and 2 (N).....	68

LISTA DE TABELAS

Capítulo 1

Table 1. Physical soil analysis using (1) clay (densimeter); (2) fine and coarse sand (sieving); (3) clay dispersed in water (densimeter); (4) degree of flocculation (GF).....	42
---	----

Capítulo 2

Table 1. Physical soil analysis using (1) clay (densimeter); (2) fine and coarse sand (sieving); (3) clay dispersed in water (densimeter); (4) degree of flocculation (GF).....	59
Table 2. Soil fertility analysis using (1) Mehlich-1 extractor; (2) 1.0 M KCl extractor; (3) Calcium acetate extractor at pH 7.0; (4) Welkley-Black method; (5) Base Saturation; (6) Saturation by aluminum.....	60
Table 3. Summary of the analysis of variance of the interaction between two soil types (soil 1 and soli 2) and water regime (control and dry) with the gas exchange variables (A, gs, E, Ci, A/Ci, WUE, iWUE), photochemical (PSII, Fv/Fma.m. and Fv/Fmp.m.), water potential ($\Psi_{wa.m}$ and $\Psi_{wp.m}$), green intensity (chlorophyll a, chlorophyll b, carotenoids, total chlorophyll, ratio clor a/b and SPAD), water potential ($\Psi_{wa.m}$ and $\Psi_{wp.m}$), osmoregulators (proline, protein, amino acid, total sugars, total ureides, allantoin, allantoic acid) and compounds of nitrate metabolism (nitrate reductase, nitrate) in the four days of data collection.....	71
Table 4. Unfolding the interaction between two types of soil (soil 1 and soil 2.) and water regime (control and dry) with the variables of gas exchanges (Ci, WUE, iWUE) and water potencial (Ψ_w 12hs) in stress maximum.....	72
Table 5. Unfolding the interaction between two type soil (soli 1 and soil 2.) and water regime (control and dry) with the variables of chloroplastic pigments (Cloro a, b, total, carotenoids) in stress maximum.....	72
Table 6. Unfolding the interaction between two type soil (soli 1 and soil 2.) and water regime (control and dry) with the variables of osmoregulators (proline, protein, ureids, allantoin) and compounds of nitrogen metabolism (nitrate and RN) in stress maximum.....	74

SUMÁRIO

Texto principal	
1 Apresentação.....	12
2 Revisão de literatura	14
2.1 Escassez hídrica e respostas fisiológicas nos vegetais.....	14
2.2 Solo e sua relação com os vegetais	16
2.3 <i>Schinus terebinthifolia</i> , espécie estudada.....	19
Referências	22
Capítulo 1	
Artigo 1: Can rehydration restore the photosynthetic apparatus of	
<i>Schinus terebinthifolia</i> (Raddi) after severe soil drought?.....	29
Abstract.....	30
Introducion.....	31
Material and methods.....	33
Plant material and experimental conditions	33
Leaf water potential	34
Quantum Efficiency of Photosystem II	34
Gas exchange and water use efficiency.....	35
Contents of photosynthetic pigments and SPAD Index.....	35
Biochemical analyses.....	35
Statistical analysis.....	36
Results.....	37
Discussion.....	43
References.....	47

Capítulo 2

Artigo 2: Physiological responses of <i>Schinus terebinthifolia</i> Raddi (Anacardiaceae) submitted to water deficiency followed by rehydration in two types of soil.....	55
Abstract.....	56
Introduction	57
Material and methods.....	59
Experimental design and physical-chemical analysis of soil.....	59
Determination of water potential and soil moisture.....	62
Determination of gas exchange; instantaneous water-use (iWUE) and intrinsic water use efficiency (IWUE)	62
Determination of the water potential of the leaf.....	62
Extraction of compounds.....	62
Quantification of nitrate	62
Quantification of ureides (Allantoin and Alantoic Acid)	63
Quantification of the proline content.....	63
Quantification of soluble carbohydrates.....	63
Quantification of free amino acids.....	64
Activity of the nitrate reductase enzyme.....	64
Quantification of the spectrometric content of chlorophyll.....	64
Quantification of colorimetric chlorophyll content (SPAD index).....	64
Determination of chlorophyll a fluorescence, determination of the maximum and effective quantum yield of photosystem II.....	65
Statistical analyzes.....	65
Results.....	66
Soil humidity.....	66
Gas exchange	67
Variables affected by water stress and soil type.....	69
Discussion.....	75
References	78

1 APRESENTAÇÃO

Schinus terebinthifolia Raddi é uma espécie arbórea tropical da família Anacardiaceae que tem grande importância farmacêutica, ambiental, ecológica e para projetos de restauração. É especialmente estudada na etnobotânica há algumas décadas por conta de suas propriedades médicas. Conhecida popularmente como aroeira-da-praia ou pimenta-rosa (Brazilian pepper tree - inglês), esta espécie é nativa da América do Sul (CORRÊA, 1974) e se tornou uma espécie considerada invasora em alguns países da América do Norte por conta de sua alta plasticidade e adaptabilidade (EWEL et al. 1982). No Brasil está presente em diversas localidades do litoral (ÁLVARES-CARVALHO et al. 2016), sendo encontrada principalmente em formações florestais de restinga, floresta semidecidual (CESÁRIO, 2008) e formações ombrófilas abertas, entretanto, em áreas semiáridas ainda não há registro de ocorrência dessa espécie (SpeciesLink, 2019).

Na Restinga, o período de baixa precipitação pode induzir restrição hídrica (JÚNIOR BOEGER, 2016), acompanhado de salinidade, alta radiação solar e de ventos constantes, fatores importantes quando se trata de manutenção de água na planta (MANTOVANI et al. 2010), fatores estes, que na floresta semidecidual e formações ombrófilas abertas estão amenizados, e englobam um limiar climático ótimo para esta espécie. Dessa forma, o potencial de sobrevivência de *S. terebinthifolia* ao estresse hídrico associado a outras adversidades encontradas em seu habitat natural de restinga ainda não foi totalmente descrito, e deve estar ligada a uma ampla plasticidade de respostas fisiológicas e ecológicas provavelmente presentes nesta espécie, e que poderia ser um ponto crucial para que a aroeira se propagasse por outras regiões sazonalmente secas e com diferentes tipos de solo, como o semiárido brasileiro onde a espécie ainda não é encontrada.

Ademais, o arranjo das espécies no ambiente é determinado, dentre outros fatores, pelas condições de estrutura e de nutrição do solo (CARVALHO et al. 2007; HARIDASAN, 2000). A água presente nesse substrato é um dos fatores abióticos mais limitantes, e juntamente com os nutrientes do solo, e suas características físicas, parecem ser decisivos na distribuição das espécies em seus ambientes (DUARTE et al. 2012; CARVALHO et al. 2017; BOTREL et al. 2002). Nesta dissertação são descritas respostas fisiológicas de *Schinus terebinthifolia* Raddi à restrição hídrica,

juntamente com a dissolução dessas respostas frente a três tipos diferentes de solos oriundos de diferentes tipologias da floresta atlântica neotropical.

2 REVISÃO DE LITERATURA

2.1 Escassez hídrica e as respostas fisiológicas dos vegetais

Para o sucesso na colonização do ambiente terrestre, as plantas precisaram se ajustar à alguns fatores cruciais como a luz, absorção do dióxido de carbono, oxigênio, nutrição mineral do solo e principalmente à quantidade de água. Destes, a disponibilidade de água é um dos fatores ambientais que exerce a maior influência sobre as taxas fotossintéticas em espécies vegetais (SOUZA et al. 2010). Com isso, a restrição hídrica leva ao desenvolvimento de estratégias fisiológicas para a manutenção do metabolismo e, consequentemente, à sobrevivência das espécies nestas condições. Neste sentido, podemos então dizer que, as espécies que consigam ajustar o seu metabolismo em diferentes situações de disponibilidade de água, serão as mesmas com aptidão necessária para enfrentar períodos de escassez e/ou abundância hídrica previstas para o clima na terra em um futuro próximo, que indicam drásticas mudanças nas distribuições de chuva em todo o mundo (BOISVENUE et al. 2006).

Como já mencionado, a água é classificada como um fator vital que limita a distribuição das espécies no planeta, e sua disponibilidade no ambiente irá depender de vários fatores climáticos que influenciam a distribuição das chuvas, temperatura e da radiação solar, como também das correntes marítimas e atmosféricas, além de questões geológicas e atividades antrópicas (TRENBERTH et al. 2015). Ou seja, a presença de água no ambiente natural vem acompanhada de várias outras variáveis ambientais muito importantes na fisiologia das espécies vegetais (POREMBSKI, 2007), o que consequentemente coordena as estratégias de controle da quantidade de água presente no próprio corpo vegetal. A regulação da abertura e do fechamento estomático é o que controla a transpiração através das folhas das plantas, sendo bastante eficiente para este fim (DOMBROSKI et al. 2011). Com a abertura dos estômatos, as plantas absorvem o CO₂, realizam fotossíntese e fazem da transpiração o melhor mecanismo para dissipar o excesso de calor. Mas, por outro lado, perdem bastante água para a atmosfera (SILVA et al. 2004).

O ajuste adequado da abertura e fechamento dos estômatos, nestes casos, promove maior eficiência no uso da água, o que maximiza a entrada de CO₂ e reduz a perda de água por transpiração (GONÇALVES et al. 2009). Assim, esta capacidade

de controle estomático pode ter se aperfeiçoado em algumas espécies, principalmente naquelas de ambientes tropicais de clima quente e úmido, por horas quente e seco, e, acredita-se que o nível de eficiência no uso da água de cada espécie seja o diferencial para genótipos mais adaptados.

A disponibilidade hídrica está frequentemente associada a altas temperaturas e excesso de irradiação, e todas estas situações combinadas além de influenciar na transpiração vegetal, levam à fotoinibição devido a danos ao fotossistema II (HUANG et al. 2012). Caso haja restrição hídrica e fechamento estomático, este dano pode ser acentuado, o que irá depender também das características adaptativas da espécie em questão, envolvendo a capacidade da mesma em reparar os danos causados aos fotossistemas (MURATA et al. 2007) através de dissipação de energia e atividades de enzimas antioxidantes, por exemplo.

A água, além de ser necessária para fotossíntese, é matéria prima primordial para o crescimento das plantas, levando a reprodução das células, mantendo a turgescência, saúde e crescimento das mesmas. A manutenção do turgor é importante para permitir a continuidade dos processos internos da célula, incluindo o funcionamento dos fotossistemas citados acima, e por fim, para o crescimento do vegetal em biomassa, ou seja, o déficit hídrico pode afetar a manutenção de todos os processos de produção vegetal (SANTOS, 1998; LÜTTGE, 2007).

Para auxiliar na manutenção da turgência, os vegetais regulam osmoticamente os tecidos, acumulando solutos compatíveis, não tóxicos e reaproveitáveis ao seu metabolismo como aminoácidos livres, prolina, açúcares solúveis e ureídeos, esses últimos, recentemente sugeridos como participantes em mecanismos de resistência ao déficit hídrico em plantas da família Fabaceae (SILVA et al. 2008), sendo necessários estudos em outras famílias botânicas não-nodulantes. Este ajuste osmótico se dá pelo efeito do soluto dissolvido sobre o potencial hídrico, auxiliando na translocação da água por osmose (HUANG et al. 2018; LUO e LUO, 2017). Um dos solutos mais estudados é a prolina, que é produzida à medida que o período de estresse hídrico se prolonga, demonstrando a significativa influência do déficit hídrico na produção e na formação de compostos, para redução do potencial osmótico na planta e, consequentemente, na manutenção de água nas células (ALVARENGA et al. 2011).

O acúmulo de prolina livre também pode ser causado por altas temperaturas e variações de salinidade (NOGUEIRA et al. 2000). Mas, em situações de baixa disponibilidade de água no solo, esse mecanismo adaptativo se faz necessário em todas as espécies e o auxílio deste processo para a manutenção do turgor celular é situação constante em ambientes áridos (SANTOS, 1998). Essas mudanças são vistas na expressão de genes relacionados ao estresse hídrico (SINGH-SANGWAN et al. 1994; SILVA, 2000; KRASENSKY e JONAK, 2012), genes que irão proporcionar as mudanças e alterações na síntese das clorofitas, por exemplo, que estão relacionadas com a eficiência fotossintética das plantas e, consequentemente, com seu crescimento e adaptabilidade aos diferentes ambientes (JESUS, 2008).

Diante de tudo isso, verifica-se que as variações ambientais induzem a expressão de genes que ao primeiro passo codificam sinais bioquímicos, que impulsionam mudanças fisiológicas e que ajustam o metabolismo vegetal de tal forma que resultam em mudanças morfológicas, caracterizando espécies mais adaptadas. Algumas dessas respostas já podem ser classificadas como características intrínsecas de muitas espécies, e abastecem a lista de caracteres que às identificam. Dessa forma, respostas fisiológicas e morfológicas apontam que as plantas, quando são submetidas ao déficit hídrico, regulam seu metabolismo para a conservação da água disponível no solo e no corpo vegetal, otimizando seu uso para processos mais importantes do seu desenvolvimento e buscando a continuidade de sua sobrevivência (SHARMA et al. 2012; MARTINS et al. 2018).

2.2 Os vegetais e sua relação com o solo

O solo, de maneira geral, é formado por materiais minerais e orgânicos, e, por definição, é a coleção de corpos naturais que ocupa a parte superficial da terra (EMBRAPA, 2018). Este substrato se relaciona estreitamente com as plantas, e por isso, é influenciador direto das alterações do metabolismo vegetal, ou seja, o que acontece no solo reflete e causa efeitos nas plantas e o que acontece com a planta causa efeito no solo (CARVALHO et al. 2006). Como exemplo disso, temos a presença de água, a estrutura física do solo e nutrientes presentes nele estreitando essa relação. Condições biológicas, químicas e físicas podem estar vinculadas com a distribuição da vegetação entre os ambientes, pois as necessidades nutricionais e da fixação do organismo do vegetal no solo através das raízes de cada indivíduo variam

entre as espécies botânicas (HARIDASAN, 2000). Essa relação entre os organismos e o solo, em conjunto com a atmosfera, resulta em diferentes formações vegetais, e toda essa interação entre solo-planta-atmosfera, vai ao longo do tempo, modificando a paisagem, tornando esse sistema dinâmico e possibilitando adaptação por partes de todos os organismos e determinando a composição florística das regiões do globo (KOTCHETKOFF-HENRIQUES et al. 2005).

Como dito anteriormente, as plantas têm uma relação particular com o solo, se relacionando com o mesmo, tanto de forma direta e indireta. A forma mais direta dessa relação se dá pela absorção dos nutrientes minerais e da água retirados diretamente do solo pelas raízes das plantas, que por sua vez, liberam certas substâncias no solo e também se decompõem sobre ele (VIEIRA, 1988). Além da absorção, as plantas fazem do solo seu substrato de fixação, se “agarrando” em partículas e até mesmo rochas, resultando em um aumento na coesão aparente do solo que, indiretamente, evita o desenvolvimento de erosões e lixiviação de nutrientes exenciais para o desenvolvimento das próprias plantas (WANG et al. 2016).

A composição edáfica do Brasil engloba solos sedimentares do quaternário, onde a vegetação e todas as suas interfases se estabeleceu (GOMES et al. 1998), isso significa que o substrato brasileiro é fruto de transformações ambientais geologicamente recentes. A região nordeste do Brasil engloba três tipologias ambientais de maior evidência: (1) o litoral, (2) a região de transição entre o litoral e o interior do estado, também chamado de zona da mata (3) e o semiárido. Destes, a restinga presente no litoral (ASSUMPÇÃO et al., 2000) e a caatinga presente no semiárido (TROVÃO et al., 2007) são ecologicamente classificados como ambientes naturalmente estressantes (ZAMITH, et al 2006).

Alguns estudos classificam os solos da região litorânea como podzóis hidromórficos ou Espodossolos, ou, areias quartzosas marinhas (CAMARGO et al. 1987; OLIVEIRA et al. 1992), que normalmente têm textura arenosa e é pobre em nutrientes (GOMES et al. 2004). A zona costeira tem uma vegetação característica arbustivo-arbórea, variando desde formações herbáceas, até árvores que chegam a 20 m de altura (ALMEIDA-JR et al. 2011). Essa composição vegetal é resultado da interação das especificações de um solo arenoso, com uma vegetação adaptada a

este tipo de substrato e clima específico dessa região, que inclue maior radiação e ventos constantes (MANTOVANI et al. 2010).

A segunda região mais importante do Nordeste é usualmente chamada de zona da mata e comprehende o domínio da mata atlântica, sendo uma área acobertada por um regime regular de chuvas, e, por ser relativamente próxima a zona litorânea, chega a receber influencia da dinâmica dos oceanos em seu ciclo de chuvas. Essa dinâmica resulta em uma estabilidade de clima e influênciia diretamente na disponibilidade de água no solo , garantindo o sucesso da colonização vegetal e animal. É uma região de solo francoargilo-arenoso fértil de amplo desenvolvimento agronômico e favorável a habitação humana (SANTOS, et al. 2019).

Já o solo do semiárido é convencionalmente classificado como um tipo de solo com horizonte mineral superficial fracamente desenvolvido, resultado de um reduzido teor de colóides minerais ou orgânicos e pelas condições externas de clima que esta região se encontra. A maioria das regiões do semiárido do Nordeste do Brasil apresenta uma estrutura física pedregosa em sua maior parte, e argilosa em algumas regiões, em geral, o solo da caatinga não retêm água em abundância (ASAAD, et al 2001) e com isso pode influenciar na ocorrência e frequênciia de algumas espécies especialmente adaptadas a estas condições (SILVA et al. 2009). É conhecida por ter um mosaico de florestas secas e vegetação arbustiva, com enclaves de floresta tropical montanhosa, com predominância de vegetação xerófila, aclimatadas a climas secos (LEAL et al. 2003).

Contudo, dentro de uma mesma família botânica, como Anacardeaceae, os vegetais são capazes de sobreviver em diferentes zonas climáticas como também em diferentes tipos de solo, podendo colonizar tanto regiões de restinga como regiões de zona da mata, e em caso raros, também o semiárido, ou ambientes equivalentes em todo o globo, fato que só é possivel graças aos ajustes genéticos, fisiológicos e morfológicos que as plantas apresentam em cada situação, mecanismos diversos que ainda necessitam ser descritos.

2.3 *Schinus terebinthifolia* Raddi

S. terebinthifolia, pertencente à família Anacardiaceae, é conhecida popularmente como aroeira-da-praia e por suas propriedades culinárias também é chamada de pimenta-rosa ou pimenta-vermelha em algumas regiões brasileiras. Esta espécie é natural da América do Sul, especificamente da Argentina, Paraguai e Brasil, utilizada como erva medicinal desde os ancestrais indígenas (JONES et al. 1997) e é amplamente explorada até os dias de hoje pela população. A pressão do extrativismo sobre esta espécie é preocupante, a colheita exagerada dos frutos causa redução no banco de sementes disponíveis na natureza, e está dificultando a distribuição e propagação natural da espécie no Brasil, levando as populações de plantas a entrar em declínio e impedindo a colonização em novos ambientes (ÁLVARES-CARVALHO et al. 2016).

Além de ter grande importância etnobotânica como erva medicinal, apresenta comprovação científica de propriedades anti-inflamatória (ROSAS et al. 2015), antialérgica (CAVALHER-MACHADO et al. 2008), antibiótica (COLE et al. 2014) e antifúngica (MOURA et al. 2010; JOHANN et al. 2008). Sua importância médica-farmacológica é imprescindível para a saúde humana, em especial no combate de doenças que acometem órgãos do sistema reprodutivo e trato urinário da mulher, sendo amplamente usada em cosméticos e fármacos relacionados a esses problemas.

Dentre os ambientes que a aroeira-da-praia é encontrada, a zona litorânea brasileira é o mais estressante deles. Comumente, a vegetação de restinga é caracterizada por um mosaico de habitats, indo da praia até uma formação de vegetação florestal densa, e entre esses extremos encontra-se uma organização vegetal em aglomerados com vários espaços abertos de areia nua (MANTOVANI et al. 2010). As espécies vegetais presentes na restinga são submetidas a combinações ambientais extremas, como alta salinidade, baixa disponibilidade de nutrientes, inundações, seca severa e irradiância elevada (MANTUANO et al. 2006). Por ocorrer neste ambiente, a plasticidade ecológica de *S. terebinthifolia* pode ser a justificativa de esta espécie poder habitar facilmente toda faixa litorânea do Brasil, próxima a rios, córregos e várzeas úmidas de formações secundárias, como também nas dunas existentes na restinga (SILVA et al. 2008) e, mesmo enfrentando tantas adversidades

ambientais, não se conhece a distribuição dessa espécie em ambientes semiáridos (SpeciesLink, 2019).

S. terebinthifolia se tornou uma espécie invasora em muitos países, incluindo partes da América Central, Bermudas, nas Ilhas Bahamas, as Índias Ocidentais, Flórida, Arizona do Sul, Califórnia, Havaí, Norte da África, sul da Ásia e África do Sul e Europa Mediterrânea, compreendendo regiões de clima semiárido frio (BESTELMEYER et al, 2015), onde é internacionalmente conhecida como “Brazilian pepper tree”, (BARBOSA, et al. 2009; SHETTY et al. 2010). Por ser uma espécie pioneira (SILVA et al. 2008), se estabeleceu facilmente nestes locais e acredita-se que sua alta plasticidade ambiental contribuiu para o sucesso de suas colonizações.

Estudos com esta espécie comumente trazem a descrição das respostas de algumas variáveis fisiológicas sob consequência de um determinado ambiente, clima ou situação de estresse. A maioria desses estudos justificam seus resultados com o fato de a aroeira ser pioneira, heliófita e por possuir uma grande capacidade de invasão de novos ambientes (EWE et al, 2004; EWE et al, 2007; ESTEVES et al, 2010; DOS ANJOS, et al 2014). Estudando *S. terebinthifolia* e outras espécies arbóreas pioneiras e não-pioneiras, Coutin (2016) constatou a capacidade fotoprotetora e antioxidante dessas espécies frente a exposição solar e concluiu que plantas pioneiras de pleno sol acumulam vitamina E para evitar a peroxidação lipídica, ou seja, apenas um dos mecanismos bioquímicos fisiológicos que essa espécie possa ter para atingir sucesso em ambiente tropicais estressantes.

Estudos de comportamento silvicultural de árvores nativas após o plantio e suas respostas às condições ambientais também podem ser vistos na literatura. Rorato (2018) por exemplo, estudou o comportamento de seis espécies arbóreas frente variações de clima no inverno (geada) e, dentre estas espécies, estava *S. terebinthifolia* que juntamente com *Eugenia involucrata* foram consideradas espécies resistentes à geada, conclusão tirada pela medida de área foliar, altura e diâmetro do caule dessas plantas sob estas condições. Ou seja, *S. terebinthifolia*, é uma espécie de grande interesse científico, sendo estudada pela genética (ÁLVARES-CARVALHO et al, 2016; CHIOCHETA et al, 2017), em estudos de composição florística (CESÁRIO et al 2007), de interesse em reflorestamento (RORATO et al, 2017), e o potencial invasivo dessa espécie (PRATT et al 2006) frente a salinidade (KOUALI et al, 2017),

alagamento (MIELKE et al, 2005), radiação (SABBI et al, 2010) e escassez hídrica (SILVA et al, 2008; VASCONCELOS-SILVA et al, 2011; PIMENTA-BARRIOS et al, 2012)., Estudos que envolvem o cultivo da espécie em solos da Caatinga ainda não foram verificados na literatura.

Este estudo foi proposto para investigar e descrever as respostas fisiológicas e metabólicas de *S. terebinthifolia* em condições de restrição hídrica e posterior reidratação, em três variações de solo típicos das principais tipologias da Floresta Atlântica da região neotropical, considerando as previsões de mudanças na precipitação e nas variáveis climáticas juntamente com a natureza do ambiente que a espécie foco desse estudo está presente, e a possível colonização do ambiente semiárido brasileiro pela mesma, juntamente com o sucesso de invasão já constatado para esta espécie em outros ambientes. Adicionalmente, estes dados podem servir de base para futuras descrições fisiológicas de espécies com as mesmas características. E por fim, ressaltar a conservação dessa espécie, com o conhecimento de sua importância ecológica e das consequências do efeito da escassez hídrica sobre a mesma.

3 REFERÊNCIAS

- ALMEIDA-JR, E. B.; ZICKEL, C. S.. Análise fitossociológica do estrato arbustivo-arbóreo de uma floresta de restinga no Rio Grande do Norte. *Brazilian Journal Of Agricultural Sciences*. Agrár. Recife, v.7, n.2, p.286-291, 2012.
- ALVARENGA, I. C. A.; QUEIROZ, G. A.; HONÓRIO, I. C. G.; VALADARES, R. V.; MARTINS, E. R.. Prolina livre em alecrim-pimenta sob estresse hídrico antes da colheita. *Revista Brasileira Plantas Medicinais*, Botucatu, v.13, p.539-541, 2011.
- ÁLVARES-CARVALHO, S. V.; DUARTE J. F.; SANTOS T. C. R.; SANTOS, R. M.; SILVA-MANN, R.; CARVALHO D.. Structure and genetic diversity of natural Brazilian pepper populations (*Schinus terebinthifolia* Raddi). *Genetics and Molecular Research* n. 15 (2), 2016.
- BARBOSA, L. C. A; DEMUNER, A. J; CLEMENTE A. D.. Seasonal variation in the composition of volatile oils from *Schinus terebinthifolia* Raddi. *Quimica Nova*, Vol. 30, N.. 8, p. 1959-1965, 2009.
- BESTELMEYER, Brandon T et al. Desertification, land use, and the transformation of global drylands. *Frontiers In Ecology And The Environment*, [s.l.], v. 13, n. 1, p.28-36, fev. 2015. Wiley. <http://dx.doi.org/10.1890/140162>.
- BOISVENUE, C.; RUNNING, S.. Impacts of climate change on natural forest productivity – evidence since the middle of the 20th century. *Global Change Biology* n. 12, p. 862–882, 2006.
- BOTREL, R. T.; OLIVEIRA FILHO, A. T.; RODRIGUES, L. A.; CURI, N.. Influência do solo e topografia sobre as variações da composição florística e estrutura da comunidade arbóreo-arbustiva de uma floresta estacional semidecidual em Ingaí, MG. *Revista Brasil. Bot.*, V.25, n.2, p.195-213, jun. 2002.
- CAMARGO, M. N.; KLAMT, A.; KAUFFMAN, J. H.. Sistema Brasileiro de classificação de solos. International Soil Reference and Information Centre. Wageningen, Holanda, 1987.

CARVALHO, M. H. C.. Drought stress and reactive oxygen species. Plant Signaling & Behavior, [s.l.], v. 3, n. 3, p.156-165, mar. 2008. Informa UK Limited. <http://dx.doi.org/10.4161/psb.3.3.5536>.

CARVALHO, F.; CORREA, M. M.; CARVALHO, G. C.; NETO, F. C. R.; MARINHO, G. P. A.; ANDRADE, S. B.. Enzymatic activity of three sugarcane varieties under salt stress. Revista Brasileira de Engenharia Agrícola e Ambiental, [s.l.], v. 20, n. 9, p.806-810, set. 2016. FapUNIFESP (SciELO). <http://dx.doi.org/10.1590/1807-1929/agriambi.v20n9p806-810>.

CAVALHER-MACHADO S. C.; ROSAS, E. C.; BRITO F. A.; HERINGE A. P.; OLIVEIRA R. R.; KAPLAN M. A. C.; FIGUEIREDO M. R.; HENRIQUES M. G. M. O.. The anti-allergic activity of the acetate fraction of *Schinus terebinthifolia* leaves in IgE induced mice paw edema and pleurisy. International Immunopharmacology n. 8, p. 1552–1560, 2008.

CESÁRIO, L. F.; GAGLIANONE, M. C.. Biologia floral e fenologia reprodutiva de *Schinus terebinthifolia* Raddi (Anacardiaceae) em Restinga do Norte Fluminense. Acta bot. bras. 22(3): 828-833. 2008.

COLE E. R.; DOS SANTOS R. B. V.; LACERDA, J.; MARTINS J. D. L.; GRECO S. J.; CUNHA-NETO, A.. Chemical composition of essential oil from ripe fruit of *Schinus terebinthifolia* Raddi and evaluation of its activity against wild strains of hospital origin. Brazilian Journal of Microbiology n. 45, Vol. 3, p. 821-828, 2014.

CORRÊA, M. P.. Dicionário de plantas úteis do Brasil e das exóticas cultivadas, vol. 3. Imprensa Nacional, Rio de Janeiro, Brazil, pp. 125–126, 1974.

DOMBROSKI, J. L. D.; PRAXEDES, S. C.; FREITAS, R. M. O.; PONTES, F. M.. Water relations of Caatinga trees in the dry season. South African Journal of Botany, n. 77, p, 430–434, 2011.

DUARTE et. al. Eficiência do uso da água na produção de óleo do girassol (*Helianthus annuus* L.), sob suspensão hídrica. Revista Brasileira de Agricultura Irrigada v.6, n. 3, p. 166 - 175, 2012.

EWEL, J. J; OJIMA, D. S; KARL, D. A; DEBUSK, W. F; Schinus in successional ecosystems of Everglades National Park. South Florida Research Report, T-676, Everglades National Park, Homestead, Florida, 1982.

GOMES, F. P.. Novos aspectos do problema do tamanho ótimo das parcelas em experimentos com plantas arbóreas. v.23, n.1, jan. 1988.

GOMES, J. M.; SILVA, A. R.. Os substratos e sua influência na qualidade de mudas. In: BARBOSA, J.G. et al.. (Ed.). Nutrição e adubação de plantas cultivadas em substrato. Viçosa, MG: Universidade Federal de Viçosa, 2004. p.190-225.

GONÇALVES, J. F. C.; SILVA, C. E. M.; GUIMARÃES, D. G.. Características fotossintéticas e potencial hídrico foliar de plantas jovens de andiroba submetidas à deficiência hídrica e à reidratação. Pesquisa Agropecuária Brasileira, v. 44, p. 8-14, 2009.

HARIDASAN, N.. Nutrição mineral de plantas nativas do cerrado. R. Bras. Fisiol.Veg., 12(1):54-64, 2000.

HUANG, W.; FU, P.; JIANG, Y.; ZHANG, J.; ZHANG, S.; HU, H.; CAO, K.. Differences in the responses of photosystem I and photosystem II of three tree species *Cleistanthus sumatranus*, *Celtis philippensis* and *Pistacia weinmannifolia* exposed to a prolonged drought in a tropical limestone forest. Tree Physiology, n. 33, p. 211–220, 2012.

LUO, Z. B.; LUO, J. Uncovering the physiological mechanisms that allow nitrogen availability to affect drought acclimation in *Catalpa bungei*. Tree Physiology, v. 37, p. 1453–1456, 2017. LÜTTGE, U. Physiological ecology of tropical plants. 2º Ed. Springer, Berlin, GE, 2007. 456p.

JESUS, S. V.; MARENCO, R. A.. O SPAD-502 como alternativa para a determinação dos teores de clorofila em espécies frutíferas. Acta Amazônica. v. 38, p. 815- 818, 2008.

JOHANN, S.; SILVA, D. L.; MARTINS, C. V. B.; ZANI C. L.; PIZZOLATTI M. G.; RESENDE M. A.. Inhibitory effect of extracts from Brazilian medicinal plants on the adhesion of *Candida albicans* to buccal epithelial cells. World J Microbiol Biotechnol n. 24:2459–2464, 2008

- JONES, D.. Biology of Brazilian pepper management plan for Florida. Florida, Florida Exotic Pest Plant Council. 1997.
- JÚNIOR, J.C.F.M; BOEGER, M.R.T. Leaf traits and plastic potential of plant species in a light-edaphic gradient from a restinga in southern Brazil. *Acta Biológica Colombiana*. 2016.
- KOTCHETKOFF-HENRIQUES, O.; JOLY, C. A.; BERNACCI, L. C.. Relação entre o solo e a composição florística de remanescentes de vegetação natural no Município de Ribeirão Preto, SP *Revista Brasil. Bot.*, V.28, n.3, p.541-562, jul.-set. 2005.
- KRASENSKY, J.; JONAK, C. Drought, salt, and temperature stress-induced metabolic rearrangements and regulatory networks. *Journal of Experimental Botany*, v. 63, p. 1593-1608, 2012.
- LEAL, I. R.; TABARELLI, M.; SILVA J. M. C.. Ecologia e conservação da caatinga. Ed Universitária da UFPE, 2003.
- LUO, Z. B.; LUO, J. Uncovering the physiological mechanisms that allow nitrogen availability to affect drought acclimation in *Catalpa bungei*. *Tree Physiology*, v. 37, p. 1453–1456, 2017. LÜTTGE, U. *Physiological ecology of tropical plants*. 2º Ed. Springer, Berlin, GE, 2007. 456p.
- MANTOVANI, A.; IGLESIAS, R. R.. The effect of water stress on seed germination of three terrestrial bromeliads from restinga. *Revista Brasil. Bot.*, V.33, n.1, p.201-205, 2010
- MANTUANO, D. G.; BARROS, C. F.; SCARANO, F. R.. Leaf anatomy variation within and between three “restinga” populations of *Erythroxylum ovalifolium* Peyr. (Erythroxylaceae) in Southeast Brazil. *Revista Brasileira de Botânica*, V.29, n.2, p.209-215, 2006.
- MARTINS, K. et al. Plant Growth, Antioxidative Enzymes, Lipid Peroxidation and Organic Solute Contents in Mulungu Seedlings (*Erythrina velutina*) Under Different Field Capacities. *Journal of Agricultural Science*, v. 10, p. 443, 2018.

MOURA, T. F. A. L.; RAFFI F. N.; SANTOS A. L. R.. Evaluation of a preservative system in a gel containing hydroalcoholic extract of *Schinus terebinthifolia*. Brazilian Journal of Pharmacognosy 21(3): 532-536, May./Jun. 2010

MURATA, N.; TAKAHASHI, S.; NISHIYAMA, Y.; ALLAKHVERDIEV, S. I.. Photoinhibition of photosystem II under environmental stress. Biochimica et Biophysica Acta, v. 1767, n. 6, p. 414–42, 2007.

NOGUEIRA. R. J. N. C.; MORAES, J. A. P. V.; BURITY, H. A.; NETO, E. B.. Alterações na resistência à difusão de vapor das folhas e relações hídricas em aceroleiras submetidas a déficit de água. Revista Brasileira de Fisiologia Vegetal, n. 13(1):75-87, 2000.

OLIVEIRA L. M. B.; BEVILAQUA C. M. L.; MACEDO I. T. F.; MORAIS S. M.; MACHADO L. K. A.; CAMPELLO C. C.; MESQUITA M. A.. Effects of Myracrodroon urundeuva extracts on egg hatching and larval exsheathment of *Haemonchus contortus*. Parasitology Research. n. 109: 893–898, 2011.

POREMBSKI, S. Tropical inselbergs: habitat types, adaptive strategies and diversity patterns. Revista Brasileira de Botânica, v. 30(4), p. 579–586, 2007.

ROSAS E. C.; CORREA L. B.; PÁDUA T. A.; COSTA, T. E. M. M.; MAZZEI J. L.; HERINGER, A. P.; BIZARRO C. A.; KAPLAN, M. A. C.; FIGUEIREDO M. R.; HENRIQUES M. G.. Anti-inflammatory effectof *Schinus terebinthifolia* Raddi Hydroal coholic extracton neutrophil migration inzy mosan-induced arthritis Journal of Ethnopharmacology. n. 17, p. 5490–498, 2015.

ASSAD MLL; SANS, LMA; ASSAD, ED; ZULLO JR, J. Relação entre água retida e conteúdo de areia total em solos brasileiros. Revista Brasileira de Agrometeorologia, Passo Fundo, v.9, n.3, p.588-596, 2001.

SANTOS, R. F.; CARLESSO, E. R.. Déficit Hídrico e os Processos Morfológico e Fisiológico Das Plantas. Revista Brasileira de Engenharia Agrícola e Ambiental. Campina Grande, PB. v.2, n.3, p.287-294, 1998.

SANTOS, P. Z. F; CROUZEILLES, R; SANSEVERO, J. B. B. Can agroforestry systems enhance biodiversity and ecosystem service provision in agricultural

landscapes? A meta-analysis for the Brazilian Atlantic Forest. *Forest Ecology And Management*, [s.l.], v. 433, p.140-145, fev. 2019. Elsevier BV.

SANTOS, H. G.; JACOMINE, P. K. T.; ANJOS, L. H. C. dos; OLIVEIRA, V. A. de; LUMBRERAS, J. F.; COELHO, M. R.; ALMEIDA, J. A. de; ARAUJO FILHO, J. C. de; OLIVEIRA, J. B. de; CUNHA, T. J. F. Sistema Brasileiro de Classificação de Solos. 5. ed. rev. e ampl. Brasília, DF. 356 p. il. color. ISBN: 978-85-7035-800-4. 2018.

SHARMA, P. et al. Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *Journal of Botany*, v. 2012, ID: 217037, 2012.

SHETTY, K. G.; MINNIS, A. M.; ROSSMAN, A. Y.; JAYACHANDRAN, K.. The Brazilian peppertree seed-borne pathogen, *Neofusicoccum batangarum*, a potential biocontrol agent. *Biological Control* n. 56, p. 91–97, 2010.

SILVA, F. A. M.; MELLONI, R.; MIRANDA, J. R. P.; CARVALHO, J. G.. efeito do estresse salino sobre a nutrição mineral e o crescimento de mudas de aroeira (*Myracrodruon urundeuva*) cultivadas em solução nutritiva. *CERNE*, V.6, N.1, P.052-059, 2000.

SILVA, M. A. V.; NOGUEIRA, R. J. M. C.; OLIVEIRA, A. F. M.; SANTOS, V. F.; Resposta estomática e produção de matéria seca em plantas jovens de aroeira submetidas a diferentes regimes hídricos. *Revista Árvore*, Viçosa-MG, v.32, n.2, p.335-344, 2008.

SILVA, E. C. et al. Aspectos ecofisiológicos de dez espécies em uma área de caatinga no município de Cabaceiras, Paraíba, Brasil. *Iheringia*, Porto Alegre, RS, v. 59(2), p. 201-205, 2004.

SINGH-SANGWAN, N.; FAROOQI, A. H. A.; SANGWAN, R. S.. Effect of drought stress on growth and essential oil metabolism in lemongrasses. *New Phytol.* 128, 173-179, 1994.

SOUZA, B. D.; MEIADO, M. V.; RODRIGUES, B. M.; SANTOS, M. G.. Water relations and chlorophyll fluorescence responses of two leguminous trees from the Caatinga to different watering regimes. *Acta Physiologiae Plantarum*, v. 2, p. 235-244, 2010.

TRENBERTH, K. E.; FASULLO, J. T.; SHEPHERD, T. G.. Attribution of climate extreme events. *Nature climate change*, v. 5, p. 725-730, august 2015.

VIEIRA, L. S.. Manual da ciência do solo; com ênfase aos solos tropicais. São paulo, SP (Brazil). Editora Agronômica Ceres Ltda. 2. ed.. 464 p. 1988.

WANG, G.. Adaptation of Asia-Pacific forests to climate change. *Journal Of Forestry Research*. Springer Nature. [s.l.], v. 27, n. 3, p.469-48818 março 2016.

**CAPÍTULO 1: A reidratação imediata restaura o aparelho fotossintético
de *Schinus terebinthifolia* (Raddi) após seca severa no solo?**

Can rehydration restore the photosynthetic apparatus of *Schinus terebinthifolia* (Raddi) after severe soil drought?

Abstract

Schinus terebinthifolia Raddi is a pioneer tree species that is native to the Atlantic forest of South America, but which is considered invasive in some places. Information on the tolerance of this species to stress conditions, in particular low water availability, is still limited. Therefore, the aim of this study was to evaluate the physiological and enzymatic metabolism adjustments in response to water deficit in *S. terebinthifolia*. The experiment was conducted in a greenhouse in a completely randomized distribution with two water treatments: [a] a control, and; [b] a suspended irrigation treatment, which continued until photosynthesis values approached zero after which the plants were rehydrated. We analyzed gas exchange, chlorophyll fluorescence and osmoregulatory compounds such as carbohydrates, amino acids, proline, ureids, nitrate. After seven days without irrigation photosynthesis was close to zero. At this time there was an increase in amino acid and proline content in the leaves, which prevented the carbohydrates and photosynthetic apparatus from being directly affected by the lack of water. We also observed stomatal closure and preservation of chloroplast pigments. The plants recovered gas exchange capabilities five days after rehydration. Young plants of this species tolerated severe drought with a fast recovery capacity, further supporting the claim that this species is very suitable for restoring forests and colonizing new environments.

Key words: Intrinsic water use efficiency. Leaf gas exchange. Stomatal conductance. Forest restoration. Antioxidant enzyme. Water deficit.

Introduction

The Atlantic Forest is one of the most degraded biomes in Brazil, having been heavily exploited since the arrival of the Portuguese in the mid 1500's (Zamith *et al.* 2006). One of the first steps to ensure a successful restoration of the remaining forest fragments is to identify suitable species for this purpose. The vegetation of the Atlantic forest has been classified into various sub-types, including dense ombrophylous forest, open ombrophylous forest, semideciduous seasonal forest, among others. Of these, the open forest is probably the most degraded by human actions (IBGE 2006) and is in urgent need of restoration. However, such initiatives require highly resilient species that are able to establish in stressful environments characterized by poor soil quality and variable precipitation. Such resistant species are especially valuable for restoration since their establishment can create valuable microhabitats for more sensitive species to establish (Mielke *et al.* 2005).

Degraded habitats are often limited in water and soil nutrients, both of which are essential for plant growth (Cabral *et al.* 2004), osmotic, physiological regulation and photosynthesis (Caron *et al.* 2014). The main effects of water stress on plant metabolism is loss of cellular turgor and decreasing in stomatal opening. These changes, although contributing to water use efficiency, end up limiting the assimilation of carbon and, consequently, the growth of the plant (Trovão *et al.* 2007). Water deficit also causes a decrease in transpiration, which is largely responsible for the absorption and transport of essential nutrients present in the soil as nitrate. The activity of the nitrate reductase enzyme, regulated by the presence of nitrate, is also quite sensitive to water deficiency. Nitrates are required for the synthesis of biomolecules such as nucleic acids, amino acids, and proteins (Kóvacs 2005; Firmano *et al.* 2009), meaning that all plant metabolism is compromised when water is scarce in the soil. Effective species for restoration should therefore be resilient to the adverse conditions found in degraded environments (Volpato *et al.* 2013; Rorato *et al.* 2018; Chiochetta *et al.* 2017).

Schinus terebinthifolia (the Brazilian pepper tree) is a good potential candidate for use in restoration projects. It is a pioneer species that is native to the Atlantic forest, but which has also become established in several regions of South America and most of the neotropical region of the globe (Jones *et al.* 1997; Silva *et al.*

2008). *S. terebinthifolia* thrives in anthropogenic environments (Pratt *et al.* 2006) and has already been identified as having great potential for the restoration of degraded areas (Coutin *et al.* 2017; Rorato *et al.* 2018). However, its ecophysiological response to these environments has not been fully investigated, and its mechanisms of drought tolerance are not well understood.

Current knowledge of this species indicates that its hydraulic characteristics make it more tolerant to water stress compared to native species of environments that it has invaded (Pratt *et al.* 2006). Moreover, it is known that the antioxidant and photoprotective apparatus of this type of pioneer species in tropical regions can be very efficient (Coutin *et al.* 2016). This suggests that its water and saline resistance may be linked to its photosynthetic and photochemical characteristics (Ewe *et al.* 2003; Ewe *et al.* 2005; Ewe *et al.* 2007; Mielke *et al.* 2005). The aim of the current study was to provide a detailed description of the adaptive response of *S. terebinthifolia* to water stress and rehydratation, including gas exchange, accumulation of organic compounds and photochemical efficiency. This information can be used to identify the main adaptive mechanisms used by this species to overcome water stress, clarifying and complementing existing physiological knowledge of this species.

Material and methods

Plant material and experimental conditions

This study was carried out in the Agrarian Sciences Center of the Federal University of Alagoas (UFAL), Rio Largo - AL ($9^{\circ} 28' 00''$ S, $35^{\circ} 49' 34''$ W, at 131 m elevation). The experiment was conducted in a greenhouse, using seedlings of *S. terebinthifolia* at approximately one year of age. The seedlings were transplanted to 20 L plastic pots filled with 20 kg of sieved soil, collected in a forested area ($9^{\circ} 27' 58.4''$ S, $35^{\circ} 49' 31.7''$ W). The experimental design was completely randomized, consisting of two treatments with eight replicates each.

The treatments were composed of two water regimes: [1] control (irrigated), consisting of plants irrigated daily, and; [2] water stressed, composed of plants submitted to water deficit by total suspension of irrigation and subsequent rehydration. Irrigation was suspended until the photosynthetic rate reached values close to $0 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ (the “maximum stress”), later the plants were rehydrated until the photosynthetic rates were equal to those of the control treatment. Physiological and biochemical analyses were performed at the end of each period. The water deficit lasted seven days and rehydration lasted five days.

The sediment used in the experiment was classified according to the Brazilian Soil Classification System (SBCS). A 0.5 kg sample of sediment was used in the evaluation of physical and chemical properties using: (1) clay (densimeter); (2) fine and coarse sand (sieving); (3) clay dispersed in water (densimeter); (4) degree of flocculation (GF). Nutritional analyses were performed using: (1) Mehlich-1 extractor; (2) 1.0 M KCl extractor; (3) Calcium acetate extractor at pH 7.0; (4) Welkley-Black method; (5) Base saturation; (6) Aluminum saturation. The soil of the experiment was classified as sandy clay loam: fine sand (320 g / kg), coarse sand (150 g / kg), clay (300 g / kg) and silt (230 g / kg). The soil had pH in water of 4.8 and contained sodium 10 (mg / dm³), phosphorus 39 (mg / dm³), potassium 200 mg / dm³), calcium 2.78 (mg / dm³), magnesium 1.52 (mg / dm³) and aluminum 0.38 (mg / dm³).

At the end of each experimental period, the soil water potential (MPa) was measured using WP4C Water Potential Meters (Decagon Devices) and the humidity (%) with a portable moisture meter model SM200, coupled to a model HH2 sensor. Temperature (Fig. 1A), relative air humidity (Fig. 1B), solar radiation (Fig. 1C) were recorded every 30 minutes by a meteorological station (Weather Station model WS -

GP1 / AT DELTA - T Devices, Cambridge - England) located within the greenhouse. From this data, the Vapor Pressure Deficit (DPV) was calculated based on Allen (1998), resulting in a value corresponding to the evaporative pressure of the environment towards the atmosphere, expressed in KPa (Fig. 1D).




Fig. 1 Average hourly values of air temperature ($^{\circ}\text{C}$), relative air humidity (%) and vapor pressure deficit (Kpa), recorded every thirty minutes and during twelve days of the experiment.

Leaf water potential

Leaf water potential (Ψ_w – MPa) was determined using a Scholander-type pressure chamber (Soil Moisture Equipment, Santa Barbara, CA, USA) (Scholander *et al.* 1965), in the periods of the pre-dawn (05:00 am) and midday (12:00 pm) on the third leaf from the apex. The pressure reading was performed according to the methodology proposed by Turner (1981).

Quantum Efficiency of Photosystem II

The quantum efficiency of PSII was performed with a portable fluorometer (PAM - 2500, Handy PEA, Hansatech, Norfolk, UK). The maximum quantum efficiency of the PSII represented by the Fv/Fm ratio was evaluated in leaves adapted to the dark for 20 minutes by means of leaf forceps, in the period of the pre-dawn (04:00 – 05:30 am) and at noon (12:00 – 12:30 pm) (Maxwell & Johnson 2000). The effective quantum efficiency of PSII (Φ_{PSII}) was determined on the same leaves at noon (11:00 - 12:30 pm) (Baker 2008).

Gas exchange and water use efficiency

The rates of photosynthesis ($A - \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$), transpiration ($E - \text{mmol H}_2\text{O m}^{-2} \text{ s}^{-1}$) and stomatal conductance to water vapor ($gs - \text{mol H}_2\text{O m}^{-2} \text{ s}^{-1}$) were measured on mature leaves and completely expanded between 8:00 and 11:00 a.m. using an infrared gas analyzer (IRGA) (LI-6400 Li-color, USA) equipped with a 6400-02B Red Blue artificial light source, set at $1200 \mu\text{mol m}^{-2} \text{ s}^{-1}$ of luminous intensity. The instantaneous water use efficiency (iWUE) and intrinsic water use efficiency (IWUE) were calculated by the A/E and A/gs ratios, respectively, and expressed in $\mu\text{mol CO}_2 \text{ mmol}^{-1} \text{ H}_2\text{O}$.

Contents of photosynthetic pigments and SPAD Index

The extraction of the photosynthetic pigments was done according to the method of Arnon (1949). Approximately 50 mg of fresh leaves were placed into test tubes protected with aluminum foil containing 5 mL of 80% acetone. After 48 hours, the absorbance was quantified at wavelength of 663, 645 and 480 nm using a spectrophotometer. The values of chlorophyll *a*, *b*, total and carotenoids were calculated following Hendry & Grime (1993), and subsequently the chlorophyll *a* / *b* ratio was calculated. The SPAD index was determined by the non-destructive method using a portable chlorophyllometer (SPAD-502 - Minolta, Japan). Each sample was quantified using the average of 5 measurements performed on different leaves.

Biochemical analyses

Soluble carbohydrates were extracted and quantified according to the methodology of Dubois *et al.* (1956). The readings were performed in a spectrophotometer at 490 nm and the absorbance compared to the standard curve of glucose concentration (μg), values were expressed in $\text{mg g}^{-1} \text{ FW}$. The free proline was extracted and quantified based on the methodology described by Bates *et al.* (1973). The reading was performed in a spectrophotometer at 520 nm and the absorbances compared to the standard proline curve, values were expressed in $\mu\text{mol g}^{-1} \text{ FW}$.

Nitrogen compounds (nitrate, ureids and free amino acids) were extracted from 500 mg of fresh plant material in 5 mL of MCW (60% methanol: 25% chloroform: 15% H_2O) according to methodology of Bielecki & Turner (1966). The nitrate was quantified according to the methodology of Cataldo *et al.* (1975), the readings were performed in a spectrophotometer at 410 nm and the absorbance compared to the standard NaNO_3

curve. Ureids (total ureids, allantoin and alantoic Acid) were quantified according to the methodology of Vogels & Van Der Drift (1970). The readings were performed in a spectrophotometer at 535 nm and the absorbances compared to the standard allantoin curve. Total free amino acids were determined according to Yemm & Cocking (1995). The reading was performed in a spectrophotometer at 570 nm and the absorbance compared to the standard curve of amino acids (pool of amino acids - arginine, glycine, glutamic acid and phenylalanine), values were expressed in $\mu\text{mol g}^{-1}$ FW.

Statistical analyses

The data were submitted to analysis of variance (ANOVA) and the means were compared by the F test at 5% probability ($P<0.05$), using the statistical program SISVAR version 5.6 build 86.

Results

Soil of plants in the control treatment, retained an average of 30% of moisture and -0.5 MPa of water potential during the experiment (Fig. 2). At the day of maximum stress for the treatment group, soil moisture was approximately 5% and the water potential was -5.0 MPa. Both the moisture and the soil water potential increased after rehydration, reaching values close to the soil evaluated in the control plants. Plants reached maximum stress at the seventh day of experiment, and after rehydration, photosynthesis rates reached values close to irrigated (control) plants on the twelfth day.




Fig. 2 (A) Soil water potential and (B) Soil moisture during the experiment.

When photosynthesis rates reached values close to zero (day of maximum stress) (Fig. 3A) we observed decreases in stomatal conductance (Fig. 3B), transpiration (Fig. 3C) and carboxylation efficiency (Fig. 3D) - direct consequences of the reduction in water availability. The values of these variables in the control treatment maintained a constant range throughout the experiment. After rehydration, stressed plants re-established pre-treatment photosynthetic rates of around $23 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$, indicating that this species can completely recover after a period of severe water stress.

The values of the instantaneous water-use efficiency (iWUE) (Fig. 3E) and the intrinsic water-use efficiency (IWUE) (Fig. 3F) were higher in control plants until the day of maximum stress, but after rehydration, stress treatment plants exceeded the values of the control treatment plants by optimizing water use soon after the severe water deficit.




Fig. 3 Gaseous exchanges of *Schinus terebinthifolia* Raddi accompanied during twelve days of experiment: (A) photosynthesis, (B) stomatal conductance, (C) transpiration, (D) carboxylation efficiency, (E) instantaneous water use efficiency and (F) intrinsic water use efficiency.

Pre-treatment leaf water potential was around -0.3 MPa at pre-dawn and at midday (data not shown). After seven days without watering, verified leaf potentials were approximately -1.2 MPa in the stress treatment plants and -0.4 MPa in the control treatment at pre-dawn (Fig. 4A). However, at midday reduced air humidity and increased temperatures led to high values of DPV and a considerable decline in leaf water potential. The capacity of osmotic adjustment of *S. terebinthifolia* was approximately -3.6 MPa and -4.2 MPa for the stress and control treatments, respectively (Fig. 4B). After rehydration, irrigated plants had water potential of around -0.4 MPa, while stress treatment plants had values of -0.3 MPa at pre-dawn, and -1.2 MPa at midday (Fig. 4A-B).




Fig. 4 Values of leaf water potential in *Schinus terebinthifolia* Raddi plants were verified seventh and twelfth day of the experiment at two times of day: (A) beforehand and (B) Midday.

Carbohydrate content was similar in leaves of both control and treatments plants (Fig. 5A). After rehydration a considerable increase in the carbohydrate content was observed on the twelfth day of the experiment. Amino acid levels increased in the drought stressed plants ($24.4 \mu\text{mol g}^{-1}$ FW at maximum stress), while control plants had values of approximately $3.5 \mu\text{mol g}^{-1}$ FW (Fig. 5B). Amino acids accumulate in stress situations, acting as osmotic regulators of the cells and adjusting the water status of the plants. After rehydration the content of these compounds decreased in the stress treatment to approximately $14.7 \mu\text{mol g}^{-1}$ FW. The control plants had an amino acid content of approximately $11.7 \mu\text{mol g}^{-1}$ FW. This result suggest that they

had an increase in amino acids after the seventh day of the experiment, which may be a consequence of atmospheric variables, since plants of the control treatment were kept hydrated throughout the experiment period. We observed a peak in proline in drought-stressed plants ($202.3 \mu\text{mol g}^{-1} \text{ FW}$) and a drastic reduction of this value ($9.1 \mu\text{mol g}^{-1} \text{ FW}$) in the same plants after rehydration. The control plants maintained a range between 22 and $28 \mu\text{mol g}^{-1} \text{ FW}$ proline on the seventh day and twelfth day of the experiment, respectively (Fig. 5C).




Fig. 5 (A) Quantification of soluble carbohydrates, (B) free amino acid content and (C) proline in leaves of *Schinus terebinthifolia* Raddi plants verified on the seventh and twelfth day of experiment.

Levels of allantoic acid did not change at the maximum stress point. After rehydration (on the twelfth day) we noticed a considerable increase in its levels in the stress treatment, from $1.9 \mu\text{mol g}^{-1} \text{ FW}$ (Fig. 6A), whereas it remained at higher levels in control plants throughout the experiment ($12 \mu\text{mol g}^{-1} \text{ FW}$ in 7^o day, and $9.2 \mu\text{mol g}^{-1} \text{ FW}$ in the 12^o day). Reductions in allantoin content were verified in stress

treatment after rehydration, decreasing from $9.8 \text{ } \mu\text{mol g}^{-1} \text{ FW}$, recorded on the maximum stress day to $5.7 \text{ } \mu\text{mol g}^{-1} \text{ FW}$ (Fig. 6B). This result suggests that water stress increases the nitrogen requirement during rehydration (Fig. 6C). Additionally, stressed plants had increased nitrate levels ($0.11 \text{ } \mu\text{mol g}^{-1} \text{ FW}$) on the day of maximum stress compared to control plants ($0.09 \text{ } \mu\text{mol g}^{-1} \text{ FW}$) on the same day (Fig. 6D). After rehydration, a decrease in nitrate levels (to $0.07 \text{ } \mu\text{mol g}^{-1} \text{ FW}$) in stressed plants was observed, with as the control plants maintain the same levels observed on the day of maximum stress.




Fig. 6 Ureide content in leaves of *Schinus terebinthifolia* Raddi plants on the seventh and twelfth day of experiment: (A) allantoic acid, (B) allantoin and (C) total ureides and (D) nitrate.

Stressed plants had chlorophyll *a* at values close to those found in control, both during maximum stress and after rehydration (Tab. 1.). The SPAD index did not markedly change during the twelve days of the experiment, though there was a slight increase in the stress treatment after rehydration (Tab. 1.).

Table 1. Index SPAD, total chlorophyll (Chlo *T*), chlorophyll *a* (Chlo *a*), chlorophyll *b* (Chlo *b*), carotenoids (Carot) and Chlorophyll *a/b* (Chlo *a/b*) at maximum stress. Means followed by the same lowercase letter in the columns do not differ significantly from each other by the Tukey test ($P < 0.05$).

Treatment	SPAD	Chlo <i>T</i> µg/L	Chlo <i>a</i> µg/L	Chlo <i>b</i> µg/L	Carot µg/L	Chlo <i>a/b</i> µg/L
Control	53.1a	1.64a	1.72A	0.45a	0.72a	4.84a
Dry	50.1a	2.18a	1.38a	0.25a	0.91a	6.81A
CV	9.44	26.8	18.9	8.4	21.6	10.08

Apparently, chlorophyll was not severely impaired by the severe (but short duration) water stress. It has been observed that values below 0.725 F_v/F_m , may cause leaves to suffer photoinhibitory damage. Drought stress reduced quantum efficiency at pre-dawn and midday (Fig. 7). Specifically, on the day of maximum stress F_v/F_m decreased to 0.709 in stressed plants (Fig. 7A), but increased to 0.754 after rehydration (Fig. 7B). Control plants showed little variation in quantum efficiency during the experiment. After rehydration, stressed plants quickly restored the electron transport efficiency of their photochemical apparatus (Fig. 7C).




Fig. 7 Photochemical efficiency of *Schinus terebinthifolia* Raddi observed at the first, seventh and twelfth day of experiment: (A) Maximum quantum yield (F_v/F_m) evaluated at 4:00 am; (B) Maximum quantum production (F_v/F_m) evaluated at 12:00 o'clock; (C) Effective quantum efficiency of photosystem II.

Discussion

As expected, gas exchange was negatively affected in stressed plants. Studies using experimental drought or performed in the dry season usually have this type of result, with plants decreasing the water lost from transpiration mainly through reduction of the stomatal aperture (Mariano *et al.* 2009; Marenco *et al.* 2014; Albuquerque *et al.* 2013; Costa *et al.* 2015). These findings suggest that the photosynthetic apparatus of *S. terebinthifolia* may be the main reason for its observed success in invading multiple environments, showing characteristics similar to that of plants of seasonally dry environments. Ewe (2003) studied *S. terebinthifolia* on a plateau at different times of the year in South Florida, observing higher mesophilic conductivity, higher intrinsic water use efficiency and greater use of nitrogen than most native species. It also had the highest gas exchange rates, indicating that this species has an important physiological advantage for invading new environments.

Other variables directly affected by stomatal aperture reduction are photosynthetic and transpiratory rates, which commonly show decreases as stress becomes severe. This balance between stomatal opening and CO₂ capture directly reflects the amount of water used by plants for key physiological processes, since the reduction of stomatal openings prevents water loss, but also reduces CO₂ entry and assimilation (Ai *et al.* 1990). The instantaneous efficiency of water use and the intrinsic efficiency of water use evaluate the ability of a plant to perform photosynthesis in the face of water scarcity, which consequently impacts on biomass production (Souza *et al.* 2010). In our study, stressed plants showed higher values of instantaneous efficiency of water use than control plants in the days prior to maximum stress, when photosynthesis was being carried out with the minimum amount of water that could be lost.

Water deficit also negatively affected the electron transfer efficiency of the photosystems. As observed, the photoinhibitory damage caused to photosystem II during the stress period was reversible (Critchley *et al.* 1995; Hasan *et al.* 2018; Maxwell *et al.* 2000; Araújo *et al.* 2004). This is typically seemed in species of seasonally dry forests that have photoprotective mechanisms (Trovão *et al.* 2007). These species have near-maximum quantum efficiency, even during periods of water stress, and can return to their maximum yield shortly after rehydration (Björkman *et al.* 1987). *S. terebinthifolia* has the plasticity of a pioneer species that grows on the edges

of the forest (Rabelo 2013). The maximum photosystem II quantum yield (F_v/F_m) and the photosynthetic maximum photochemical yield of photosystem II (F_v/F_0) also showed the same pattern of response to drought stress. This is also a consequence of the decreased in the amount of water in the plant organs, which could dissipate excess of heat through transpiration, and because of the stomatal closure. In addition, due to the inhibition of CO_2 entry (also caused by the stomatal closure) the electron transport chain of photosynthesis is disrupted and the contribution of NADP^+ to the photosystem I is impaired, producing the superoxide anion ($\text{O}_2\cdot^-$) molecule that highly toxic to plant cells (Martins *et al.* 2018). This is caused by the reduction in the plant's capacity to transport electrons through PSII, resulting from photoinhibition.

Our results suggest that *S. terebinthifolia* is highly resistant to periods of water scarcity and able to quickly overcome damages suffered due to lack of water in the metabolism. In this sense it belongs to a group of species that can successfully establish themselves in the midst of water scarcity (Sheil *et al.* 2018). The reduction of leaf Ψ_w verified in plants under stress reflect the effect of water restriction on their vegetable water status (Schimpl *et al.* 2019). The ability to reduce leaf Ψ_w in stressed plants indicates that this species has one of the main drought tolerant traits - a strategy to maintain the water potential gradients necessary for the plant to absorb water even in low humidity soils (Silva *et al.* 2016; Albrecht *et al.* 2006). These osmotic changes can be seen in the leaves, as has been observed in "aroeira" and also in leaves and fruits of other species in other studies (Fernandes *et al.* 2018; Falahi *et al.* 2018). Due to atmospheric variables, that are extremely different between the pre-dawn and midday hours, these responses may vary between species. Water deficit has been observed to induce lower leaf water potentials and has a significant effect on the gas exchange parameters (Sellin 2014; Silva-Braco 2017), which was also verified here.

Water stress can cause organic compounds that have nitrogen as feedstock to be degraded or for their synthesis to be drastically reduced (Golçalves *et al.* 2012). The increase in proline concentration is due to the reduction of protein synthesis and the action of proteases usually occurs in stressed plants (Stewart *et al.* 1973), which also explains the observed increase in amino acid content. Our results indicated that soluble carbohydrates remained stable during water stress. However, when rehydrated, there was a peak in sugar production, probably caused by the resumption of CO_2 assimilation and starch synthesis (Santos *et al.* 1988). The increase in soluble sugars concentration probably occurred due to the decrease in starch synthesis during

stress, and also through hydrolysis of the starch reserves to generate ATP. Some researchers have noted an increase in carbohydrates during water stress, linking this to osmotic adjustment in the cells (Filho *et al.* 2001; Costa *et al.* 2015; Mendes *et al.* 2017).

Osmotic adjustment in *S. terebinthifolia* may have been performed by other solutes, such as nitrate, total amino acids and especially proline, with a marked increase in stress and a decrease in rehydration. The products of protein lysis are proline, and other amino acids, which act as osmoprotectants and serve as substrate for the synthesis of more proline (Falahi *et al.* 2018; Nounjan *et al.* 2018; Lee *et al.* 2013). However, during rehydration the proline acts as a reservoir of nitrogen and a carbon source for post-stress growth, resulting in a pronounced drop in proline content and a marked increase in synthesized carbohydrates after the resumption of carboxylation using the haptonic skeleton of these compounds (Hossain *et al.* 2008).

Nitrate may have accumulated as a result of changes in nitrogen metabolism, since nitrate reductase enzyme is inhibited by drought stress (Shane *et al.* 1967; Marur *et al.* 2000; Omena-Garcia *et al.* 2015). As no nitrate toxicity in cells was observed in this study, the nitrate probably influenced the osmotic adjustment of the cells (Ariz *et al.* 2010; Beur *et al.* 1992; Beninni *et al.* 2002). Such responses activate the degradation pathways of biomolecules and cause the synthesis and accumulation of other organic solutes, such as nitrate, and compounds from the urea group. We observed that stressed plants had allantoic acid values close to the values found in well irrigated plants but that these ureids peaked after rehydration. This result does not support the hypothesis of accumulation of these compounds during water stress, as has been observed in legumes (King 2005). Allantoin and allantoic acid - considered the main forms of nitrogen reserve in tropical tree plants and better known in the family Fabeaceae (Kabbadj *et al.* 2017) - had a decreasing trend until the end of the experiment. This suggests that the decrease in the assimilation of these compounds is inversely proportional to water stress in *S. terebinthifolia*. It is well known that chlorophyll content responds to shading, soil fertility, nutritional status of the plant and to the water regime or with these factors in combination (Rego *et al.* 2006; Nogueira *et al.* 2001; Bayoumi *et al.* 2013), decreasing according to the level of stress. Salla (2007) demonstrated that total chlorophyll content is strongly correlated to SPAD index values using data from several tree species. Both these measures can estimate the loss of photosynthetic pigments in plants of *S. terebinthifolia* submitted to water deficit,

with a significant decrease in total chlorophyll (Rabelo *et al.* 2013). One of the probable reasons for such variation is that the nitrogen metabolism, which is involved in the initial stages of chlorophyll production, is directly impaired by lack of water.

S. terebinthifolia is considered a resilient species in tropical environments. Our study suggests that young plants of this species are highly tolerant to water stress with a rapid and total recovery from short term severe stress. Water stress tolerance appears to be largely mediated through regulation of osmotic potential with associated increases amino acids and proline, ‘saving’ carbohydrates for the maintenance of biomass. *S. terebinthifolia* protects its photosynthetic apparatus by regulating stomatal closure, which also prevents the degradation of chloroplast pigments. Despite making a full physiological recovery after rehydration, there were many changes in its metabolism under water stress.

References

- Ai Z, Zhang J, Liu H, Xin Q, Xue S, Liu G. 2017. Soil nutrients influence the photosynthesis and biomass in invasive *Panicum virgatum* on the Loess Plateau in China. *Plant Soil* 418: 153–164. doi: 10.1007/s11104-017-3286-x.
- Albuquerque MPF, Moraes FKC, Santos RIN, Castro GLS, Ramos EMLS, Pinheiro HA. 2013. Ecofisiologia de plantas jovens de mogno-africano submetidas a déficit hídrico e reidratação. *Pesquisa Agropecuária Brasileira* 48: 9-16. doi: 10.1590/S0100-204X2013000100002.
- Allen RG, Pereira LS, Raes D, Smith M. 1998. Crop evapotranspiration - Guidelines for computing crop water requirements. FAO Irrigation and Drainage Papers 56: 326 p.
- Araújo SAC, Deminicis BD. 2009. Fotoinibição da Fotossíntese. *Brazilian Journal of Biosciences* 7: 463-472.
- Ariz I, Esteban R, García-Plazaola JI, Becerril JM, Aparicio-Tejo PM, Moran JF. 2010. *Journal of Plant Physiology* 167: 1038–1045. doi: 10.1016/j.jplph.2010.02.014.
- Arnon DI. 1949. Copper enzymes in isolated chloroplasts. Polyphenoloxidase in Beta vulgaris. *Plant physiology*. 24: 1-15.
- Baker, NR. 2008. Chlorophyll Fluorescence: A Probe of Photosynthesis In Vivo. *Annual Review of Plant Biology*. 59: 89-113. doi: 10.1146/annurev.arplant.59.032607.092759.
- Bates LS, Waldren RP, Teare ID. 1973. Rapid determination of free proline for water-stress studies. *Plant Soil* 39:205–207. doi: 10.1007/BF00018060.
- Bayoumi TY, Amal M, EL-Mageed A *et al.* 2015. The Impact of Drought Stress on some Morpho-Physiological Traits and RAPD Markers in Wheat Genotypes. *Journal of Plant Production Sciences* 4: 27-37. doi: 10.21608/jpps.2015.7396.

Behr U; Wlebe HJ. 1992. Relation between photosynthesis and nitrate content of lettuce cultivars. *Scientia Horticulturae* 49: 175-179. doi: 10.1016/0304-4238(92)90155-6.

Beninni ERY, Takahashi HW, Neves CSVJ, Fonseca, ICB. 2002. Teor de nitrato em alface cultivada em sistemas hidropônico e convencional. *Horticultura Brasileira* 20:183-186. doi: 10.1590/S0102-05362002000200013.

Bielecki RL, Turner NA. 1966. Analytical Separation and estimation of amino acids in crude plant extracts by thin-layer electrophoresis and chromatography. *Biochemistry* 17:278-293. doi: 10.1016/0003-2697(66)90206-5

Björkman O, Demmig B. 1987. Photon yield of O₂ evolution and chlorophyll fluorescence characteristics at 77K among vascular plants of diverse origins. *Planta* 170: 489-504. doi: 10.1007/BF00402983.

Branco, MCS. 2017. Influence of low light intensity and soil flooding on cacao physiology. *Scientia Horticulturae* 217: 243-257. doi: 10.1016/j.scienta.2017.01.038.

Cabral EC, Barbosa DCA, Simabukuro EA. 2004. Crescimento de plantas jovens de *Tabebuia aurea* (Manso) Benth. & Hook. f. ex S. Moore submetidas a estresse hídrico. *Acta Botanica Brasilica* 18: 241-251. doi: 10.1590/S0102-33062004000200004.

Caron BO, Perrando ER, Schmidt D et al. 2014. Relações fisiológicas em mudas de pata-de-vaca (*Bauhinia forficata* Link). *Revista Brasileira de Plantas Medicinais*. 16:196-201. doi: 10.1590/S1516-05722014000200005.

Castro DS, Santos AO, Lobato AKS et al. 2007. Concentrações de Prolina e Carboidratos Solúveis Totais em Folhas Teca (*Tectona grandis* L.f) Submetida ao Estresse Hídrico. *Revista Brasileira de Biociências* 5: 921-923.

Cataldo DA, Haroon M, Schradev LE, Youngs VL. 1975. Rapid colorimetric determination of nitrate in plant tissue by nitration of salicylic acid. *Communications in Soil Science and Plant Analysis*, 6: 71-80. doi: 10.1080/00103627509366547.

Chiochetta AG, Toumi H, Böhm RFS et al. 2017. Use of phytoproductivity data in the choice of native plant species to restore a degraded coal mining site amended with a

stabilized industrial organic sludge. Environmental Science and Pollution Research 24: 24624-24633. doi: 10.1007/s11356-017-0128-6.

Contin DR, Munné-Bosch S. 2016. Interspecific variation in vitamin E levels and the extent of lipid peroxidation in pioneer and non-pioneer species used in tropical forest restoration. Tree Physiology 36: 1151-1161. doi:10.1093/treephys/tpw018.

Costa AS, Freire ALO, Bakke IA, Pereira FHF. 2015. Respostas fisiológicas e bioquímicas de plantas de aroeira (*Myracrodruon urundeuva* allemão) ao déficit hídrico e posterior recuperação. Irriga 20: 705-717.

Critchley C. 1998. Photoinhibition. In: Raghavendra, AS (Ed.). Photosynthesis: A comprehensive treatise. Cambridge, Cambridge University Press. p. 264-272.

Dubois M, Gilles KA, Hamilton JK, Rebers PA, Smith F. 1956. Colorimetric method for determination of sugars and related substances. Analytical Chemistry 28: 350-356. doi: 10.1021/ac60111a017.

Ewe SML, Sternberg LSL. 2003. Seasonal gas exchange characteristics of *Schinus terebinthifolia* in a native and disturbed upland community in Everglades National Park, Florida. Forest Ecology and Management 179: 27-36. doi: 10.1016/S0378-1127(02)00531-5.

Ewe SML, Sternberg LSL. 2005. Growth and gas exchange responses of Brazilian pepper (*Schinus terebinthifolia*) and native South Florida species to salinity. Trees 19: 119-128. doi: 10.1007/s00468-004-0370-7.

Ewe SML, Sternberg LSL. 2007. Water Uptake Patterns of an Invasive Exotic Plant in Coastal Saline Habitats. Journal of Coastal Research 23: 255-264. doi: 10.2112/1551-5036.

Ewe SML. 2001. Ecophysiology of *Schinus terebinthifolia* contrasted with native species in two South Florida ecosystems. PhD Thesis, University of Miami, Miami. SERC Dissertations & Theses (Working copies).

Ewel JJ, Ojima DS, Karl DA, Debusk WF. 1982. *Schinus* in successional ecosystems of Everglades National Park. South Florida Research Center Report T-676.

Fernandes RDM, Cuevas MV, Diaz-Espejo A, Hernandez-Santana V. 2018. Effects of water stress on fruit growth and water relations between fruits and leaves in a hedgerow olive orchard. Agricultural Water Management 210: 32-40. doi: 10.1016/j.agwat.2018.07.028.

Falahi H, Sharifi M, Ahmadian CN, Zare MH. 2018. Water stress alleviation by polyamines and phenolic compounds in *Scrophularia striata* is mediated by NO and H₂O₂. Plant Physiology and Biochemistry 130: 139-147. doi: 10.1016/j.plaphy.2018.07.004.

Hossain Z, Mandal AKA, Datta SK. 2008. Development of NaCl-tolerant line in *Chrysanthemum morifolium* Ramat. through induced somaclonal variation. Journal of Biotechnology 136: 160-167. doi: 10.1016/j.jbiotec.2008.07.343.

Filho JTC, Stacciarini-Seraphin E. 2001. Alteração no potencial osmótico e teor de carboidratos solúveis em plantas jovens de lobeira (*Solanum lycocarpum* St.-Hil.) em resposta ao estresse hídrico. Brazilian Journal of Botany 24: 199-204. doi: 10.1590/S0100-84042001000200010.

Gonçalves JFC, Silva CEM, Guimarães DG. 2009. Fotossíntese e potencial hídrico foliar de plantas jovens de andiroba submetidas à deficiência hídrica e à reidratação. Pesquisa Agropecuaria Brasileira 44: 8-14. doi: 10.1590/S0100-204X2009000100002.

Gonçalves JFC, Melo EGF, Silva CEM, Ferreira JM, Justino GC. 2012. Estratégias no uso da energia luminosa por plantas jovens de *Genipa spruceana* Steyermark submetidas ao alagamento. Acta Botanica Brasilica. 26:391-398. doi: 10.1590/S0102-33062012000200014.

Hasan MM, Ma F, Prodhan ZH et al. 2018. Molecular and Physio-Biochemical Characterization of Cotton Species for Assessing Drought Stress Tolerance. International Journal of Molecular Sciences 19: 2636. doi: 10.3390/ijms19092636.

Jones D. 1997. Biology of Brazilian pepper. In: Ferriter A (Ed). Brazilian pepper management plan for Florida. Florida, Florida Exotic Pest Plant Council.

- Kabbadj A, Makoudi B, Mohammed M, Pauly N, Frendo P, Ghoulam C. 2017. Physiological and biochemical responses involved in water deficit tolerance of nitrogen-fixing *Vicia faba*. Plos One 12: e0190284. doi: 10.1371/journal.pone.0190284.
- King CA, Purcell LC. 2005. Inhibition of N₂ fixation in soybean is associated with elevated ureids and amino acids. Plant Physiology 137: 1389-1396. doi: 10.1104/pp.104.056317.
- Kouali A, Mouradi M, Latrach L et al. 2017. Study of effects of seawater salt spray on growth, chlorophyll fluorescence and chlorophyll content in three coastal species of Morocco. Journal of Materials and Environmental Sciences 8: 4385-4390.
- Kovács GJ. 2005. Modelling of adaptation processes of crops to water and nitrogen stress. Physics and Chemistry of the Earth 30: 209-216. doi: 10.1016/j.pce.2004.08.022.
- Lee BR, Muneer S, Park SH, Zhang Q, Kim TH. 2013. Ammonium- induced proline and sucrose accumulation, and their significance in antioxidative activity and osmotic adjustment. Acta Physiologiae Plantarum, 35: 2655-2664. doi: 10.1007/s11738-013-1297-7.
- Marenco RA, Antezana-Vera SA, Gouvêa PRS, Camargo MAB, Oliveira MFO, Santos JKS. 2014. Fisiologia de espécies florestais da Amazônia: fotossíntese, respiração e relações hídricas. Revista Ceres 61: 786-789. doi: 10.1590/0034-737X201461000004.
- Mariano KR, Barreto LS, Silva AHB, Neiva GKP, Amorim S. 2009. Fotossíntese E tolerância protoplasmática foliar em *Myracrodruon urundeuva* FR. ALL. submetida ao déficit hídrico. Revista Caatinga (Mossoró) 22: 72-77.
- Marur CJ, Mazzafera P, Magalhães AC. 2000. Atividade da enzima redutase do nitrato em algodoeiro submetido ao déficit hídrico e posterior recuperação da turgescência. Scientia Agricola 57: 277-281. doi: 10.1590/S0103-90162000000200013.
- Maxwell K, Johnson GN. 2000. Chlorophyll fluorescence: a practical guide. Journal of Experimental Botany 51: 659–668. doi: 10.1093/jexbot/51.345.659.

Mendes FS, Castro DS, Neto CFO, Lobato AKS, Cunha RLM, Costa RCL. 2007. Níveis de Prolina e Carboidratos Solúveis Totais em Folhas de Mogno (*Swietenia macrophylla* King R.A) Induzidas ao Estresse Hídrico e a Reidratação. Revista Brasileira de Biociências 5: 939-941.

Mielke MS, Almeida AAF, Gomes FB, Mangabeira PAO, Silva DC. 2005. Effects of soil flooding on leaf gas exchange and growth of two neotropical pioneer tree species. New Forest 29: 161-168. doi: 10.1007/s11056-005-0247-7.

Nogueira RJMC, Moraes JAPV, Burit HA, Neto EB. 2001. Alterações na resistência à difusão de vapor das folhas e relações hídricas em aceroleiras submetidas a déficit de água. Revista Brasileira de Fisiologia Vegetal 13: 75-87. doi: 10.1590/S0103-31312001000100009.

Nounjan N, Chansongkrow P, Charoensawan V *et al.* 2018. High Performance of Photosynthesis and Osmotic Adjustment Are Associated With Salt Tolerance Ability in Rice Carrying Drought Tolerance QTL: Physiological and Co-expression Network Analysis. Frontiers in Plant Science 9: 1135. doi: 10.3389/fpls.2018.01135.

Omena-Garcia RP, Justino GC, Araújo VBF, Souza LAG, Camargos LS, Gonçalves JFC. 2014. Mineral Nitrogen Associated Changes in Growth and Xylem-N Compounds in Amazonian Legume Tree. Journal of Plant Nutrition 38: 584-595. doi: 10.1080/01904167.2014.957389

Pratt RB, Black RA. 2006. Do invasive trees have a hydraulic advantage over native trees?. Biological Invasions 8:1331–1341. doi: 10.1007/s10530-005-0422-y.

Rabelo GR, Vitória AP, Silva MVA *et al.* 2013. Structural and ecophysiological adaptations to forest gaps. Trees 27: 259-272. doi: 10.1007/s00468-012-0796-2.

Rego GM, Possamai, E. 2011. Efeito do Sombreamento sobre o Teor de Clorofila e Crescimento Inicial do Jequitibá-rosa. Pesquisa Florestal Brasileira 53: 179-194.

Rorato DG, Araujo MM, Tabaldi LA *et al.* 2018. Tolerance and resilience of forest species to frost in restoration planting in southern Brazil. Restoration Ecology 26: 537–542. doi: 10.1111/rec.12596.

- Salla L, Rodrigues JC, Marenco RA. 2007. Teores de clorofila em árvores tropicais determinados com o SPAD-502. Revista Brasileira de Biociências 5: 159-161.
- Santos RF, Carlesso ER. 1998. Déficit Hídrico e os Processos Morfológico e Fisiológico Das Plantas. Revista Brasileira de Engenharia Agrícola e Ambiental 2: 287-294. doi: 10.1590/1807-1929/agriambi.v2n3p287-294.
- Scholander PF, Hammel HT, Hemingsen EA, Bradstreet ED. 1965. Hydrostatic pressure and osmotic potential in leaves of mangroves and some other plants. Proceedings of the National Academy of Sciences of the United States of America 52: 119–125. doi:10.1073/pnas.52.1.119.
- Sellin A, Niglas A, Ōunapuu-Pikas E, Kupper P. 2014. Rapid and long-term effects of water deficit on gas exchange and hydraulic conductance of silver birch trees grown under varying atmospheric humidity. BMC Plant Biology 14: 72. doi: 10.1186/1471-2229-14-72.
- Shaner DL, Boyer JS. 1976. Nitrate Reductase Activity in Maize (*Zea mays L.*) Leaves: I. Regulation by Nitrate Flux. Plant Physiology 58: 499-504. doi: 10.1104/pp.58.4.499.
- Sheil D. 2018. Forests, atmospheric water and an uncertain future: the new biology of the global water cycle. Forest Ecosystems 5:19. doi: 10.1186/s40663-018-0138-y.
- Silva MAV, Nogueira RJMC, Oliveira AFM, Santos VF. 2008. Resposta estomática e produção de matéria seca em plantas jovens de aroeira submetidas a diferentes regimes hídricos. Revista Árvore. 32: 335-344. doi: 10.1590/S0100-67622008000200016.
- Silva MR, Klar AE, Passos JR. 2004. Efeitos do manejo hídrico e da aplicação de potássio nas características morfofisiológicas de mudas de *Eucalyptus Grandis W.* (Hill ex. Maiden). Irriga 9: 31-40. doi: 10.15809/irriga.2004v09n1p31-40.
- Silva-Pinheiro J, Lins L, Souza FC et al. 2016. Drought-stress tolerance in three semi-arid species used to recover logged areas. Brazilian Journal of Botany 39: 1031-1038. doi: 10.1007/s40415-016-0309-4.

- Souza BD, Meiado MV, Rodrigues BM; Santos MG. 2010. Water relations and chlorophyll fluorescence responses of two leguminous trees from the Caatinga to different watering regimes. *Acta Physiologiae Plantarum* 32: 235–244. doi: 10.1007/s11738-009-0394-0.
- Trovão DMBM, Fernandes PD, Andrade LA, Neto JD. 2007. Variações sazonais de aspectos fisiológicos de espécies da Caatinga. *Revista Brasileira de Engenharia Agrícola e Ambiental* 11: 307–311. doi: 10.1590/S1415-43662007000300010.
- Trovão DMBM, Fernandes PD, Andrade LA, Neto JD, Oliveira AB, Queiroz JA. 2004. Avaliação do potencial hídrico de espécies da Caatinga sob diferentes níveis de umidade no solo. *Revista de Biologia e Ciências da Terra* 4(2).
- Turner NC. 1981. Techniques and experimental approaches for the measurement of plant water status. *Plant and soil* 58: 339-366.
- Verbruggen N, Hermans C. 2008. Proline accumulation in plants: a review. *Amino Acids* 35: 753-759. doi: 10.1007/s00726-008-0061-6.
- Viégas LB, Simões D, Lima KB, Silva MR (2018) Water Management for *Schinus terebinthifolia* Raddi Seedlings in Degradable Containers. *Floresta e Ambiente* 25: e20160084. doi: 10.1590/2179-8087.008416.
- Vogels GD, Drift C. 1970. Differential analyses of glyoxylate derivates. *Analytical Biochemistry* 33: 143-157. doi: 10.1016/0003-2697(70)90448-3.
- Volpato GH, Martins SV. 2013. The bird community in naturally regenerating *Myracrodruon urundeuva* (Anacardiaceae) forest in Southeastern Brazil. *Revista de Biología Tropical* 61: 1585-1595.
- Yemm EW, Cocking EC. 1955. The determination of amino acids with ninhydrin. *Analyst* 80: 209-213. doi: 10.1039/AN9558000209.
- Zamith LR, Scarano FR. 2006. Restoration of a Restinga Sandy Coastal Plain in Brazil: Survival and Growth of Planted Woody Species. *Restoration Ecology* 14: 87-94. doi: 10.1111/j.1526-100X.2006.00108.x.

**Capítulo 2: Respostas fisiológicas de *Schinus terebinthifolia* Raddi
(Anacardiaceae) submetida à deficiência hídrica seguida por reidratação em
dois tipos de solos**

Physiological responses of *Schinus terebinthifolia* Raddi (Anacardiaceae) submitted to water deficiency followed by rehydration in two types of soils

Abstract - In plant metabolism water and soil nutrients are involved in all physiological processes and the ability to adapt to variations and changes in soil and water conditions is a key factor in resilience to environmental change. The adaptability of ecologically important trees is major importance in restoration initiatives, since these often take place in degraded habitats under sub-optimal environmental conditions. Here, we evaluate the physiological responses of a putatively resilient and widespread native Brazilian tree, *Schinus terebinthifolia* Raddi, under conditions of water deficit and cultivated in two different soil types. Our aim is to investigate the capacity of the species to withstand drought stress in soil types from ecologically distinct biomes (coastal forest and savanna). The study was carried out in greenhouses at the Rio Largo Campus of the federal University of Alagoas ($9^{\circ} 28' S$, $35^{\circ} 49' W$, at 127 meters altitude) with one year old plants. We adopted a completely randomized design in a 2×2 factorial scheme (two types of soil x water management), with five replications of each treatment. The analyses were standardized at two time intervals: (1) maximum stress and (2) rehydration. Data were collected on soil moisture, gas exchange, nitrogen metabolism compounds and compatible metabolites (proline, amino acids and ureides). Photochemical efficiency variables also measured, as well as microclimate and physicochemical data of the soil. Differences between the interactions of soil types versus water treatments were verified in C_i , WUE, iWUE, Ψ_{w12hs} , chlorophyll *a*, *b*, total and carotenoids, proline index, proteins, total ureides, alanthine, nitrate and also in the activity of the enzyme nitrate reductase on the day of greatest stress. Gaseous exchange gradually decreased as the available water in the soil was depleted. In the dry treatment of soil type 2, photosynthesis zeroed eleven days after irrigation was suspended, while soil 1 it occurred after seventeen days. Water treatment was more important than soil type for promoting physiological changes. Soil type influenced the time needed to reach peak stress and the recovery trajectory after water stress.

Key words: invasive species, water stress, aroeira.

Introducion

The coastal region of Brazil is mainly composed of vast sedimentary plains and contains various ecologically distinct vegetation formations, including natural fields, restingas (coastal forest), mangroves and other types of vegetation such as seasonal forest, deciduous forest and savannas (Law 11.428). Of these, the coastal restingas and the semi-arid savannas represent the greatest abiotic challenges to native plants due to their climate, salinity, strong winds, and in many cases, soil types that make it even more difficult to absorb water (Gomes et al., 2007; Costa et al., 2006).

The edaphic composition of this region includes sedimentary soils of the quaternary where the coastal restinga vegetation was established (GOMES et al., 1998). Some studies have classified restinga soils as quartz sands (Camaro et al., 1987; Oliveira et al., 1992; EMBRAPA, 2006), which normally have a sandy texture and are poor in nutrients (GOMES et al., 2007). Thus, the shrub-tree vegetation that is characteristic of the restinga is a result of the interaction of the regional pool of vegetation with the sandy soil and the climate of that region. Nevertheless, there is a considerable diversity of vegetation occurring in Brazilian restingas including several botanical families varying from herbaceous formations to trees that reach 20m in height.

The soil of the savannahs of eastern Brazil has a poorly developed surface mineral horizon, resulting from a reduced content of mineral or organic colloids or from the external climatic and vegetation conditions of this region (EMBRAPA, 2006). It has a clayey physical structure with the presence of boulders, being able to retain abundant water – though this is not necessarily available to the savannah vegetation due to both physical barriers and high osmotic potential. Soil types with higher water retention capacity or that are more fertile can influence the occurrence and frequency of some plant species (SILVA et al, 2009).

Water and soil nutrients are key factors in all plant growth processes (CABRAL et al, 2004), osmotic regulation, physiological regulation and photosynthetic processes (CARON et al, 2014). Thus, in order to thrive native tree species need to be able to adapt to changes in soil and water conditions caused by abiotic variations and anthropogenic degradation in natural environments. The ecological resilience and adaptability of ecologically important trees in their habitats is therefore of great interest to plant physiologists, ecologists and conservationists, since information on these

characteristics is of enormous value for sustainable management and the restoration programs of anthropically degraded areas (TRENBERTH et al. 2015).

One of the most resilient native Brazilian tree species is the Brazilian pepper tree, *Schinus terebinthifolia* Raddi, a wide-ranging and adaptable species that can be found in the Atlantic forest, restinga, and in transition areas between these two biomes (SILVA et al, 2008). Its great adaptability is reflected in the fact that it has become established in several extrazonal neotropical regions (JONES et al, 1997), and has even invaded several countries from outside the neotropics (EWE, 2008). The Brazilian pepper tree is highly cited in the literature as an appropriate species for the restoration of degraded environments in the tropics (LIMA et al, 2018) due to its seemingly high plasticity and ability to withstand periods of drought stress (VOLPATO; MARTINS, 2013). Here, we try to uncover the mechanisms underlying the high resilience of *S. terebinthifolia*, experimentally exposing individual plants to conditions of high water deficit while cultivated on two types of soil (restinga and savannah).

Material and methods

Experimental design and physical-chemical analysis of soil

One-year-old plants of *Schinus terebinthifolia* were transplanted to pots with 20 kg of soil collected in two sites. Soil 1 was collected at 9°47'52.7"S 35°51'57.0"W, in a restinga native area in the city of Barra de São Miguel. Soil 2 was collected at 9°23'28.1"S 37°12'38.9"W, in a caatinga area native in the city of Santana do Ipanema. Samples soils a sent for physical and chemical analysis in the Laboratory of Soil, Water and Plant of the Federal University of Alagoas (UFAL). From which sample, a portion of approximately 0.5 kg was separated and classified according to the Brazilian Soil Classification System (SBCS) using: (1) clay (densimeter); (2) fine and coarse sand (sieving); (3) clay dispersed in water (densimeter); (4) degree of flocculation (GF). As well as nutritional analyzes were performed using: (1) Mehlich-1 extractor; (2) 1.0 M KCl extractor; (3) Calcium acetate extractor at pH 7.0; (4) Welkley-Black method; (5) Base Saturation; (6) Saturation by aluminum.

Soil 1 from a restinga area and soil 2 from native savannah area of a semi-arid region was collected in the state of Alagoas. Restinga soil comes from the natural habitat of the Pepper tree while the savannah soil comes from a region (and biome) where this species does not naturally occur. The two types of soil have, as expected, very different physical and nutritional characteristics (Table 1. and Table 2.). Specifically, the soil of the semiarid region belongs to the loam clay sandy category according to SBCS's textural classification. Restinga soil is categorized as sandy, according to the same determinations and methodologies. Among the nutritional characteristics evaluated, the main difference between the two types of soil was the amount of potassium, which together with the other nutritional quantifications indicated that the savannah soil 2 was more fertile than the restinga soil.

Table 1. Physical soil analysis using (1) clay (densimeter); (2) fine and coarse sand (sieving); (3) clay dispersed in water (densimeter); (4) degree of flocculation (GF).

Determinations	Coast (%)	Semiarid (%)
Thick sand (g / kg)	940	410
Fine sand (g / kg)	20	180
Silt (g/kg)	20	190
Clay (g/kg)	20	220
Silt / clay ratio	1.00	0.86
Textural Classification (SBCS)	sand	Franco-Argiloarenosa

Biophysical data were monitored during two experimental periods. Period 1 lasted twelve days and included the maximum stress period and period 2 included the rehydration period (Figure 1.). The mean values of the air temperature, relative air humidity and the DPV are shown in the Figure 1. As can be seen, between 11:00 am and 14:00 pm there is an increase in DPV and air temperature, the opposite occurring with the relative humidity of the air, which at the same time has a considerable decrease, corresponding to the hottest and driest period of the day. These microclimatic variations have a great influence on the physiological responses of the plants to drought, as we will see later.

Table 2. Soil fertility analysis using (1) Mehlich-1 extractor; (2) 1.0 M KCl extractor; (3) Calcium acetate extractor at pH 7.0; (4) Welkley-Black method; (5) Base Saturation; (6) Saturation by aluminum.

Determinations	Coast (%)	Semiarid (%)
pH em água (1:2,5)	5.3	5.7
Na (mg/dm ³) ⁽¹⁾	15	20
P (mg/dm ³) ⁽¹⁾	3	14
K (mg/dm ³) ⁽¹⁾	15	90
Ca (cmol _c /dm ³) ⁽²⁾	1.08	3.48
Mg (cmol _c /dm ³) ⁽²⁾	0.87	1.67
Al (cmol _c /dm ³) ⁽²⁾	0.16	0.09
H + Al (cmol _c /dm ³) ⁽³⁾	3.48	2.28
CTCefetiva (cmol _c /dm ³)	2.22	5.56
CTC total (cmol _c /dm ³)	5.54	7.75
MO (g/kg) ⁽⁴⁾	18.5	11.8
V (%) ⁽⁵⁾	37	71
m (%) ⁽⁶⁾	7	2
Sat.de Ca (%)	19.5	44.9
Sat.de Mg (%)	15.7	21.5
Sat.de K (%)	0.7	3.0
Sat.de Na (%)	1.3	1.2

Seedlings of adequate phytosanitary status were standardized in a uniform phytosanitary state and transferred to a greenhouse in the Laboratory of Plant Physiology of the Center of Agricultural Sciences (CECA / UFAL). The plants were divided in four treatments, with five plants in each treatment: (a) irrigated plants in soil 1; (b) irrigated plants in soil 2; (c) plants with suspended irrigation in soil 1 and (d) plants with suspended irrigation in soil 2. When the photosynthesis of the plants with suspended irrigation reached zero, all plants were rehydrated. The treatments were

arranged in a completely randomized distribution. The analyses were performed at two time periods: (1) the period of maximum stress and (2) the period of rehydration.




Fig 1. Average hourly values of air temperature ($^{\circ}\text{C}$), relative humidity (%), radiation ($\mu\text{mol}/\text{m}^2/\text{sec}$) and vapor pressure deficit (Kpa), recorded every fifteen minutes and during the experiment in the greenhouse.

Determination of soil moisture

Measurements were taken on the first day of the experiment, on the day of maximum stress and after complete rehydration. Soil moisture was recorded with a portable humidity meter model SM200, coupled to a model HH2 sensor.

Determination of gas exchange; instantaneous water-use (iWUE) and intrinsic water-use efficiency (IWUE)

Gas exchange by fully expanded leaves was measured between 8:00 AM and 11:00 AM using an infrared gas analyzer (IRGA) model LI-6400 (Li-color, USA) equipped with a 6400-02B Red Blue artificial light source. The IRGA chamber (Li-cor 6400) was set to work with a flow of $1200 \mu\text{mol}/\text{s}^{-1}$. With IRGA, photosynthesis, stomatal conductance and transpiration data were collected. From these variables, the

values of instantaneous water-use efficiency (iWUE) and the intrinsic water-use efficiency (IWUE) were calculated as: iWUE ($\mu\text{mol CO}_2 \text{ mmol}^{-1} \text{ H}_2\text{O}$ = photosynthesis / transpiration) and IWUE ($\mu\text{mol CO}_2 \text{ mmol}^{-1} \text{ H}_2\text{O}$) = photosynthesis / stomatal conductance).

Determination of the water potential of the leaf

Leaf water potential was verified through the pressure chamber (SCHOLANDER et al. 1965) in the predawn and at midday, using the third leaf from the apex, showing good phytosanitary status, according to Turner's (1981) methodology.

Extraction of compounds

The extraction of the nitrogen compounds followed Bielecki and Turner (1966), where 500 mg of fresh material were macerated together with 5 ml of MCW solution (60% ml methanol, 25% ml chloroform, 15% ml H_2O). The material was centrifuged with 1 ml of chloroform and 1.5 ml of H_2O was added to each 4 ml of supernatant. After separation of the reaction phases, the supernatant (water soluble) was used to determine the compounds described below.

Quantification of nitrate

Nitrate was quantified following Cataldo (1975), where 0.2 ml of MCW extraction solution was added with 0.8 ml of 5% salicylic acid and concentrated sulfuric acid. After cooling the samples, 19 ml of 2N sodium hydroxide were added and read in a spectrophotometer at 410 nm.

Quantification of ureides (Allantoin and Alantoic Acid)

Was performed according to the method of Vogels and Van Der Drift (1970). A drop of phenylhydrazine was placed in test tubes with 250 μl of MCW extraction supernatant, 500 μl of distilled H_2O and 250 μl of 0.5 M NaOH and the assay was heated at 100 ° C in a 20% water bath. minutes and then cooled to room temperature. Once cooled, 250 μl of 0.65 N HCl is added and heated at 100 ° C again in a water bath for 4 minutes. After that time, the assay was cooled to room temperature and 250 μl of 0.4 M phosphate buffer pH 7.0 or 250 μl of 0.33% phenylhydrazine solution were added to this reaction and then, the assay was cooled in an ice bath for 5 minutes.

After cooling, 1.25 ml of pre-cooled, concentrated HCl and 250 μ l of 1.65% Potassium Ferrocyanide solution were added. After another ice bath, the assay was brought to room temperature and after 15 minutes a spectrophotometer reading was performed at 535 nm. The concentration of ureides was estimated based on the standard curve of the allantoin solution; compared to the standard proline curve (0 to 100 μ g mL⁻¹).

Quantification of the proline content

Was performed based on the methodology described by Bates (1973) where 50 μ L of the MCW extraction solution were collected and added to 2 mL of ninhydrin acid solution (1.25 g nihidrine, 30 mL glacial acetic acid , 20 mL of 6 M phosphoric acid) and 2 mL of glacial acetic acid in 15 mL microcentrifuge tubes. Samples were incubated at 100°C for about one hour, when the test reached purple color were placed on ice to paralyze the reaction. The samples were then added with 4 mL of toluene and constantly stirred for 20 seconds on the tube shaker for complete extraction of proline. The supernatant was used for 520 nm reading spectrophotometry and absorbance.

Quantification of soluble carbohydrates

Was performed according to Dubois (1956), where 50 μ L of the MCW extraction solution was added with 5% phenol and 250 μ L of concentrated sulfuric acid. Thereafter, the assay was shaken and cooled in an ice bath and then the spectrophotometer read at 490 nm. Carbohydrate concentration values were determined by the standard curve of glucose concentration (μ g).

Quantification of free amino acids

Using 1 ml of the supernatant (water soluble) from the MCW extraction was performed where 500 ul of citrate buffer + 200 ul of 5% of ninhydrin solution in methylglycol + 1 ml of KCN solution 0.0002M and this assay was heated at 100 ° C for 20 minutes and, after being cooled for 10 minutes, 1 ml of 60% ethyl alcohol was added according to Yemm and Cocking (1995), followed by spectrophotometer reading at 570 nm . The concentration of amino acids was estimated according to the standard curve of the Leucine solution.

Activity of the nitrate reductase enzyme

The enzymatic activity of nitrate reductase was determined following Stewart (1986), with 0.2 g leaf discs packed in 10 mL syringes and 5 mL of reaction medium (0.05 M K₂HPO₄, pH 7.5 0.1 M KNO₃, + 1% propanol). After insertion of the plunger, the syringe will be reversed to expel the air from the syringe. The technique of infiltration of the buffer during the determination of nitrate retune enzyme activity will be performed according to Majerowics et al. (2003). The exit of the syringe was closed with the finger and the plunger will be pulled 5 times in order to make the vacuum to totally submerge plant material. The syringes will be conditioned in the dark for 1 hour at room temperature (30 °C). After that time, the reaction was stopped and the samples used for the dosing of the enzymatic activity that was expressed in µmol of NO₂⁻ h⁻¹ g MF⁻¹.

Quantification of the spectrometric content of chlorophyll

The procedure was carried out by minimizing the luminosity to avoid degradation of the pigments following Arnon (1949) methodology. Samples of 50 mg of fresh leaf were minced and placed into the threaded test tube containing 5 ml of acetone (80%), for 48 hours. After this, determination was made spectrophotometrically with optical densities obtained in three wavelengths of (664, 665 and 750 nm). The method provides data on chlorophyll *a*, chlorophyll *b* and carotenoids in unit volume (µg/L).

Quantification of colorimetric chlorophyll content (SPAD index)

The index of chlorophyll content was determined by non-destructive method using chlorophyll meter (SPAD-502 - Minolta, Japan). Five readings were performed on randomly chosen leaflets from each plant, and with the mean we obtained the final value of the SPAD index.

Determination of chlorophyll a fluorescence, determination of the maximum and effective quantum yield of photosystem II

Chlorophyll a fluorescence was determined using a portable fluorometer (Handy PEA, Hansatech, Norfolk, UK) between 4:00 and 5:30 hours and between 11:00 and 12:30 hours on mature, healthy and fully expanded leaves. The leaves were subjected to 20 minutes of darkness by leaf tweezers and after that time, subjected to a pulse of

saturating light of 3000 ($\mu\text{mol m}^{-2}\text{s}^{-1}$) and the wavelength of 650 nm for 1 second. At midday the determination of the maximum and effective quantum yield of photosystem II with the same equipment was also performed.

Statistical analyzes

Significant differences between the two treatments and the two soils type were assessed with the Tukey Test (significant difference at 5% probability, $P<0.05$).

Results

Soil humidity

The control treatment plants of the two soil types were always maintained with the soil at field capacity. Soil moisture clearly reflected the water regimes. Soil 1 of the control treatment had values around 15 to 20% (Figure 3), values relatively lower than those found in pots with soil 2 in the same treatment (around 20 to 30%). A clear result of the granulometry of the two types of soil, since the sandy soil (soil 1) retains less water than the loam clay soil (soil 2).

The loam clay soil (soil 2), which theoretically retains more water than sandy soils, lost almost 20% of its moisture in 11 days (Figure 3), reaching the maximum stress for plant at this point. In contrast, the pots containing sandy soil (soil 1) maintained soil moisture (with a minimum of 7%) until the seventeenth day of the experiment (Figure 3), the day of maximum stress for plants in this soil (Figure 3).

We monitored physiological variables when photosynthesis reached values near or equal to zero in plants of the dry treatment of the two types of soil. Plants in soil 2 took ten to eleven days to zero their photosynthetic rates. Plants in soil 1 took seventeen days (Figure 3b). Thus, data collection at the point of maximum stress were carried out on different days, but were compared to each other, taking into account the stress level of the plant.

Photosynthesis was also used to determine the level of recovery after rehydration, that is, for dry treatment plants of the two types of soil that reached values close to zero and were rehydrated. Physiological monitoring data were therefore collected when photosynthesis reached values close to the values in control plants after rehydration. For plants in soil 2 this collection was performed on the seventeenth day and for plants in soil 1, on the nineteenth day of experiment (Figure 3b). For the control treatment plants the time of the data collections were carried out according to the corresponding soil type.





Fig 2. Temporal monitoring of soil moisture in both soil types (soil 1 and soil 2) and in both water treatments (control and drought). The arrow indicates the day of greatest stress (maximum stress).

Gas exchange

Rates of CO₂ assimilation reduced while the soil water was depleting in plants in the dry treatment. The peak of water stress for plants in soil type 2 occurred eleven days after the watering and after seventeen days for plants in soil type 1. At this time stomatal conductance was around zero. In this situation, stomata generally closed to avoid water loss by transpiration, which could consequently reduce photosynthesis, A/Ci, WUE and iWUE (Figure 4). Initially, plants cultivated in soil 1 under control conditions had higher photosynthesis rates. This is understandable considering that the restinga (Soil 1) is a natural habitat of the aroeira. However, by the eighth day the plants of this treatment matched their photosynthetic rates by dribbling the soil factor in their development (Figure 4a). Control and rehydrated plants maintained their photosynthetic rates between 13 and 17 µmol CO₂ m⁻² s⁻¹. The initial stomatal conductance, even under hydration conditions, was lower in soil 2. After 9 days this had equaled the rates verified in control plants in the soil 1. The internal CO₂ concentration of the control treatment remained around 400 µmol CO₂ m⁻² s⁻¹ throughout the experiment regardless of soil type (Figure 4g), and consequently it maintained a stable A/Ci ratio (Figure 4i).






Fig 3. Gaseous exchanges of young plants of *Schinus terebinthifolia* Raddi for 19 days. A Control 1 and 2 (A); A Dry only 1 and 2 (B); gs control soil 1 and 2 (C); gs dry soil 1 and 2 (D); E control solo 1 and 2 (E); E dry only 1 and 2 (F); Ci control soil 1 and 2 (G); Ci Dry 1 and 2 (H); A/Ci controls only 1 and 2 (I); A/Ci Dry 1 and 2 (J); EUA control soil 1 and 2 (K); EUA dry soil 1 and 2 (L); EIUA control only 1 and 2 (M); EIUA dry soil 1 and 2 (N).

In the dry treatments, the variables Ci and A/Ci had considerable peaks in the days preceding the maximum stress, regardless the type of soil, that is, when the stomata closed and photosynthesis reached levels close to zero. The same happened with effective water use efficiency (WUE) and the intrinsic water use efficiency (iWUE), that both had their highest values in the days before the maximum stress (Figure 4l and 4n.). This occurred when the capacity of photosynthesize and decrease of water loss by transpiration was optimized in dry treatment plants. WUE and iWUE values almost tripled after rehydration of soil 2, a much more pronounced effect than that seen in soil 1 (Figure 4l and 4n).

Variables affected by water stress and soil type

Differences in the interaction of soil types and water treatments were verified for Ci, WUE, iWUE, Ψ_{w12hs} , chlorophyll *a*, *b*, total and carotenoids, proline, protein, total ureides, and allantoin and nitrate content and also in the nitrate reductase activity at the maximum stress (Table 3). When we compare the "water regime" factor alone, almost all variables are higher in the control treatment reflecting the loss of water from the metabolism under drought conditions. The exceptions were the green index SPAD and chlorophyll *a/b* ratio that were not measurably affected by stress (Table 4). We also observed substances that increased with increasing water stress, such as proline, proteins, amino acids, total ureides and allantoin (Table 6).

However, isolating "soil type" we can see that within the same treatment there were some differences. For example, in the dry treatment of soil 2 Ci was 253% higher than in soil 1 of the same treatment , indicating that soil type directly influences the physiological responses of the plants (Table 4). The same pattern was observed with some variables in the control treatment; for example, chloro was lower in soil 1 than in soil 2 in control treatment plants (Table 5). In summary, few variables were directly influenced by the interaction between soil type and water treatment. Thus, we will focus on those variables sensitive to the set of changes: "soil + water regime". Similarly, rehydration appeared to be total and reversible in both types of soil.

Table 3. Summary of the analysis of variance of the interaction between two soil types (soil 1 and soli 2) and water regime (control and dry) with the gas exchange variables (A, gs, E, Ci, A/Ci, WUE, iWUE), photochemical (PSII, $F_v/F_{ma.m}$. and $F_v/F_{mp.m}$), water potential ($\Psi_{wa.m}$ and $\Psi_{wp.m}$), green intensity (chlorophyll a, chlorophyll b, carotenoids, total chlorophyll, ratio clor a/b and SPAD), water potential ($\Psi_{wa.m}$ and $\Psi_{wp.m}$), osmoregulators (proline, protein, amino acid, total sugars, total ureides, allantoin, allantoic acid) and compounds of nitrate metabolism (nitrate reductase, nitrate) in the four days of data collection.

SV	d.f.	Variables	F value		
			initial day	maximum stress	rehydration
S vs Wr	1	A	11.005 ^{ns}	2.87 ^{ns}	0.55 ^{ns}
Residue	16	gs	0.027 ^{ns}	2.16 ^{ns}	0.0001 ^{ns}
		E	2.83 ^{ns}	0.04 ^{ns}	0.021 ^{ns}
		Ci	1446.71 ^{ns}	742615.69*	239.43 ^{ns}
		A/Ci	0.00003 ^{ns}	0.0003 ^{ns}	0.00002 ^{ns}
		WUE	1.63 ^{ns}	1253.73**	0.21 ^{ns}
		iWUE	709.4 ^{ns}	405276.14**	78.76 ^{ns}
		$F_v/F_{0.12hs}$	0,0058 ^{ns}	0.0042 ^{ns}	0.0009 ^{ns}
		F_v/F_{m4hs}	0,00046 ^{ns}	0.000045 ^{ns}	0.00002 ^{ns}
		F_v/F_{m12hs}	0,00035 ^{ns}	0.00050 ^{ns}	0.0004 ^{ns}
		Ψ_w4hs	0,048 ^{ns}	0.038 ^{ns}	0.009 ^{ns}
		Ψ_w12hs	0,00045 ^{ns}	0.75*	0.00002 ^{ns}
		Clor a	0.00033 ^{ns}	1.14**	0.00002 ^{ns}
		Clor b	0.055 ^{ns}	0.19*	0.00002 ^{ns}
		Clor T	0.046 ^{ns}	0.11*	0.0001 ^{ns}
		Carotenoids	0.0049 ^{ns}	2.29**	0.08*
		Clor a/b	12.85 ^{ns}	0.11 ^{ns}	0.02 ^{ns}
		SPAD	58,82 ^{ns}	39.48 ^{ns}	29.04 ^{ns}
		Proline	0.0015 ^{ns}	4.61*	0.09 ^{ns}
		Protein	0.091 ^{ns}	8.24*	2.92 ^{ns}
		Amino acid	0.31**	0.80 ^{ns}	5.48**
		Total sugars	1.52**	0.18 ^{ns}	0.78*
		Total ureides	0.024 ^{ns}	256.68*	883.78**
		Allantoin	0.050 ^{ns}	247.45*	883.12**
		Allantoic acid	0.14 ^{ns}	0.08 ^{ns}	0.00012 ^{ns}
		Nitrate reductase	0.069**	0.16**	2.92*
		Nitrate	0.15 ^{ns}	0.40*	0.09 ^{ns}

SV: Sources of variation; soils (s); water regime (wr); d.f.:Degrees of freedom; * and ** significant at 5 and 1% probability, respectively; ns: not significant.

Table 4. Unfolding the interaction between two type soil (soil 1 and soil 2.) and water regime (control and dry) with the variables of gas exchanges (Ci, WUE, iWUE) and water potential (Ψ_w 12hs) in stress maximum.

Water regime	Ci		WUE		iWUE		Ψ_w 12hrs.	
	Soils							
	S1	S2	S1	S2	S1	S2	S1	S2
Control	395.35aA	357.0bA	4.01aA	5.65bA	42.6aA	59.5bA	-0.66aA	-0.40bA
Dry	289.01aB	1021.4aA	3.07aB	6.3aA	44.9aB	631.3aA	-0.94aB	-1.46aA
CV	6.37		9.06		6.72		-38.39	

Means followed by the same lowercase letter, in the columns, and upper case, in the lines, do not differ significantly from each other by the Tukey test (p<0,05).

Chlorophylls were one of the variables least affected by water stress or soil type. Chlorophyll a increased with the water stress in plants in soil 2, starting at 1.83 µg/L on the first day of experiment (data not shown) and rising to 2.13 µg/L on the day of maximum stress. Chlorophyll b and carotenoid content remained stable throughout the experiment (Table 5), contributing to the stability of total chlorophylls even under stress condition when grown in soil 2. In relation to Fv/Fm 4hs, there was a decrease at the maximum stress in the two soil types, which coincidentally had values of 0.788 µmol m⁻² s⁻¹. In the control treatments these values were 0.805 and 0.792 µmol m⁻² s⁻¹ (soil 1 and 2 respectively). At midday, even under the atmospheric influence, there were no significant changes in values (0.738 µmol m⁻² s⁻¹ for the dry treatment plants in soil 2, and 0.733 µmol m⁻² s⁻¹ for dry treatment plants in soil 1).

Table 5. Unfolding the interaction between two type soil (soil 1 and soil 2.) and water regime (control and dry) with the variables of chloroplastic pigments (Cloro a, b, total, carotenoids) in stress maximum.

Water regime	Clor a		Clor b		Clor T		Carot	
	Soils							
	S1	S2	S1	S2	S1	S2	S1	S2
Control	2.29bA	1.91aB	0.89aA	0.71aA	3.19aA	2.63aB	0.76aA	0.69aA
Dry	1.55aB	2.13aA	0.62bA	0.83aA	2.17bB	2.97aA	0.55bA	0.79aA
CV	13.02		24.12		11.40		16.08	

Means followed by the same lowercase letter, in the columns, and upper case, in the lines, do not differ significantly from each other by the Tukey test (p<0,05).

Based on predawn measurements of Fv/F0, water stress and atmospheric factors induced a decrease in the effective quantum efficiency of photosystem II, which reached levels of 0.457 and 0.420 in dry treatment plants in soil 2 and 1, respectively, at the time of maximum stress. Control treatment plants in both soil types had Fv/F0 of above 0.520.

As anticipated, nitrate reductase activity was affected by water stress. In dry treatment plants of soil 2, it reached values close to zero on the day of maximum stress (Table 6). This effect was milder in plant of the soil 1. The enzyme rapidly recovered after rehydration in plants of the two types of soil (Table 6). Control plants cultivated in soil 1 had higher values than control plants in soil 2. This result implies that the soil type influences nitrate assimilation. Consequently, it was verified that nitrate accumulates even after hydration. There was a negative effect of drought on the nitrate reductase activity and an accumulation of nitrate in leaves. In control plants, the accumulation of nitrate was not inversely proportional to enzyme activity (Table 6). Dry treatment plants accumulated higher levels of nitrate when grown on soil 2 (Figure 6b), where was enzymatic activity was lowest (Figure 5b).

Individuals decreased total sugar content at maximum stress, a probable consequence of the decrease in CO₂ assimilation verified in both types of soil. Conversely, there was an accumulation of total amino acids and proline at the maximum stress in both types of soil. Ureids and protein content were not significantly influenced by water regime or soil type (Table 6).

Table 6. Unfolding the interaction between two type soil (soil 1 and soil 2.) and water regime (control and dry) with the variables of osmoregulators (proline, protein, ureids, allantoin) and compounds of nitrogen metabolism (nitrate and RN) in stress maximum.

Water regime	Proline		protein		ureids		allantoin		nitrate		RN	
	Soils											
	S1	S2	S1	S2	S1	S2	S1	S2	S1	S2	S1	S2
Control	1.18b	0.88b	3.82bA	4.28aA	51.72aB	64.97bA	37.87bB	48.87aA	0.56aA	0.63bA	1.14bB	0.73aA
Dry	14.74a	16.37a	5.43aA	5.24aA	66.67aA	65.59aA	53.72aA	50.65aA	0.63aB	1.27aA	0.05aA	0.008bA
CV	11.41		25.05		8.96		12.77		18.92		11.16	

Means followed by the same lowercase letter, in the columns, and upper case, in the lines, do not differ significantly from each other by the Tukey test ($p<0,05$).

Discussion

Water stress conditions appear to cause more severe consequences for *S. terebinthifolia* when grown in clay soil (soil 2). In this soil the gas exchange reached zero in less time than in the sandy soil (soil 1), corroborating previous studies (AI et al, 2017). In contrast, the recovery of CO₂ assimilation after rehydration was considerably faster in soil 1. Sandy soils are the natural habitat of this species, and stress and recovery would seem to be related to a combination of water availability of the soil and the water absorption capacity of the plant (AI et al. 2017).

Previous studies have shown that the rates of liquid photosynthetic, stomatal conductance and photochemical efficiency (Fv/Fm) may differ significantly between soil types (AI et al, 2017). Soil 2 has better water retention, higher surface area of particles, higher amount of organic matter and greater presence of clay. It also has the greatest capacity to retain water, the scarcest for the plant to absorb, which involves physical and chemical issues (AI et al. 2017). The experimental plants tolerated a greater water shortage in the sandy soil, probably because greater water availability between the particles and dissolved solids (SAAD et al, 2009). In addition, when the water in the clay soil begins to evaporate, clefts form further increasing contact with the atmosphere leading to higher evaporation and damaging the roots by exposing them to the atmosphere.

The key value in our study was the level of soil moisture that could bring about changes in the species' metabolism. Indeed, Lopéz (2016) suggest that the effects of shade and soil type are less important than the effects of water deficit. Nevertheless, the level of water stress in the current study was insufficient to damage PSII functioning – corroborating the work of Bosque et al. (2018) on *Lithraea Caustica* (liters), an Anacardiaceae. This plant evolved in a typical mediterranean climate, characterized by summers of low soil water availability, high solar radiation and at high temperatures. This species probably has the capacity to minimize or delay the inactivation of photosystem proteins through a mechanism of more efficient dissipation of excess energy, as observed in cultivated plants (ZEGADA-LIZARAZU et al., 2015)..

Of the variables significantly influenced by water stress and soil type, Ci (along with WUE and iWUE) showed the greatest difference between dry soil 2 plants, probably due to the rapid stomatal closure response that occurred in dry plants during the days which preceded the maximum stress (CHIRICO et al. 2017; PEÑA-ROJAS et al., 2018). When the volume of transpired water exceeds the volume of water

absorbed, the plant enters in a state of deficit and dehydrates, as observed in *Handroanthus impetiginosus* (Pessoa et al. 2017). The same process was observed in drought exposed *Myracrodruon urundeuva* (Anacardeaceae) and representatives of the family Arecaceae and Meliaceae (OLIVEIRA et al, 2002; MARIANO et al. 2009; MARENCO et al. 2014; ALBUQUERQUE et al., 2013; COSTA, et al 2015).

Proline accumulation has been reported to occur after certain external disturbances (VERBRUGGEN et al. 2008; HUANG et al. 2018). In our study, the increase of proline was considered as a sign of drought stress (SCHIMPL et al. 2019), together with the degradation of total soluble sugars (LUO and LUO, 2017) (SCHIMPL et al. 2019). As this increase was not verified among the same treatments in different soil types, we conclude that soil type did not contribute to stress in this species. Only lack of water affected the osmoprotective mechanism.

Ureides have also been implicated in resistance to water stress (KING et al. 2005). Their content was higher in dry treatment plants when compared to irrigated treatment in each soil type. The use of urea-derived compounds in tree species is very scarce, and is restricted to legumes or cultivated species (QUILES et al. 2019). Concentrations of these compounds are usually increased in stress conditions, corroborating our results (LAMBERT et al. 2016; LAMBERT et al. 2017; KABBADJ et al. 2017).

Nitrate is toxic in large quantities and was observed to accumulate in aroeira leaves under water stress condition (SHANER et al. 1967). This is because water stress inhibits the enzyme nitrate reductase in both soil types (CASTRO et al. 2007; OLIVEIRA et al. 2011; OMENA-GARCIA et al. 2015). As no leaf necrosis caused by nitrate toxicity was observed in this study, following BENINNI (2002) we believe that nitrate probably influences the osmotic adjustment of the cells in non-toxic amounts. We tried to relate this to the value of the water potential of the leaf and were able to verify that the soil interaction versus water treatment was significant at noon and stressed plants had lower values than control plants. Dry treatment plants in soil 2 decreased their soil water potential more than those in soil 1, probably due to the difficulty of absorbing water from the clay soil. This lead to changes in the osmotic adjustment of leaves (FERNANDES et al. 2018), photosynthetic responses, and photochemical responses, mainly during the day (SELLIN et al. 2014; SILVA-BRANCO et al. 2017). It is important to note that other factors, such as water, atmospheric

variables, and soil, are extremely important to evaluate in this context (SILVA et al, 2006; TROVÃO et al. 2004; ALBUQUERQUE et al. 2013).

Plants grown in soil 2 increased their pigment content, including Clor a and Clor T. The opposite pattern was observed in plants cultivated in soil 1, which decreased their chlorophyll content under stress. This corroborates several previous studies in other species (REGO, et al. 2006; NOGUEIRA et al 2001; BAYOUMI et al. 2015; SILVA et al. 2003) *S. terebinthifolia* is halophytic and maintains chlorophyll content even under severe stress conditions (BRITO et al. 2018).

In conclusion, water stress most strongly stimulated the observed physiological changes with a more limited influence of soil type mediated through the time in which the species takes to reach its peak of stress and recovery. When hydrated, soil type did not appear to limit the growth and functioning of the plant.

References

- AI, Zemin et al. Soil nutrients influence the photosynthesis and biomass in invasive *Panicum virgatum* on the Loess Plateau in China. *Plant And Soil*, [s.l.], v. 418, n. 1-2, p.153-164, 22 maio 2017. Springer Nature. <http://dx.doi.org/10.1007/s11104-017-3286-x>.
- Albuquerque MPF, Moraes FKC, Santos RIN, Castro GLS, Ramos EMLS, Pinheiro HÁ (2013) Ecofisiologia de plantas jovens de mogno-africano submetidas a deficit hídrico e reidratação. *Pesq. agropec. bras.*, v.48, n.1, p.9-16. doi: 10.1590/S0100-204X2013000100002
- Allen RG, Pereira LS, Raes D, Smith M (1998) Crop evapotranspiration - Guidelines for computing crop water requirements. FAO Irrigation and Drainage Papers.56:326.
- Arnon DI (1949) Copper enzymes in isolated chloroplasts. Polyphenoloxidase in *Beta vulgaris*. *Plant physiology*. 24:1-15.
- Bates LS, Waldren RP, Teare ID (1973) Rapid determination of free proline for water-stress studies. *Plant Soil* 39:205–207. doi: 10.1007/BF00018060
- Bayoumi TY, Amal M, EL-Mageed A; Ibrahim ES, Mahmoud SA, El-Demardash IS, Abdel-Raheem A (2015) The Impact of Drought Stress on some Morpho-Physiological Traits and RAPD Markers in Wheat Genotypes. *Journal of Plant Production Sciences*. 4:27-37.
- Beninni ERY, Takahashi HW, Neves CSVJ, Fonseca, ICB (2002) Teor de nitrato em alface cultivada em sistemas hidropônico e convencional. *Horticultura Brasileira*. 20:183-186.
- Bieleski RL, Turner NA (1966) Analytical Separation and estimation of amino acids in crude plant extracts by thin-layer electrophoresis and chromatography. *Biochemistry* 17:278-293. doi: 10.1016/0003-2697(66)90206-5
- Cabral EC, Barbosa DCA, Simabukuro EA (2004) Crescimento de plantas jovens de *Tabebuia aurea* (Manso) Benth. & Hook. f. ex S. Moore submetidas a estresse hídrico. *Acta bot. bras.* 18: 241-251.

Caron BO, Perrando ER, Schmidt D, Manfron PA, Behling A, Elli EF, Eloy E (2014) Relações fisiológicas em mudas de pata-de-vaca (*Bauhinia forficata* Link). Revista Brasileira Plantas Medicinais. 16:196-201.

Castro DS, Santos AO, Lobato AKS, Gouvea DDS, Neto CFO, Cunha RLM, Costa RCL (2007) Concentrações de Prolina e Carboidratos Solúveis Totais em Folhas Teca (*Tectona grandis* L.f) Submetida ao Estresse Hídrico. Revista Brasileira de Biociências. 5:921-923.

Cataldo DA, Harron M, Schradev LE (1975) Rapid colorimetric determination of nitrate in plant tissue by nitration of salicylic acid. Communications in Soil Science and Plant Analysis, 6:71 – 80.

Cesário LF, Gaglianone MC (2008) Biologia floral e fenologia reprodutiva de *Schinus terebinthifolia* Raddi (Anacardiaceae) em Restinga do Norte Fluminense. Acta bot. Bras. 22:828-833.

Chirino, Esteban et al. Morpho-functional traits and plant response to drought conditions in seedlings of six native species of Ecuadorian Ecosystems. Flora, [s.l.], v. 233, p.58-67, ago. 2017. Elsevier BV. <http://dx.doi.org/10.1016/j.flora.2017.05.012>.

Costa AS, Freire ALO, Bakke IA, Pereira FHF (2015) Respostas fisiológicas e bioquímicas de plantas de aroeira (*Myracrodruon urundeuva* allemão) ao déficit hídrico e posterior recuperação. Irriga, Botucatu. 20:705-717.

Dubois M, Gilles K A, Hamilton JK, Rebers PA, Fred S (1956) Colorimetric method for determination of sugars and related substances Anal. Chem. 28:3. 350-356. DOI 10.1021/ac60111a017

Ewe, SML, STERNBERG LSL (2007) Water Uptake Patterns of an Invasive Exotic Plant in Coastal Saline Habitats. Journal Of Coastal Research. 231:255-264. Coastal Education and Research Foundation. <http://dx.doi.org/10.2112/1551-5036>.

Fernandes RDM, Cuevas MV, Diaz-Espejo A, Hernandez-Santana V (2018) Effects of water stress on fruit growth and water relations between fruits and leaves in a hedgerow olive orchard. Agricultural Water Management 210:32–40. doi: 10.1016/j.agwat.2018.07.028

Gonçalves JFC, Carlos Eduardo Moura da Silva e Diogo Gato Guimarães (2009) Fotossíntese e potencial hídrico foliar de plantas jovens de andiroba submetidas à deficiência hídrica e à reidratação. *Pesq. agropec. bras.*, Brasília. 44:8-14.

JONES, D.. (1997) Biology of Brazilian pepper management plan for Florida. Florida, Florida Exotic Pest Plant Council.

Kabbadj, Abla et al.. (2017) Physiological and biochemical responses involved in water deficit tolerance of nitrogen-fixing *Vicia faba*. *Plos One*. 12:1-12. <http://dx.doi.org/10.1371/journal.pone.0190284>.

King CA, Purcell LC (2005) Inhibition of N₂ fixation in soybean is associated with elevated ureides and amino acids. *Plant Physiol*. 137:1389-1396

Lambert, R., Cabello-Diaz, J.M., Quiles, F.A., Piedras, P., 2016. Identification of nucleases related to nutrient mobilization in senescing cotyledons from French bean. *ActaPhysiol. Plant.* 38, 11.

Lambert, R., Quiles, F.A., Gálvez-Valdivieso, G., Piedras, P., 2017. Nucleases activities during French bean leaf aging and dark-induced senescence. *J. Plant Physiol*. 218,235–242.

Lima AD, Bezerra FMS, Neves ALR, Sousa CHC, Lacerda CF, Bezerra AME (2018) Response of four woody species to salinity and water deficit in initial growth phase. *Revista Brasileira de Engenharia Agrícola e Ambiental*. 22:753-757. FapUNIFESP (SciELO). <http://dx.doi.org/10.1590/1807-1929/agriambi.v22n11p753-757>.

Luo, Z. B.; Luo, J. Uncovering the physiological mechanisms that allow nitrogen availability to affect drought acclimation in *Catalpa bungei*. *Tree Physiology*, v. 37, p. 1453–1456, 2017.

Majerowicz, N. et al (2003) *Fisiologia Vegetal - Curso Prático*, Âmbito Cultural Edições Ltda., RJ.

Maggio A, Miyazaki S, Veronese P, Fujita T, Ibeas JI, Damsz B, Narasimhan ML, Hasegawa PM, Joly RJ, Bressan RA (2002) Does proline accumulation play an active role in stress-induced growth reduction? *Plant J*. 31:699–712.

Marenco RA, Antezana-Vera SA, Gouvêa PRS, Camargo MAB, Oliveira MFO, Santos JKS (2014) Fisiologia de espécies florestais da Amazônia: fotossíntese, respiração e relações hídricas. Rev. Ceres, Viçosa. 61:786-789. doi: dx.doi.org/10.1590/0034-737X201461000004

Mariano KRS, Barreto LS, Silva AHB, Neiva GKP, Amorim S (2009) Fotossíntese E tolerância protoplasmática foliar em *Myracrodroon urundeava* FR. ALL. submetida ao déficit hídrico. Revista Caatinga [en linea] doi: 0100-316X

Nogueira RJNC, Moraes JAPV, Burit HÁ, Neto EB (2000) Alterações na resistência à difusão de vapor das folhas e relações hídricas em aceroleiras submetidas a déficit de água. R. Bras. Fisiol. Veg. 13:75-87

Oliveira LMB, Bevilaqua CML, Macedo ITF, Moraes SM, Machado LKA, Campello CC, Mesquita MA (2011) Effects of *Myracrodroon urundeava* extracts on egg hatching and larval exsheathment of *Haemonchus contortus*. Parasitology Research. 109:893-898. Springer Nature. <http://dx.doi.org/10.1007/s00436-011-2331-6>.

Omena-Garcia RP, Justino GC, Araújo VBF, Souza LAG, Camargos LS, Gonçalves JFC (2014) Mineral Nitrogen Associated Changes in Growth and Xylem-N Compounds in Amazonian Legume Tree. Journal of Plant Nutrition. doi: 10.1080/01904167.2014.957389

Peña-rojas, Karen et al. Respuestas morfo-fisiológicas de plantas de *Lithraea caustica* (Anacardiaceae) sometidas a restricción hídrica controlada. Bosque (valdivia), [s.l.], v. 39, n. 1, p.27-36, 2018. SciELO Comision Nacional de Investigacion Cientifica Y Tecnologica (CONICYT). <http://dx.doi.org/10.4067/s0717-92002018000100027>.

Pessoa, Jéssica Lima; FREIRE, Antonio Lucineudo Oliveira; COSTA, Amanda Silva. Trocas gasosas de plantas de *Handroanthus impetiginosus* (Mart. ex DC) Mattos submetidas ao déficit hídrico e posterior reidratação. Revista de Ciências Agroveterinárias, [s.l.], v. 16, n. 3, p.269-276, 23 out. 2017. Universidade do Estado de Santa Catarina. <http://dx.doi.org/10.5965/223811711632017269>.

Quiles FA, Galvez-Valdivieso G, Guerrero-Casado J, Pineda M, Piedras P (2019) Relationship between ureidic/amidic metabolism and antioxidant enzymatic activities

in legume seedlings. *Plant Physiology And Biochemistry*. 138:1-8. Elsevier BV. <http://dx.doi.org/10.1016/j.plaphy.2019.02.016>.

Rego GM, Possamai, E (2011) Efeito do Sombreamento sobre o Teor de Clorofila e Crescimento Inicial do Jequitibá-rosa. *Pesquisa Florestal Brasileira*. 53:179. ISSN 1983-2605

Schimpl, Flávia Camila et al. Physiological responses of young Brazil nut (*Bertholletia excelsa*) plants to drought stress and subsequent rewetting. *Flora*, [s.l.], v. 252, p.10-17, mar. 2019. Elsevier BV. <http://dx.doi.org/10.1016/j.flora.2019.02.001>.

Saad JCC, Lopes JLW, Santos TA (2009) Manejo hídrico em viveiro e uso de hidrogel na sobrevivência pós-plantio de *Eucalyptus urograndis* em dois solos diferentes. *Eng. Agríc.*, Jaboticabal. 29:404-411.

Salla L, Rodrigues JC, Marenco RA (2007) Teores de clorofila em árvores tropicais determinados com o SPAD-502. *Revista Brasileira de Biociências*. 5:159-161

Scholander PF, Hammel HT, Hemingsen EA, Bradstreet ED (1965) Hydrostatic pressure and osmotic potentials in leaves of mangroves and some other plants. *Proc Natl Acad Sci USA*. 51:119–125

Sellin A, Niglas A, Õunapuu-Pikas E, Kupper P (2014) Rapid and long-term effects of water deficit on gas exchange and hydraulic conductance of silver birch trees grown under varying atmospheric humidity. *BMC Plant Biology*.14-72.

Shaner DL, Boyer JS.. Nitrate Reductase Activity in Maize (*Zea mays L.*) Leaves: I. Regulation by Nitrate Flux. *Plant Physiology*. American Society of Plant Biologists (ASPB). 58:499-504. <http://dx.doi.org/10.1104/pp.58.4.499>.

Silva MAV, Nogueira RJMC, Oliveira AFM, Santos VF (2008) Resposta estomática e produção de matéria seca em plantas jovens de aroeira submetidas a diferentes regimes hídricos. *Revista Árvore*. 32:2, p.335-344

Silva MR, Klar AE, Passos JR (2004) efeitos do manejo hídrico e da aplicação de potássio nas características morfofisiológicas de mudas de *Eucalyptus Grandis* W. (Hill ex. Maiden) 9:1. p. 31-40. doi:10.15809/irriga.2004v09n1p31-40

Silva, E. C. A. Respostas fisiológicas bioquímicas e enzimáticas em mudas de Moringa oleifera Lam. submetidas a estresses abióticos. 2013. Dissertação (Mestrado em Ciências Florestais). Universidade Federal Rural de Pernambuco, Recife, PE, 2013.

Silva-Pinheiro J, Lins L, Souza FC, Silva CEM, Moura FBP, Endres L, Justino GC (2016) Drought-stress tolerance in tree semi-arid species used to recover logged area. *Braz J Bot* 39(4):1031–1038. doi: 10.1007/s40415-016-0309-4

Souza BD, Meiado MV, Rodrigues BM; Santos MG (2010) Water relations and chlorophyll fluorescence responses of two leguminous trees from the Caatinga to different watering regimes. *Acta Physiol Plant* 32:235–244. doi: 10.1007/s11738-009-0394-0

Stewart GR, Popp M, Holzapfel I, Stewart JA, Dickie-eskew A(1986) Localization of nitrate reduction in ferns and its relationship to environment and physiological characteristics. *New Phytologist*. 104:373-384. Wiley. <http://dx.doi.org/10.1111/j.1469-8137.1986.tb02905.x>.

Trenberth KE, Fasullo JT, Shepherd TG (2015) Attribution of climate extreme events. *Nature Climate Change*. 5:725-730. Springer Nature. <http://dx.doi.org/10.1038/nclimate2657>.

Trovão DMBM, Fernandes PD, Andrade LA, Neto JD, Oliveira AB, Queiroz JA (2004) Avaliação do potencial hídrico de espécies da Caatinga sob diferentes níveis de umidade no solo. *Revista de Biologia e Ciências da Terra [on line]*. 4:1519-5228

Turner, NC (1981) Techniques and experimental approaches for the measurement of plant water status. *Plant and soil*, 58:339-366

Verbruggen N; Hermans C (2008) Proline accumulation in plants: a review. *Amino Acids* (2008) 35:753–759. doi: 10.1007/s00726-008-0061-6

Vogels GD, Drift C (1970) Differential analyses of glyoxylate derivates. *Anal. Biochem.* 33:143-157.

Volpato GH, Martins SV (2013) The bird community in naturally regenerating *Myracrodruon urundeuva* (Anacardiaceae) forest in Southeastern Brazil. Rev. Biol. Trop. 61:4 - 1585-1595

WANG, M. et al. The critical role of potassium in plant stress response. International Journal of Molecular Sciences, v. 14(4), p. 7370-7390, 2013.

Yemm EW, Cocking EC (1955). The determination of amino acids with ninhydrin. Analyst. 80:209-213.

Zegada-lizarazu W, Luna DF, Monti A (2015) Photosynthetic acclimation of sweet sorghum under progressive water stress. Industrial Crops And Products. 66: 216-219. Elsevier BV. <http://dx.doi.org/10.1016/j.indcrop.2014.12.045>.